

# UAW-GM Cohort Study

With additional model terms; exposure lagged 21 years

*November 12, 2019*

## Introduction

In previous survival analyses, hazard ratios associated with exposure to the three metalworking fluid types were estimated simultaneously in the same Cox proportional hazards model. There was a concern that those estimates may have been biased or misleading, as those models assumed independent covariate (statistical) effects e.g. that the effect of exposure to straight metalworking fluids was constant across levels of exposure to other metalworking fluid types. One way we attempted to address this concern was to fit independent models for each exposure-outcome pair of interest with one additional term to capture changes in the log-hazard associated with the combination of zero exposure to one metalworking fluid type and nonzero exposure to some other metalworking fluid type.

## Model

Let  $\mathbf{W}(t)$  represent the vector of potential confounders at time  $t$ . Let  $X_{St}$ ,  $X_{So}(t)$ , and  $X_{Sy}(t)$  represent cumulative exposure to straight, soluble, and synthetic metalworking fluids, respectively at time  $t$ . If we were interested in straight metalworking fluids, then the log hazard for the outcome could be modeled:

$$\begin{aligned} \log [h(t | \mathbf{W}(t) = \mathbf{w}(t), \mathbf{X}(t) = \mathbf{x}(t), \hat{\beta})] = & \log [h_0(t)] \\ & + \hat{\beta}_1 \mathbb{1} [x_{St}(t) \in (0, St_{low}]] \\ & + \hat{\beta}_2 \mathbb{1} [x_{St}(t) \in (St_{low}, St_{mid}]] \\ & + \hat{\beta}_3 \mathbb{1} [x_{St}(t) \in (St_{mid}, St_{high}]] \\ & + \hat{\beta}_4 \mathbb{1} [x_{So}(t) \in (0, So_{low}]] \\ & + \hat{\beta}_5 \mathbb{1} [x_{So}(t) \in (So_{low}, So_{mid}]] \\ & + \hat{\beta}_6 \mathbb{1} [x_{So}(t) \in (So_{mid}, So_{high}]] \\ & + \hat{\beta}_7 \mathbb{1} [x_{Sy}(t) \in (0, Sy_{low}]] \\ & + \hat{\beta}_8 \mathbb{1} [x_{Sy}(t) \in (Sy_{low}, Sy_{mid}]] \\ & + \hat{\beta}_9 \mathbb{1} [x_{Sy}(t) \in (Sy_{mid}, Sy_{high}]] \\ & + \hat{\beta}_{10} \mathbb{1} [x_{St}(t) = 0] \mathbb{1} [x_{So}(t) + x_{Sy}(t) > 0] \\ & + \hat{\beta}_{11} w_1(t) + \hat{\beta}_{12} w_2(t) + \hat{\beta}_{13} w_3(t) + \dots \end{aligned}$$

where  $St_{low}$ ,  $So_{low}$ , and  $Sy_{low}$  represent the upper boundaries of low exposure to the three types of metalworking fluids among cases at time of death, respectively;  $St_{mid}$ ,  $So_{mid}$ , and  $Sy_{mid}$  represent the upper boundaries of moderate exposure among cases at time of death; and  $St_{high}$ ,  $So_{high}$ , and  $Sy_{high}$  represent maximum exposure among cases at time of death. If we were interested in exposure to soluble metalworking fluids, we would replace the 10<sup>th</sup> covariate with  $\mathbb{1} [x_{So}(t) = 0] \mathbb{1} [x_{St}(t) + x_{Sy}(t) > 0]$ . For synthetic, we would replace with  $\mathbb{1} [x_{Sy}(t) = 0] \mathbb{1} [x_{St}(t) + x_{So}(t) > 0]$ . Note that if all three terms were included simultaneously, we would have a rank-deficient model matrix.

## Results

Table 1: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **straight** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI	
Laryngeal cancer (73 cases)						
0	mg/m <sup>3</sup> .years	40				
> 0 to 0.5	mg/m <sup>3</sup> .years	17	1.10	0.78	(0.55, 2.18)	
> 0.5	mg/m <sup>3</sup> .years	16	0.80	0.52	(0.41, 1.58)	
Trend				0.26		
Lung cancer (1894 cases)						
0	mg/m <sup>3</sup> .years	967				
> 0 to 0.3	mg/m <sup>3</sup> .years	309	1.06	0.42	(0.91, 1.24)	
> 0.3 to 1.6	mg/m <sup>3</sup> .years	309	0.95	0.48	(0.81, 1.10)	
> 1.6	mg/m <sup>3</sup> .years	309	0.90	0.15	(0.78, 1.04)	
Trend				0.25		
Esophageal cancer (176 cases)						
0	mg/m <sup>3</sup> .years	83				
> 0 to 0.4	mg/m <sup>3</sup> .years	31	1.16	0.54	(0.71, 1.89)	
> 0.4 to 2.1	mg/m <sup>3</sup> .years	32	1.13	0.62	(0.70, 1.84)	
> 2.1	mg/m <sup>3</sup> .years	30	1.15	0.57	(0.72, 1.84)	
Trend				0.57		
Stomach cancer (194 cases)						
0	mg/m <sup>3</sup> .years	104				
> 0 to 0.3	mg/m <sup>3</sup> .years	30	1.02	0.95	(0.63, 1.65)	
> 0.3 to 2.9	mg/m <sup>3</sup> .years	30	0.74	0.22	(0.45, 1.20)	
> 2.9	mg/m <sup>3</sup> .years	30	1.68	0.03	(1.05, 2.66)	*
Trend				0.06		.
Colon cancer (407 cases)						
0	mg/m <sup>3</sup> .years	211				
> 0 to 0.5	mg/m <sup>3</sup> .years	66	0.89	0.49	(0.65, 1.23)	
> 0.5 to 2.1	mg/m <sup>3</sup> .years	65	0.91	0.56	(0.65, 1.26)	
> 2.1	mg/m <sup>3</sup> .years	65	0.91	0.54	(0.67, 1.24)	
Trend				0.60		
Rectal cancer (83 cases)						
0	mg/m <sup>3</sup> .years	44				
> 0 to 1	mg/m <sup>3</sup> .years	20	0.79	0.49	(0.40, 1.54)	
> 1	mg/m <sup>3</sup> .years	19	0.88	0.71	(0.46, 1.70)	
Trend				0.84		
Bladder cancer (138 cases)						
0	mg/m <sup>3</sup> .years	73				
> 0 to 0.3	mg/m <sup>3</sup> .years	20	0.97	0.90	(0.54, 1.71)	
> 0.3 to 1.8	mg/m <sup>3</sup> .years	22	0.72	0.24	(0.41, 1.26)	
> 1.8	mg/m <sup>3</sup> .years	23	0.75	0.27	(0.44, 1.25)	
Trend				0.41		
Liver cancer (123 cases)						
0	mg/m <sup>3</sup> .years	55				
> 0 to 0.5	mg/m <sup>3</sup> .years	22	1.01	0.98	(0.57, 1.80)	

Table 1: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **straight** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI	
> 0.5 to 1.6	mg/m <sup>3</sup> .years	23	1.69	0.07	(0.96, 2.94)	.
> 1.6	mg/m <sup>3</sup> .years	23	1.17	0.57	(0.68, 2.01)	
Trend				0.98		
Pancreatic cancer (315 cases)						
0	mg/m <sup>3</sup> .years	154				
> 0 to 0.3	mg/m <sup>3</sup> .years	54	1.19	0.36	(0.83, 1.70)	
> 0.3 to 1.1	mg/m <sup>3</sup> .years	53	1.10	0.61	(0.76, 1.60)	
> 1.1	mg/m <sup>3</sup> .years	54	0.82	0.27	(0.58, 1.16)	
Trend				0.21		
Skin cancer (69 cases)						
0	mg/m <sup>3</sup> .years	32				
> 0 to 0.9	mg/m <sup>3</sup> .years	18	1.45	0.28	(0.73, 2.88)	
> 0.9	mg/m <sup>3</sup> .years	19	1.54	0.21	(0.78, 3.04)	
Trend				0.53		
Prostate cancer (418 cases)						
0	mg/m <sup>3</sup> .years	192				
> 0 to 0.5	mg/m <sup>3</sup> .years	76	1.10	0.56	(0.81, 1.49)	
> 0.5 to 2	mg/m <sup>3</sup> .years	75	1.08	0.65	(0.78, 1.48)	
> 2	mg/m <sup>3</sup> .years	75	1.01	0.93	(0.75, 1.37)	
Trend				0.78		
Brain and nervous system cancers (128 cases)						
0	mg/m <sup>3</sup> .years	74				
> 0 to 1	mg/m <sup>3</sup> .years	27	0.69	0.20	(0.40, 1.21)	
> 1	mg/m <sup>3</sup> .years	27	0.80	0.42	(0.47, 1.36)	
Trend				0.76		
Leukemia (200 cases)						
0	mg/m <sup>3</sup> .years	100				
> 0 to 0.3	mg/m <sup>3</sup> .years	34	1.10	0.68	(0.69, 1.75)	
> 0.3 to 2.3	mg/m <sup>3</sup> .years	33	0.79	0.33	(0.49, 1.27)	
> 2.3	mg/m <sup>3</sup> .years	33	0.95	0.81	(0.60, 1.49)	
Trend				0.86		
Breast cancer (76 cases)						
0	mg/m <sup>3</sup> .years	43				
> 0 to 0.7	mg/m <sup>3</sup> .years	16	1.02	0.95	(0.52, 2.03)	
> 0.7	mg/m <sup>3</sup> .years	17	1.83	0.07	(0.96, 3.50)	.
Trend				0.02		*

Table 2: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **soluble** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

	Number of cases	HR	<i>p</i>	95% CI
Laryngeal cancer (73 cases)				

Table 2: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **soluble** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI	
0 to 5	mg/m <sup>3</sup> .years	41				
> 5 to 17.9	mg/m <sup>3</sup> .years	16	1.07	0.85	(0.54, 2.11)	
> 17.9	mg/m <sup>3</sup> .years	16	1.65	0.18	(0.79, 3.42)	
Trend				0.04		*
Lung cancer (1894 cases)						
0 to 5	mg/m <sup>3</sup> .years	1083				
> 5 to 9.8	mg/m <sup>3</sup> .years	271	0.92	0.31	(0.79, 1.08)	
> 9.8 to 19.2	mg/m <sup>3</sup> .years	270	1.01	0.88	(0.87, 1.18)	
> 19.2	mg/m <sup>3</sup> .years	270	1.04	0.63	(0.89, 1.22)	
Trend				0.37		
Esophageal cancer (176 cases)						
0 to 5	mg/m <sup>3</sup> .years	93				
> 5 to 9.5	mg/m <sup>3</sup> .years	27	0.94	0.81	(0.57, 1.55)	
> 9.5 to 16.2	mg/m <sup>3</sup> .years	28	1.36	0.22	(0.83, 2.24)	
> 16.2	mg/m <sup>3</sup> .years	28	0.89	0.67	(0.53, 1.50)	
Trend				0.52		
Stomach cancer (194 cases)						
0 to 5	mg/m <sup>3</sup> .years	114				
> 5 to 8.3	mg/m <sup>3</sup> .years	27	1.31	0.27	(0.81, 2.09)	
> 8.3 to 17.8	mg/m <sup>3</sup> .years	26	0.83	0.44	(0.51, 1.35)	
> 17.8	mg/m <sup>3</sup> .years	27	0.83	0.45	(0.50, 1.36)	
Trend				0.50		
Colon cancer (407 cases)						
0 to 5	mg/m <sup>3</sup> .years	218				
> 5 to 10.1	mg/m <sup>3</sup> .years	63	0.88	0.43	(0.64, 1.21)	
> 10.1 to 21.2	mg/m <sup>3</sup> .years	63	0.85	0.32	(0.61, 1.17)	
> 21.2	mg/m <sup>3</sup> .years	63	0.90	0.54	(0.65, 1.26)	
Trend				0.93		
Rectal cancer (83 cases)						
0 to 5	mg/m <sup>3</sup> .years	41				
> 5 to 9.3	mg/m <sup>3</sup> .years	22	2.46	0.00	(1.33, 4.57)	*
> 9.3	mg/m <sup>3</sup> .years	20	1.01	0.98	(0.52, 1.96)	
Trend				0.67		
Bladder cancer (138 cases)						
0 to 5	mg/m <sup>3</sup> .years	75				
> 5 to 9.3	mg/m <sup>3</sup> .years	20	0.98	0.94	(0.57, 1.68)	
> 9.3 to 21.3	mg/m <sup>3</sup> .years	21	0.83	0.49	(0.48, 1.43)	
> 21.3	mg/m <sup>3</sup> .years	22	1.11	0.71	(0.64, 1.93)	
Trend				0.27		
Liver cancer (123 cases)						
0 to 5	mg/m <sup>3</sup> .years	73				
> 5 to 12.7	mg/m <sup>3</sup> .years	25	1.02	0.93	(0.60, 1.74)	
> 12.7	mg/m <sup>3</sup> .years	25	0.97	0.90	(0.56, 1.68)	
Trend				0.27		
Pancreatic cancer (315 cases)						

Table 2: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **soluble** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI
0 to 5	mg/m <sup>3</sup> .years	179			
> 5 to 8.3	mg/m <sup>3</sup> .years	46	1.28	0.19	(0.89, 1.86)
> 8.3 to 16.2	mg/m <sup>3</sup> .years	45	1.08	0.68	(0.74, 1.58)
> 16.2	mg/m <sup>3</sup> .years	45	0.87	0.48	(0.59, 1.28)
Trend				0.27	
Skin cancer (69 cases)					
0 to 5	mg/m <sup>3</sup> .years	43			
> 5 to 16.2	mg/m <sup>3</sup> .years	13	0.87	0.72	(0.42, 1.83)
> 16.2	mg/m <sup>3</sup> .years	13	1.37	0.44	(0.62, 3.02)
Trend				0.18	
Prostate cancer (418 cases)					
0 to 5	mg/m <sup>3</sup> .years	183			
> 5 to 10.9	mg/m <sup>3</sup> .years	79	0.97	0.87	(0.72, 1.32)
> 10.9 to 24.2	mg/m <sup>3</sup> .years	78	0.88	0.40	(0.64, 1.19)
> 24.2	mg/m <sup>3</sup> .years	78	1.12	0.48	(0.82, 1.53)
Trend				0.17	
Brain and nervous system cancers (128 cases)					
0 to 5	mg/m <sup>3</sup> .years	78			
> 5 to 11.9	mg/m <sup>3</sup> .years	25	1.21	0.49	(0.70, 2.08)
> 11.9	mg/m <sup>3</sup> .years	25	1.09	0.77	(0.61, 1.93)
Trend				0.95	
Leukemia (200 cases)					
0 to 5	mg/m <sup>3</sup> .years	119			
> 5 to 9.2	mg/m <sup>3</sup> .years	27	0.99	0.98	(0.62, 1.60)
> 9.2 to 19.9	mg/m <sup>3</sup> .years	27	0.85	0.52	(0.52, 1.39)
> 19.9	mg/m <sup>3</sup> .years	27	0.91	0.71	(0.55, 1.50)
Trend				0.70	
Breast cancer (76 cases)					
0 to 5	mg/m <sup>3</sup> .years	70			
> 5	mg/m <sup>3</sup> .years	6	0.38	0.06	(0.13, 1.05)

Table 3: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **synthetic** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI
Laryngeal cancer (73 cases)					
0	mg/m <sup>3</sup> .years	54			
> 0	mg/m <sup>3</sup> .years	19	1.54	0.31	(0.66, 3.59)
Lung cancer (1894 cases)					
0	mg/m <sup>3</sup> .years	1377			
> 0 to 0.3	mg/m <sup>3</sup> .years	173	0.92	0.42	(0.75, 1.13)
> 0.3 to 1.4	mg/m <sup>3</sup> .years	172	1.05	0.65	(0.85, 1.29)

Table 3: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **synthetic** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI
> 1.4	mg/m <sup>3</sup> .years	172	0.99	0.90	(0.81, 1.21)
Trend				0.99	
Esophageal cancer (176 cases)					
0	mg/m <sup>3</sup> .years	126			
> 0 to 0.7	mg/m <sup>3</sup> .years	25	1.08	0.80	(0.59, 1.99)
> 0.7	mg/m <sup>3</sup> .years	25	1.26	0.45	(0.69, 2.30)
Trend				0.19	
Stomach cancer (194 cases)					
0	mg/m <sup>3</sup> .years	150			
> 0 to 0.5	mg/m <sup>3</sup> .years	22	1.16	0.64	(0.63, 2.14)
> 0.5	mg/m <sup>3</sup> .years	22	1.03	0.92	(0.56, 1.90)
Trend				0.79	
Colon cancer (407 cases)					
0	mg/m <sup>3</sup> .years	310			
> 0 to 0.4	mg/m <sup>3</sup> .years	33	0.82	0.39	(0.52, 1.29)
> 0.4 to 1.7	mg/m <sup>3</sup> .years	32	0.99	0.97	(0.62, 1.58)
> 1.7	mg/m <sup>3</sup> .years	32	0.93	0.77	(0.59, 1.47)
Trend				0.99	
Rectal cancer (83 cases)					
0	mg/m <sup>3</sup> .years	59			
> 0 to 0.8	mg/m <sup>3</sup> .years	12	1.26	0.62	(0.51, 3.12)
> 0.8	mg/m <sup>3</sup> .years	12	1.36	0.50	(0.56, 3.31)
Trend				0.49	
Bladder cancer (138 cases)					
0	mg/m <sup>3</sup> .years	104			
> 0 to 0.5	mg/m <sup>3</sup> .years	17	1.07	0.85	(0.53, 2.15)
> 0.5	mg/m <sup>3</sup> .years	17	0.83	0.58	(0.42, 1.62)
Trend				0.18	
Liver cancer (123 cases)					
0	mg/m <sup>3</sup> .years	89			
> 0 to 0.4	mg/m <sup>3</sup> .years	17	0.99	0.98	(0.49, 2.01)
> 0.4	mg/m <sup>3</sup> .years	17	0.69	0.30	(0.34, 1.39)
Trend				0.02	*
Pancreatic cancer (315 cases)					
0	mg/m <sup>3</sup> .years	227			
> 0 to 0.3	mg/m <sup>3</sup> .years	30	0.95	0.83	(0.58, 1.55)
> 0.3 to 0.9	mg/m <sup>3</sup> .years	29	1.23	0.41	(0.75, 2.04)
> 0.9	mg/m <sup>3</sup> .years	29	0.83	0.45	(0.50, 1.36)
Trend				0.32	
Skin cancer (69 cases)					
0	mg/m <sup>3</sup> .years	52			
> 0	mg/m <sup>3</sup> .years	17	0.49	0.09	(0.22, 1.11)
Prostate cancer (418 cases)					
0	mg/m <sup>3</sup> .years	301			
> 0 to 0.5	mg/m <sup>3</sup> .years	39	1.17	0.49	(0.75, 1.84)

Table 3: Cox model estimates of the hazard ratio for selected cancer outcomes associated with exposure to **synthetic** metalworking fluids, controlling for other fluid types, calendar year, calendar year of hire, age, race, sex, and plant.

		Number of cases	HR	<i>p</i>	95% CI	
> 0.5 to 2	mg/m <sup>3</sup> .years	39	1.36	0.19	(0.86, 2.14)	
> 2	mg/m <sup>3</sup> .years	39	1.59	0.03	(1.04, 2.45)	*
Trend				0.18		
Brain and nervous system cancers (128 cases)						
0	mg/m <sup>3</sup> .years	94				
> 0 to 0.6	mg/m <sup>3</sup> .years	17	1.29	0.48	(0.63, 2.62)	
> 0.6	mg/m <sup>3</sup> .years	17	1.04	0.92	(0.51, 2.10)	
Trend				0.75		
Leukemia (200 cases)						
0	mg/m <sup>3</sup> .years	142				
> 0 to 0.9	mg/m <sup>3</sup> .years	29	1.30	0.36	(0.75, 2.25)	
> 0.9	mg/m <sup>3</sup> .years	29	1.66	0.07	(0.97, 2.84)	.
Trend				0.29		
Breast cancer (76 cases)						
0	mg/m <sup>3</sup> .years	60				
> 0	mg/m <sup>3</sup> .years	16	0.79	0.53	(0.39, 1.64)	

With additional model terms; exposure lagged 21 years; soluble referent at 5 mg/m<sup>3</sup>.y

