## **UAW-GM Cohort Study**

Suicide and fatal overdose after job loss (Cox analyses)

March 12, 2020

## **Tables and figures**

Table 1: Summary of the **UAW-GM Cohort** restricted to men employed in or after 1970.

	All records		Subset with complete records <sup>b</sup>		
N (person⋅years)	26 890	(933 187)	19 663	(650 685)	
Race, n (%)					
White	19 388	(72 %)	13 293	(68 %)	
Black	5 262	(20 %)	4 142	(21 %)	
Unknown	2 240	(8 %)	2 228	(11 %)	
Plant <sup>a</sup> , n (%)				•	
Plant 1	6 935	(26 %)	6 627	(34 %)	
Plant 2	10 351	(38 %)	6 412	(33 %)	
Plant 3	9 604	(36 %)	6 624	(34 %)	
Complete work records	19 663	(73 %)	19 663	(100 %)	
Year of hire	1967	(1956, 1975)	1965	(1953, 1971)	
Age at hire	24	(20, 31)	25	(20, 33)	
Year of birth	1942	(1927, 1950)	1936	(1923, 1947)	
Year of worker exit	1991	(1982, 1995)	1987	(1979, 1993)	
Age at worker exit	49	(40, 58)	52	(39, 61)	
Age at death among deceased	70	(60, 79)	71	(61, 80)	
Year of death among deceased	1999	(1989, 2008)	1997	(1988, 2007)	
Suicide cases	203	•	179		
Fatal overdose cases	55		34		

Notes: Statistics shown are median (first quartile, third quartile), unless otherwise indicated.

<sup>&</sup>lt;sup>a</sup> Some subjects worked at several sites; plant indicates the site of longest work record time.

 $<sup>^{\</sup>it b}$  Left work by December31, 1994 when employment records were truncated.

Figure 1: Crude rate (per 10,000 person-years) of suicide by calendar year in the **UAW-GM Cohort** restricted to men employed in or after 1970.




Table 2: Adjusted hazard ratio estimates for **suicide** by employment status in the **UAW-GM Cohort** restricted to men employed in or after 1970.

	Recorded worker exit date				Fuzzy worker exit date <sup>‡</sup>		
Job exit status	n	HR	95% CI	n	HR	95% CI	
At work	21	1.0	_	27	1.0		
Not at work	158	17.7	10.8, 29.1	152	12.5	7.9, 19.7	

Abbreviations: CI, confidence interval; HR, hazard ratio

Notes: Estimates were adjusted for plant, race, year of hire, and calendar year. Risk sets were indexed by age. Women were excluded from this analysis.

<sup>&</sup>lt;sup>‡</sup> Cases that occurred within a week after the recorded worker exit date were assumed to have occurred while still employed.

Table 3: Adjusted hazard ratio estimates for **suicide** and the combined outcome of **suicide** and **fatal overdose** in the **UAW-GM Cohort** restricted to men employed in or after 1970.

	Suid	Suicide			Suicide and fatal overdose		
Age at worker exit	n	HR	95% CI	n	HR	95% CI	
55 or older	42	1.0	_	45	1.0	_	
40 to 54	48	1.2	0.8, 1.8	51	1.2	0.8, 1.8	
30 to 39	41	1.7	1.1, 2.6	56	2.1	1.4, 3.1	
19 to 29	27	1.4	0.8, 2.3	38	2.0	1.2, 3.1	

Abbreviations: CI, confidence interval; HR, hazard ratio

Notes: Estimates were adjusted for race, plant, and worker exit date. Risk sets were indexed by time since worker exit. Women and those with unknown date of worker exit were excluded from this analysis.

Figure 2: Crude rate (per 10,000 person-years) of the combined outcome of suicide and fatal overdose by calendar year in the **UAW-GM Cohort** restricted to men employed in or after 1970.

