

Random Fourier Features for Kernel Ridge Regression

Approximation Bounds and Statistical Guarantees

Kailong Wang¹

1766

¹Ph.D. of ECE
Rutgers University University

ECE 539 HDP, May 3, 2023



Table of Contents

① Motivation


② Random Fourier Features



Linear Classification with Non-linear Input


Consider a binary classification problem with non-linear (e.g. polynomial) samples. This is not separable with linear function.

$$\text{(e.g. } X = \begin{bmatrix} x_{1,1} & x_{1,2} \\ x_{2,1} & x_{2,2} \\ \vdots & \vdots \\ x_{N,1} & x_{N,2} \end{bmatrix} \in \mathbb{R}^{N \times 2}.)$$



Lifting

One idea is to **LIFT** the samples into a higher dimensional space in which the samples are linearly separable.



The Lifting function in this case is $\phi(X) = \begin{bmatrix} x_{1,1}^2 & x_{1,2}^2 & \sqrt{2}x_{1,1}x_{1,2} \\ x_{2,1}^2 & x_{2,2}^2 & \sqrt{2}x_{2,1}x_{2,2} \\ \dots & \dots & \dots \\ x_{N,1}^2 & x_{N,2}^2 & \sqrt{2}x_{N,1}x_{N,2} \end{bmatrix}$.



Curse of Dimensionality

Consider solving the above problem with *support vector machine* (SVM).

$$\mathcal{L}(\mathbf{w}, \alpha) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_n^N \sum_m^N \alpha_n \alpha_m y_n y_m (x_n^\top x_m).$$

The \mathbf{w} is the linear decision boundary and α is a vector of Lagrange multipliers.



Curse of Dimensionality

Consider solving the above problem with *support vector machine* (SVM).

$$\mathcal{L}(\mathbf{w}, \alpha) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_n^N \sum_m^N \alpha_n \alpha_m y_n y_m (x_n^\top x_m).$$

The \mathbf{w} is the linear decision boundary and α is a vector of Lagrange multipliers.

We need to use lifting function $\phi(X)$ to make the samples linearly separable. Specifically, we replace $(x_n^\top x_m)$ with $(\phi(x_n)^\top \phi(x_m))$.

$$\begin{aligned} \phi(x_n)^\top \phi(x_m) &= \begin{bmatrix} x_{n,1}^2 & x_{n,2}^2 & \sqrt{2}x_{n,1}x_{n,2} \end{bmatrix} \begin{bmatrix} x_{m,1}^2 & x_{m,2}^2 & \sqrt{2}x_{m,1}x_{m,2} \end{bmatrix}^\top \\ &= x_{n,1}^2 x_{m,1}^2 + x_{n,2}^2 x_{m,2}^2 + 2x_{n,1}x_{n,2}x_{m,1}x_{m,2} \end{aligned}$$



Curse of Dimensionality

Consider solving the above problem with *support vector machine* (SVM).

$$\mathcal{L}(\mathbf{w}, \alpha) = \sum_{n=1}^N \alpha_n - \frac{1}{2} \sum_n^N \sum_m^N \alpha_n \alpha_m y_n y_m (x_n^\top x_m).$$

The \mathbf{w} is the linear decision boundary and α is a vector of Lagrange multipliers.

We need to use lifting function $\phi(X)$ to make the samples linearly separable. Specifically, we replace $(x_n^\top x_m)$ with $(\phi(x_n)^\top \phi(x_m))$.

$$\begin{aligned} \phi(x_n)^\top \phi(x_m) &= \begin{bmatrix} x_{n,1}^2 & x_{n,2}^2 & \sqrt{2}x_{n,1}x_{n,2} \end{bmatrix} \begin{bmatrix} x_{m,1}^2 & x_{m,2}^2 & \sqrt{2}x_{m,1}x_{m,2} \end{bmatrix}^\top \\ &= x_{n,1}^2 x_{m,1}^2 + x_{n,2}^2 x_{m,2}^2 + 2x_{n,1}x_{n,2}x_{m,1}x_{m,2} \end{aligned}$$

Calculate the inner product in the \mathbb{R}^3 across all N pairs of samples is acceptable. However, the lifting function $\phi(X)$ is usually very high dimensional.



Kernel Trick

Consider the following derivation,

$$\begin{aligned}(x_n^\top x_m)^2 &= ([x_{n,1} \ x_{n,2}][x_{m,1} \ x_{m,2}]^\top)^2 \\&= (x_{n,1}x_{m,1} + x_{n,2}x_{m,2})^2 \\&= x_{n,1}^2x_{m,1}^2 + x_{n,2}^2x_{m,2}^2 + 2x_{n,1}x_{n,2}x_{m,1}x_{m,2} \\&= \phi(x_n)^\top \phi(x_m)\end{aligned}$$



Kernel Trick

Consider the following derivation,

$$\begin{aligned}(x_n^\top x_m)^2 &= ([x_{n,1} \ x_{n,2}][x_{m,1} \ x_{m,2}]^\top)^2 \\&= (x_{n,1}x_{m,1} + x_{n,2}x_{m,2})^2 \\&= x_{n,1}^2x_{m,1}^2 + x_{n,2}^2x_{m,2}^2 + 2x_{n,1}x_{n,2}x_{m,1}x_{m,2} \\&= \phi(x_n)^\top \phi(x_m)\end{aligned}$$

Instead of computing inner product in the high dimensional space, we compute the inner product in the original space.



Kernel Trick

Consider the following derivation,

$$\begin{aligned}(x_n^\top x_m)^2 &= ([x_{n,1} \ x_{n,2}][x_{m,1} \ x_{m,2}]^\top)^2 \\&= (x_{n,1}x_{m,1} + x_{n,2}x_{m,2})^2 \\&= x_{n,1}^2x_{m,1}^2 + x_{n,2}^2x_{m,2}^2 + 2x_{n,1}x_{n,2}x_{m,1}x_{m,2} \\&= \phi(x_n)^\top \phi(x_m)\end{aligned}$$

Instead of computing inner product in the high dimensional space, we compute the inner product in the original space.

The function

$$K(x_n, x_m) = (x_n^\top x_m)^2 = \phi(x_n)^\top \phi(x_m)$$

is called a **kernel function**.



There must be disadvantages...

Given training data $(x_1, y_1), (x_2, y_2), \dots, (x_N, y_N) \in \mathcal{X} \times \mathcal{Y}$, where $\mathcal{X} \subseteq \mathbb{R}^d$ and $\mathcal{Y} \subseteq \mathbb{R}$. Consider *Kernel Ridge Regression* (KRR), with $\phi(\mathcal{X}) \subseteq \mathbb{R}^k$, where $k \rightarrow \infty$

$$\mathcal{L}(\mathbf{w}, \lambda) = \underset{\mathbf{w}}{\operatorname{argmin}} \sum_n^N (y_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2 + \lambda \mathbf{w}^\top \mathbf{w}.$$

Solving it with Lagrange multipliers α , which is the solution of

$$(\mathbf{K} + \lambda \mathbf{I}_k) \alpha = \mathbf{y},$$

requires $\Theta(k^3)$ time and $\Theta(k^2)$ memory. Here $\mathbf{K} \in \mathbb{R}^{k \times k}$ is the kernel matrix or Gram matrix defined by $\mathbf{K}_{nm} \equiv K(\mathbf{x}_n, \mathbf{x}_m)$.



There must be disadvantages...

Given training data $(x_1, y_1), (x_2, y_2), \dots, (x_N, y_N) \in \mathcal{X} \times \mathcal{Y}$, where $\mathcal{X} \subseteq \mathbb{R}^d$ and $\mathcal{Y} \subseteq \mathbb{R}$. Consider *Kernel Ridge Regression* (KRR), with $\phi(\mathcal{X}) \subseteq \mathbb{R}^k$, where $k \rightarrow \infty$

$$\mathcal{L}(\mathbf{w}, \lambda) = \underset{\mathbf{w}}{\operatorname{argmin}} \sum_n^N (y_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2 + \lambda \mathbf{w}^\top \mathbf{w}.$$

Solving it with Lagrange multipliers α , which is the solution of

$$(\mathbf{K} + \lambda \mathbf{I}_k) \alpha = \mathbf{y},$$

requires $\Theta(k^3)$ time and $\Theta(k^2)$ memory. Here $\mathbf{K} \in \mathbb{R}^{k \times k}$ is the kernel matrix or Gram matrix defined by $\mathbf{K}_{nm} \equiv K(\mathbf{x}_n, \mathbf{x}_m)$.

Intuition: Can we find a kernel function which lifts \mathcal{X} to \mathbb{R}^s , where $d < s \ll k$, while not sacrifices model performance?



Some Prerequisites

Shift Invariant Kernel (Radial Basis Kernel (RBK))

A kernel function $K(\mathbf{x}_n, \mathbf{x}_m)$ is called **shift invariant** if it can be written as $K(\mathbf{x}_n, \mathbf{x}_m) = g(\mathbf{x}_n - \mathbf{x}_m)$ for some function $g(\cdot)$ (e.g. $K_{Gaussian}(\mathbf{x}_n, \mathbf{x}_m) = \exp(-\gamma \|\mathbf{x}_n - \mathbf{x}_m\|_2^2)$).

Mercer's Theorem

A continuous function $K(\mathbf{x}_n, \mathbf{x}_m)$ is a valid kernel function if and only if the kernel matrix \mathbf{K} is **positive definite**.

Bochner's Theorem

A continuous function $g(\cdot)$ is **positive definite** if and only if it is the Fourier transform of a non-negative measure.



Random Fourier Features

Conclusion

A continuous **shift invariant** kernel $K(\mathbf{x}_n, \mathbf{x}_m)$, which is **positive definite** (Mercer's Theorem), is the Fourier transform of a non-negative measure $p(\cdot)$.

$$\phi(\mathbf{x}_n)^\top \phi(\mathbf{x}_m) = K(\mathbf{x}_n, \mathbf{x}_m) = K(\mathbf{x}_n - \mathbf{x}_m) \quad (1)$$

$$= \int_{\mathbb{R}^s} p(\omega) \exp(i\omega^\top (\mathbf{x}_n - \mathbf{x}_m)) d\omega \quad (2)$$

$$= \mathbb{E}_\omega [\xi(\mathbf{x}_n) \xi(\mathbf{x}_m)^*] \quad (3)$$

Here $\xi(\mathbf{x}) = \exp(i\omega^\top \mathbf{x}) = \begin{bmatrix} \cos(\omega^\top \mathbf{x}) \\ \sin(\omega^\top \mathbf{x}) \end{bmatrix}$ and hence $\xi(\mathbf{x}_n) \xi(\mathbf{x}_m)^*$ is an unbiased estimator of $K(\mathbf{x}_n, \mathbf{x}_m)$.



Algorithm

Since both the $p(\cdot)$ and $K(\Delta)$ are real-valued, we can replace $\xi(\mathbf{x})$ with $z_\omega(\mathbf{x}) = [\sqrt{2} \cos(\omega^\top \mathbf{x} + b)]$ where ω is drawn from $p(\omega)$ and b is uniformly drawn from $[0, 2\pi]$. Then (3) becomes $\mathbb{E}_\omega[z(\mathbf{x}_n)z(\mathbf{x}_m)^\top]$

Algorithm 1 An algorithm with caption

Require: $n \geq 0$

Ensure: $y = x^n$

```
1:  $y \leftarrow 1$ 
2:  $X \leftarrow x$ 
3:  $N \leftarrow n$ 
4: while  $N \neq 0$  do
5:   if  $N$  is even then
6:      $X \leftarrow X \times X$ 
7:      $N \leftarrow \frac{N}{2}$ 
8:   else if  $N$  is odd then
9:      $y \leftarrow y \times X$ 
10:     $N \leftarrow N - 1$ 
11:  end if
```

▷ This is a comment



Two-column slide

This is a subtitle

This is a text in first column.

$$E = mc^2$$

First item ℓ

Second item

This text will be in the second column and on a second thought this is a nice looking layout in some cases.

