

1 Chapter 3.1

Problem 2. How many integers in $[100]$ are not divisible by 4, 6, or 7?

Answer:

$$\begin{aligned} N_{\geq}(\emptyset) &= 100 & N_{\geq}(d_4, d_6) &= \left\lfloor \frac{100}{2 \cdot 6} \right\rfloor = 8 \\ N_{\geq}(d_4) &= \left\lfloor \frac{100}{4} \right\rfloor = 25 & N_{\geq}(d_4, d_7) &= \left\lfloor \frac{100}{4 \cdot 7} \right\rfloor = 3 \\ N_{\geq}(d_6) &= \left\lfloor \frac{100}{6} \right\rfloor = 16 & N_{\geq}(d_6, d_7) &= \left\lfloor \frac{100}{6 \cdot 7} \right\rfloor = 2 \\ N_{\geq}(d_7) &= \left\lfloor \frac{100}{7} \right\rfloor = 14 & N_{\geq}(d_4, d_6, d_7) &= \left\lfloor \frac{100}{2 \cdot 6 \cdot 7} \right\rfloor = 1 \end{aligned}$$

The answer is $100 - (25 + 16 + 14) + (8 + 3 + 2) - 1 = 57$

Problem 6. Answer the hat-check problem (i.e., the problem of derangement) for general n . This number is known as D_n . The question is “How many ways can the hats of n people be re-distributed so that each person receives exactly one hat, but no person receives their own hat”?

Answer:

Before diving into the general n solution. Lets look at an example party problem, with 3 people, called Kaelin, Alyse and Donna. The three cases to avoid are the following:

$$\begin{aligned} h_k &:= \text{“Kaelin gets his own hat back”} \\ h_a &:= \text{“Alyse gets her own hat back”} \\ h_d &:= \text{“Donna gets her own hat back”} \end{aligned}$$

So the set $P = \{h_k, h_a, h_d\}$ and the possible subsets of the problem are:

$$J = \{\emptyset, \{h_k\}, \{h_a\}, \{h_d\}, \{h_k h_a\}, \{h_k h_d\}, \{h_a h_d\}, \{h_k h_a h_d\}\}$$

Instead of looping through each element in J and adding them individually, a better approach would be to group cases where the same amount of people receive their own hat back. For instance.

- $j = 0$, at least no one gets their hat back, $\emptyset$, 1 option
- $j = 1$, at least one person gets their hat back, $\{h_k\}, \{h_a\}, \{h_d\}$, 3 options
- $j = 2$, at least two persons get their hat back, $\{h_k h_a\}, \{h_k h_d\}, \{h_a h_d\}$, 3 options
- $j = 3$, at least three persons get their hat back, $\{h_k h_a h_d\}$, 1 option

Each of the number of options can be represented as “With 3 people, how many ways can j people get their hat back”? This is represented as $\binom{3}{j}$ or $\binom{n}{j}$ for a general amount of people, n .

Now we look at how to distribute the hats to people. If we didn’t need to fix anyone getting their hat, the ways to distribute them would be $3!$. Kaelin would have 3 options, Alyse would have 2 options and Donna would have 1 option. If instead we fixed one person, say Kaelin, then you only have 2 options to choose from, $2!$. This is represented mathematically as $(3 - j)!$ or $(n - j)!$ for the general case.

Putting this together for the 3-person example we have:

$$D_3 = \sum_{j=0}^3 \binom{3}{j} (-1)^j (3 - j)!$$

Where the $(-1)^j$ is the book keeping for ensuring odd values are subtracted and evens added. This prevents accidental double counting of values that appear in multiple cases. For the general case, where its n people, the equation is:

$$D_n = \sum_{j=0}^n \binom{n}{j} (-1)^j (n-j)!$$

Algebraically, this can be simplified starting with substituting the combination for its formula.

$$D_n = \sum_{j=0}^n \frac{n!}{j!(n-j)!} (-1)^j (n-j)! = n! \sum_{j=0}^n \frac{(-1)^j}{j!}$$

Problem 7. The number D_n is the hat check problem for general n .

- (a) Calculate $\lim_{n \rightarrow \infty} \frac{D_n}{n!}$
- (b) Prove that for any n , D_n equals the closest integer to $n!/e$.

Answer:

- (a) Looking at the limit of the hat check derangement over the total ways to distribute the hats.

$$\frac{D_n}{n!} = \frac{n!}{n!} \sum_{j=0}^n \frac{(-1)^j}{j!} = \sum_{j=0}^n \frac{(-1)^j}{j!}$$

The inverse of Euler's number can be expressed with the following infinite series.

$$\frac{1}{e} = \sum_{j=0}^{\infty} \frac{(-1)^j}{j!}$$

As $n \rightarrow \infty$ the summation approaches $1/e$. So the limit is the inverse of Euler's number.

- (b) Explanation of the alternating series remainder term theorem from calculus here.

Before jumping right into using the alternating series remainder theorem, we need to first show the series converges. The criteria is:

$$\lim_{j \rightarrow \infty} \frac{1}{j!} = 0 \quad \text{and} \quad \frac{1}{(j+1)!} < \frac{1}{j!}$$

So it passes. We can now apply the alternating series remainder theorem. This says the error of evaluated term will be less than or equal to the value of the first term left off. When $j = 0$ or 1 the series value is $1, 0$. This doesn't make sense to begin evaluating the series at these values. The first value that makes sense is $j = 2$, so two people.

$$\sum_{j=0}^2 \frac{(-1)^j}{j!} = 1 - 1 + 1/2 = 1/2$$

The next term is $j = 3$ which gives $1/6$. So the low and high side of the $j = 2$ case is $1/2 \pm 1/6 = (2/6, 4/6)$ or $(1/3, 2/3)$. The value this finally converges to is $1/e = 1.1/3$ which falls inside the boundary.

Proof. For any value of n the number of derangements is:

$$D_n = n! \sum_{j=0}^n \frac{(-1)^j}{j!}$$

The number of derangements over total permutations $D_n/n!$ is a converging series, as it meets the alternating series theorem criteria.

$$\lim_{j \rightarrow \infty} \frac{1}{j!} = 0 \quad \text{and} \quad \frac{1}{(j+1)!} < \frac{1}{j!}$$

This series converges to $1/e$ as n approaches infinity.

$$\frac{D_n}{n!} = \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} = \frac{1}{e}$$

The alternating series remainder theorem states the max difference between the summation evaluated at n and the final converged value will be less than the single value at $n+1$. For the probability of derangements this appears as

$$\sum_{j=0}^n \frac{(-1)^j}{j!} - \frac{1}{e} \leq \frac{1}{(n+1)!}$$

Multiplying through by $n!$ for number of derangements.

$$n! \sum_{j=0}^n \frac{(-1)^j}{j!} - \frac{n!}{e} \leq \frac{n!}{(n+1)n!} = \frac{1}{n+1}$$

As such the difference between D_n and $n!/e$ will always be less than an integer, allowing rounding the irrational number to provide an accurate count. $\square$

Problem 8. How many ways can you distribute 20 identical objects to 10 distinct recipients so that each recipient receives at most five objects?

Answer:

The empty set at least (base case) is how many ways can you distribute 20 identical objects to 10 distinct recipients. This is

$$N_{\geq}(\emptyset) = \left(\binom{10}{20} \right)$$

Then you need to evaluate the following cases which are:

- (a) P_1 - 1 person receives 6 or more objects
- (b) P_2 - 2 people receives 6 or more objects
- (c) P_3 - 3 people receives 6 or more objects

Taking the first case, P_1 . Out of the 10 recipients, you can select 1 recipient to receive at least 6 objects. mathematically this is $\binom{10}{1}$, then you need to distribute the remaining 14 objects to the 10 recipients. Those objects can go to any person, all that matters is that at least 1 recipient has at least 6 objects, $\left(\binom{10}{14} \right)$.

$$N_{\geq}(P_1) = \binom{10}{1} \left(\binom{10}{14} \right)$$

Second case P_2 , now two recipients get at least 6 objects. So you have 8 objects left over to distribute to the ten recipients.

$$N_{\geq}(P_2) = \binom{10}{2} \left(\binom{10}{8} \right)$$

Third case P_3 , now three recipients get at least 6 objects. So you have 2 objects left over to distribute to the ten recipients.

$$N_{\geq}(P_3) = \binom{10}{3} \left(\binom{10}{2} \right)$$

The total answer is:

$$\binom{\binom{10}{20}}{\binom{10}{1}} - \binom{\binom{10}{14}}{\binom{10}{1}} + \binom{\binom{10}{8}}{\binom{10}{2}} - \binom{\binom{10}{2}}{\binom{10}{3}}$$

Problem 9. *How many ways can you distribute k identical objects to n distinct recipients so that each recipient receives at most r objects?*

Answer: I just took problem 8 and make it a general case for problem 9.

$$\sum_{j=0}^{k/(r+1)} (-1)^j \binom{n}{j} \binom{n}{k-j(r+1)}$$

Problem 17. *A taxi drives from the intersection labeled A to the intersection labeled B in the grid of streets shown on page 94. The driver can only drive north (up) or east (right). Traffic reports indicate that there is heavy congestion at the intersections identified. How many routes from A to B can the driver take that ...*

- (a) *Avoid all the congested intersections?*
- (b) *Pass through at most one congested intersection?*

Answer: Start by defining the different types of points. Ones that can go up and right. Ones that can only go up, ones that can only go right. Use the knowledge that each point has two choices (up or right) with a few exceptions.