

1 Chapter 2.1

Problem 4. Consider the possible functions $f : [7] \rightarrow [9]$

- (a) How many have $f(3) = 8$?
- (b) How many have $f(3) \neq 8$?
- (c) How many have $f(1) \neq 5$ and are one-to-one?
- (d) How many have $f(i)$ even, for all i ?
- (e) How many have $\text{rng}(f) = \{5, 6\}$?
- (f) How many in which f^{-1} is not a function?

Answer:

- (a) $f(3)$ has 1 choice. This leaves 6 other values that can go into 9 other bins. I have nine bins, that I can place six values in, with repetition allowed. This is 9^6 .
- (b) $f(3)$ has 8 choices, while the six other values have 9 choices. This is $8 \cdot 9^6$.
- (c) $f(1)$ has 8 choices, $f(2)$ has eight choices (since one was removed by $f(1)$.) Then $f(3)$ has seven choices, $f(4)$ has six choices... This is $8 \cdot (8)_6 = \frac{8 \cdot 8!}{(8-6)!}$.
- (d) Instead of 9 choices, we now have 4 choices. So 7 values have 4 choices. This is 4^7 .
- (e) Now we have only two choices, $\{5, 6\}$ for 7 values. This is 2^7 . But, we must guarantee at least one value in $\{5\}$ and one value in $\{6\}$. So subtract out when all the values map to 5 or all the values map to 6, which is two cases. Final answer is $2^7 - 2$.
- (f) For f^{-1} to be a function, the original function f must be one-to-one. We are looking for where the original function is not one-to-one. Base options is 9 choices for 7 values. So 9^7 . But, we need to subtract out the one-to-one option, which is $(9)_7$. The final answer is $9^7 - (9)_7$.

Problem 7. Find the number of onto functions $[k] \rightarrow [4]$.

Answer: I am assuming that $k \geq 4$. The base case is 4^k options for distribution. Then need to subtract out the following:

- (a) Missing 3 of the elements in $[4]$. Answer is $4^1 = 4$.
- (b) Missing 2 of the elements in $[4]$. You have $\binom{4}{2} = 6$ elements to miss. You then have 2^k ways to distribute the values to the remaining elements. Each distribution has 2 ways that all the values end up on one element instead of both, so remove those. The answer is $6(2^k - 2)$.
- (c) Missing 1 of the elements in $[4]$. You have $\binom{4}{3} = 4$ different elements to miss. Once you miss, you can select 3^k places to go. But you need to guarantee that you don't miss more than 1 element. There are 3 ways to miss 2 elements with 2^k places to go. There are 3 ways to miss 3 elements with 1^k places to go. The answer is $4(3^k - 3(2^k - 2) - 3 \cdot 1^k)$

Final answer is $4^k - (4 + 6(2^k - 2) + 4(3^k - (3 + 3(2^k - 2))))$

Problem 11. Give a combinatorial proof. For $n \geq 1$ and $k \geq 1$, then $2^{kn} > \max\{n^k, k^n\}$. Hint: Compare relations to functions.

Answer: Before diving into the details with the general case, I'm going to do a numerical example to get us warmed up. Let us define two sets.

$$K = [4] = \{1, 2, 3, 4\} \quad N = [3] = \{1, 2, 3\}$$

All the possible relations from K to N are contained in the Cartesian product of $K \times N$.

$$K \times N = \{\{1, 1\}, \{1, 2\}, \{1, 3\} \dots \{4, 1\}, \{4, 2\}, \{4, 3\}\} \quad (1)$$

Every possible relation on $K \rightarrow N$ is a subset of $K \times N$. For instance, for the relation $>$ would have $\{\{2, 1\}, \{3, 1\}, \{3, 2\}, \{4, 1\}, \{4, 2\}, \{4, 3\}\}$. So all the possible relations is $\mathcal{P}(K \times N)$, the power set of the cartesian product of $K \times N$. So, the number of possible relations is $2^{|K \times N|}$. If $k = |K|$ and $n = |N|$ then it is 2^{kn} .

Now, we focus our attention to functions instead of relations. The basic definition of a function is that every input can only have one output. Thinking about this as distributions, the first number in K has 3 potential choices. The second number in K also has 3 choices, and so on. As a result, when looking at functions, there are $n^k = 3^4 = 81$ possible functions from $K \rightarrow N$. Likewise, if we go the opposite way, from $N \rightarrow K$, there are $k^n = 4^3 = 64$ possible functions.

What this is saying, is the number of possible relations between two sets will always be bigger than the possible functions from one set to another and vice versa.

Proof. Suppose that $k, n \geq 1$ and K, N are sets of the first positive integers k, n denoted as $K = [k]$ and $N = [n]$. Then the set of all possible relations between K and N is the powerset of $K \times N$, denoted as $\mathcal{P}(K \times N)$. The cardinality of the powerset is $|\mathcal{P}(K \times N)| = 2^{kn}$, or the number of possible relations.

For a relation to be considered a function, every value in the domain must map to only one value in the codomain. The function $f : K \rightarrow N$ has n^k possible variations. The set of all function variations is a subset of all the relation variations. As such, $n^k \leq 2^{kn}$.

The inequality can be written $n^k < 2^{kn}$ by recognizing the empty set is inside the powerset but will never be considered to meet the definition of a function.

Now reversing the roles of K and N with the function $f : N \rightarrow K$ which has k^n possible variations. The same argument from above is used to prove $k^n < 2^{kn}$. This proves that $2^{kn} > \max\{n^k, k^n\}$. $\square$

2 Chapter 2.2

Problem 4. Give combinatorial or bijective proofs of the following. Part of your job is to determine all values of n, k , and / or m for which the identities are valid.

$$(a) \quad 3^n = \sum_{k=0}^n \binom{n}{k} 2^{n-k}$$

$$(b) \quad \binom{n}{k} \binom{k}{j} = \binom{n}{j} \binom{n-j}{k-j}$$

$$(c) \quad \left\langle \binom{n}{k} \right\rangle = \left\langle \binom{k+1}{n-1} \right\rangle$$

Answer:

- (a) LHS: How many distributions can be made with n distinct objects (different colored balls) to 3 distinct recipients, {Moe, Curly, Larry}.

RHS: Focus attention on the first recipient, Moe. Start with $k = 0$, meaning Moe will get all the combinations of zero balls. The remaining balls $n - k$ are distributed in various manners to Curly and Larry. Next, $k = 1$, meaning Moe will get all the combinations of one distinct ball. Curly and Larry receive the number of balls that Moe did not get. This pattern is repeated

until Moe gets the combinations of all the balls and Curly / Larry receive none. These different cases are added together.

Example is made with two balls, red and blue $\{R, B\}$. The table below shows $k = 0, 1, 2$ and the various ball distributions. As known, $3^2 = 9$ which is the number of permutations below.

k	Moe	Curly	Larry
0	$\emptyset$	R,B	$\emptyset$
0	$\emptyset$	$\emptyset$	R,B
0	$\emptyset$	R	B
0	$\emptyset$	B	R
1	R	B	$\emptyset$
1	R	$\emptyset$	B
1	B	R	$\emptyset$
1	B	$\emptyset$	R
2	R,B	$\emptyset$	$\emptyset$

- (b) LHS: A concert has n -distinct people attending. The concert is giving away k -guitars. Of the k -guitars a small subset of size j are signed. Each person can only receive one guitar and each guitar can only be signed once. How many different ways can the unsigned and signed guitars be distributed?

RHS: The same concert has n -distinct people attending. The concert starts by giving away j -guitars that are signed. After the signed guitars are given away, a new subgroup is made of $n - j$ people that didn't receive one. These people are eligible for $k - j$ unsigned guitars. How many ways can the signed and unsigned guitars be distributed.

For the identity to be valid, $n > k > j$.

- (c) The number of size k multisets from an n -set equals the number of size k subsets from an $n+k-1$ set. The right hand and left hand of the equivalence equation are shown using non-repeating subset notation.

$$\binom{n}{k} = \binom{n+k-1}{k}, \quad \binom{k+1}{n-1} = \binom{k+1+n-1-1}{n-1} = \binom{n+k-1}{n-1}$$

As such, the multiset equivalence statement can be written as:

$$\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$$

I need to finish this later and show that with a $[n+k+1]$ set that the complement of the number of k subsets is $n-1$ subsets. So instead, we can count the complement.

Problem 5. What does $\binom{n-1}{k-1} + \binom{n-2}{k-1} + \binom{n-3}{k-1} + \dots + \binom{k-1}{k-1}$ equal? Make a conjecture and then give a combinatorial proof.

My conjecture is that:

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-2}{k-1} + \binom{n-3}{k-1} + \dots + \binom{k-1}{k-1}$$

Problem 7. Determine the number of solutions to each of the following equations. Assume all z_i are nonnegative integers unless stated otherwise.

(a) $z_1 + z_2 + z_3 + z_4 = 1$

(b) $z_1 + z_2 + 10z_3 = 8$

(c) $z_1 + z_2 + \dots + z_{20} = 401$ where each $z_i \geq 1$.

Answer:

(a) $\binom{4}{1} = 4$

(b) In this problem, z_3 can only be 0, if $z_3 > 0$ then z_1 or z_2 are forced to be negative which violates the constraints. The answer is $\binom{2}{8} = \binom{9}{8} = 9$

(c) $\binom{20}{401-20} = \binom{400}{381} = \text{big}$

3 Chapter 2.3

Problem 2. For any integer $n \geq 2$, how many onto functions $[n] \rightarrow [n-1]$ are possible? Give a formula that doesn't involve Stirling numbers.

Problem 7. Find the number of equivalence relations on an n -set.

Problem 14. Here is a simple recursive python program for computing $S(n, k)$ based on Theorem 2.3.1. The program assumes $n, k \geq 0$. It works but it is extremely wasteful. Why? Design a more efficient algorithm.

```
def Stirling(n: int, k: int):
    if n == k:
        return 1
    elif n < k:
        return 0
    elif n > 0 and k == 0:
        return 0
    else:
        return Stirling(n-1, k-1) + k*Stirling(n-1, k)
```

Yeah, this is some garbage code for show.

Problem 17. Solve a linear system to find numbers a, b, c, d, e so that the following polynomial equation is true:

$$x^4 = a \cdot x_0 + b \cdot x_1 + c \cdot x_2 + d \cdot x_3 + e \cdot x_4$$

In the equation, the following represent $x_1, x_2 \dots x_0$.

$$\begin{aligned} x_4 &= x(x-1)(x-2)(x-3) \\ x_3 &= x(x-1)(x-2) \\ x_2 &= x(x-1) \\ x_1 &= x \\ x_0 &= 1 \end{aligned}$$

Express the solution a, b, c, d, e in terms of numbers studied in this section. Use linear algebra and formulas for expanding the coefficients.