

For Doctor

Other Doctors

MDS-LANCET LABORATORIES-EMAIL

27 MASERU STREET

EAST LEGON

.... ACCRA

Patient : NANCY TWUMASI-ASIAMAH

Guarantor : MRS N TWUMASI-ASIAMAH

Doctors Ref: NOT AVAILABLE

MedAid : CASH N

Age/Sex/DOB: 41 / F / 19770828

Tel : 0244631462

Id Num : TAN0262092929

(W) NOT AVAILABLE

Lab Ref : 706249212

Collection Date : 08/09/18 1216

MRI No. : GH01417665

Received Date : 08/09/18 1224

Spec # : 0908:EA00061L

FINAL Report Date : 22/09/18 1108

Requested : ., THY AB, TSH RECEPTOR AB, THYROGLOBULIN
ENDOCRINOLOGY

Test	Result	Reference
> THYROGLOBULIN	269.0 ug/L	< 55
The presence of thyroglobulin antibodies may interfere with the thyroglobulin assay. The result of this test must be treated with reserve if the patient has a positive thyroglobulin antibody test.		
Please note that biotin may interfere with this assay leading to falsely depressed values. We recommend that patients don't take Biotin for 48 hours before testing. Please contact Chemical Pathologist for further discussion if required.		
The presence of thyroglobulin antibodies may interfere with the thyroglobulin assay. The result of this test must be treated with reserve if the patient has a positive thyroglobulin antibody test.		
> Anti - TG Antibody	10.41 IU/mL	H < 4.11
> Anti - TPO Antibody	> 1000.00 IU/mL	H < 5.61
> TSH RECEPTOR ANTIBODIES	< 0.89 IU/L	< 1.80

This test may be useful in the following clinical situations:

- *Determining the cause of thyrotoxicosis in pregnancy.
- *Patients with normal thyroid function tests but clinically suspected Graves disease.
- *Determining the risk of a pregnant woman giving birth to a newborn with neonatal

** Continued on Next Page **