



Ms. ABIR ABDUL MALIK

PID NO: P36180007457

Age: 20 Year(s) Sex: Female

Reference:

Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180107742

Registered On:

02/07/2018 03:29 PM

Collected On:

02/07/2018

Reported On:

03/07/2018 10:39 AM

Investigation	Observed Value	Unit	Biological Reference Interval
FSH - Follicle Stimulating Hormone (Serum)	6.06	mIU/mL	Normal Menstruating Women Follicular Phase : 3.0 - 12.0 Mid Cycle Phase : 8.0 - 22.0 Luteal Phase : 2.0 - 12.0 Post Menopausal : 35.0 - 151.0
LH- Luteinizing hormone (Serum,CMIA)	2.88	mIU/mL	Follicular phase: 2.4-12.6 Midcycle peak: 14.0-95.66 Luteal phase: 1.0-11.4 Post menopausal: 7.7-58.5 Post Menopausal: 7.7-58.2

Interpretation :

Intact PTH has been demonstrated to be labile and is susceptible to fragmentation. This instability depends on both time and temperature . In room temperature EDTA sample stability is 8 hours and serum is for 4 hours. At 4degree C. EDTA sample stability is 72 hours and serum is for 48 hours.



Ms. ABIR ABDUL MALIK

PID NO: P36180007457

Age: 20 Year(s) Sex: Female

Reference:

Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180107742

Registered On:

02/07/2018 03:29 PM

Collected On:

02/07/2018

Reported On:

03/07/2018 10:39 AM

Investigation

Prolactin
(Serum,CMA)

Observed Value

15.75

Unit

ng/mL

Biological Reference Interval

5.18-26.53

Interpretation:

Prolactin secretion from pituitary shows significant diurnal, episodic and cyclical variations. Following is a suggested approach to hyperprolactinaemia in females -

Serum Prolactin levels	Interpretation	Remark "
5.18 to 26.53 ng/ml	Normal	Biological Reference Interval "
26.53 to 50 ng/ml	Mild prolactin excess	Often seen with physiological conditions like physical/emotional stress, exercise, pregnancy, lactation, etc. This may not be associated with clinical hyperprolactinaemia & needs review after a month"
51 to 75 ng/ml	Moderate prolactin excess	Often associated with clinical hyperprolactinaemia(short luteal phase,oligomenorrhea), hypothyroidism (often subclinical), macroprolactinaemia."
Above 100 ng/ml	Marked prolactin excess	Often associated with clinical hyperprolactinaemia- hypogonadism, amenorrhea, galactorrhea, hypothyroidism (often subclinical), macroprolactinaemia."
Above 200 ng/ml	Marked prolactin excess	Often associated with pituitary adenoma requiring further workup. High levels may be repeated with tripooled sample.

References :

1. Diagnosis & Treatment of hyperprolactinaemia. The endocrine society clinical practice guideline, 2011
2. Diagnosis & Management of hyperprolactinemia. Canadian Medical Association CMAJ .Sept.16, 2003;169(6)

Page 2 of 4 **Referred **Mr. David Adjei Adu** sample as received
Bsc.Biomedical Scientists



Ms. ABIR ABDUL MALIK

PID NO: P36180007457

Age: 20 Year(s) Sex: Female

Reference:

Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180107742

Registered On:

02/07/2018 03:29 PM

Collected On:

02/07/2018

Reported On:

03/07/2018 10:39 AM

Investigation	Observed Value	Unit	Biological Reference Interval
---------------	----------------	------	-------------------------------

HCG Beta Subunit, Serum

HCG Beta Subunit
(Serum,CMIA)

< 2.0

mIU/mL

Less than 5.3

Post LMP Period

-

Pregnant Woman - Weeks of gestation(Gest.)	Weeks post Last Menstrual Period (LMP)	Range(mIU/ml)
1.3 to 2	3.3 to 4	16 to 156
2 to 3	4 to 5	101 to 4,870
3 to 4	5 to 6	1,110 to 31,500
4 to 5	6 to 7	2,560 to 82,300
5 to 6	7 to 8	23,100 to 151,000
6 to 7	8 to 9	27,300 to 233,000
7 to 11	9 to 13	20,900 to 291,000
11 to 16	13 to 18	6,140 to 103,000
16 to 21	18 to 23	4,720 to 80,100
21 to 39	23 to 41	2,700 to 78,100

Page 3 of 4 **Referred Mr. David Adjei Adu the sample as received
Bsc.Biomedical Scientists



Ms. ABIR ABDUL MALIK

PID NO: P36180007457

Age: 20 Year(s) Sex: Female

Reference:

Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180107742

Registered On:

02/07/2018 03:29 PM

Collected On:

02/07/2018

Reported On:

03/07/2018 10:39 AM

Investigation

25 Hydroxy (OH) Vit D
(Serum,CMLA)

Observed Value

23.80

Unit

ng/mL

Biological Reference Interval

Deficiency: < 10
Insufficiency: 10-30
Sufficiency: 30-100
Hypervitaminosis: > 100

Interpretation :

1. Vitamin D is a fat soluble vitamin and exists in two main forms as cholecalciferol(vitamin D3) which is synthesized in skin from 7-dehydrocholesterol in response to sunlight exposure & Ergocalciferol(vitamin D2) present mainly in dietary sources.Both cholecalciferol & Ergocalciferol are converted to 25(OH)vitamin D in liver.
2. Testing for 25(OH)vitamin D is recommended as it is the best indicator of vitamin D nutritional status as obtained from sunlight exposure & dietary intake. For diagnosis of vitamin D deficiency it is recommended to have clinical correlation with serum 25(OH)vitamin D, serum calcium, serum PTH & serum alkaline phosphatase.
3. During monitoring of oral vitamin D therapy- suggested testing of serum 25(OH)vitamin D is after 12 weeks or 3 mths of treatment. However, the required dosage of vitamin D supplements & time to achieve sufficient vitamin D levels show significant seasonal(especially winter) & individual variability depending on age, body fat, sun exposure, physical activity ,genetic factors(especially variable vitamin D receptor responses), associated liver or renal disease, malabsorption syndromes and calcium or magnesium deficiency influencing the vitamin D metabolism Vitamin D toxicity is known but very rare.kindly correlate clinically, repeat with fresh sample if indicated.

Associated Test Profile :

- For diagnosis of vitamin D deficiency it is recommended to have clinical correlation with serum 25(OH)vitamin D and serum PTH.An inverse relationship exists between PTH and 25(OH)D levels, Parathyroid hormone levels start to rise at 25(OH)D levels below 31 ng/mL & usually decrease after the correction of vitamin D insufficiency.Thus, restoration of PTH and 25(OH) D levels to normalcy after adequate vitamin D replacement therapy is a useful monitoring strategy.
- As a holistic & scientific approach for diagnosis and optimal treatment for vitamin D deficiency, Vitamin D plus profile (25 Hydroxy(OH) Vit D and PTH) is suggested.

-- End of Report --

Page 4 of 4 **Referred Mr. David Adjei Adu the sample as received
Bsc.Biomedical Scientists