NAME ADE SEHEED ABDULA

REFERENCE: GILEAD MEDICAL

PID NO: P36160004054

VID: 36180104191

DATE: 21/05/2018

Sample Collected At: Ghana

AGE: 41Years(s) Sex: Male

Metropolis Healthcare (Ghana) Ltd.

Mantoux test

Negative (2) TU PPD)

mm

Negative: 0-4 (10 Indeterminate: 5-9

Positive: > 10

INTERPRETATION:

 The Mantoux tuberculin skin test (TST) is the standard method of an intradermal injection of 0.1 ml of tuberculin purified

protein derivative (PPD) into the inner surface of the forearm. The skin test reaction should be read between 48 and 72

hours after administration. A patient who does not return within 72 hours will need to be rescheduled for another skin test.

 The reaction is measured in millimeters of the induration (palpable, raised, hardened area or swelling) while erythema

(redness) is excluded & not measured. The diameter of the indurated area is measured across the forearm (perpendicular to

the long axis). An induration of 5 or more millimeters may be considered positive in HIV-infected persons, Persons who are

immunosuppressed for other reasons (e.g., taking the equivalent of >15 mg/day of prednisone for 1 month or longer, taking

TNF-a antagonists), patients with organ transplants.

 Positive mantoux test suggests delayed hypersensitivity reaction(type IV cell mediated immunity) to tuberculin protein, which in turn may be due to previous TB infection, BCG vaccination, exposure to atypical &

nontuberculosis mycobacteria or Latent

& Active TB, hence clinical correlation is necessary.

False-negative reactions may be seen with the following Cutaneous anergy (inability to react to skin tests because of a

weakened immune system), recent TB infection (within 8-10 weeks of exposure), young age (less than 6 months), viral

infections (like measles, varicella, influenza, EBV), very old TB infection (many years) resulting in boosted reaction,

Overwhelming TB disease like miliary TB.

End of Report-

DAVID ADJEI ADU

Bsc. Biomedical Scientists Advances Dip. (CMH)



INNER HEALTH REVEALED Metropolis Healthcare (Ghana) Ltd.

1st Floor, Pyramid Building, Above Stanbic Bank, Near Nima Traffic Light, Opposite Nima Police Station, Ring Road, Accra, Ghana.

Tel: +233 543028001 / +233 543026795. Email: info.ghana@metropolisafrica.com Website: www.metropolisafrica.com

Collection Centre:
Shop No. 2 Ground Floor, Opposite Korle-Bu Administration Gate, Rock Chemist Building, Korle-Bu, Accra, Ghana. Tel: + 233 241714323

Metropolis services are also available at: International: India . Sri Lanka . UAE . South Africa . Kenya . Mauritius