



**Gilead Medical & Dental Centre**

[info@gileadmedical.com](mailto:info@gileadmedical.com)

House of Balm, C896/3, Kanda Highway, Accra-Ghana

Tel:0302227196

[www.gileadmedicalgh.com](http://www.gileadmedicalgh.com)

Name: Lawrence Sampson Amartey

Age: 47yrs

Gender: Male

Clinical History:

Type of Investigation: Abdominal Ultrasound

Investigation Date:09/02/2018

### **ULTRASOUND REPORT**

#### **Abdomen**

The liver is of average size, measuring 14.2cm MCL with homogenous echopattern. No focal mass or diffuse lesion noted. No intra or extrahepatic bile duct dilatation.

The gallbladder shows normal outline and wall thickness. No gallstones, sludge or mass seen. No pericholecystic fluid seen. Absent sonographic Murphy sign.

The visualized portions of the pancreas is of normal sonographic appearance. No pancreatic calcification.

The spleen is normal in size and echopattern measuring 8.5cm. No focal mass or diffuse lesion

Both kidneys are of average size measuring, right (10.6x4.3cm) and left (10.7x4.6cm) respectively with good corticomedullary differentiation and normal renal sinus. No focal loss of renal parenchyma. No hydronephrosis, calculi or focal mass noted.

No ascites, abdominal mass or lymphadenopathy. There is normal abdominal aorta and IVC.

#### **Pelvic**

The prostate measures (4.7x3.9x2.6cm) approximately = 25.7cm<sup>3</sup> and has homogeneous echo pattern. No focal mass or calcifications noted.

The urinary bladder wall is not thickened. The Urinary bladder appears normal. No focal mass or calcifications were seen in it.

Pre-void bladder volume = 327.0 cm<sup>3</sup>

Post void residual bladder volume = 10.6 cm<sup>3</sup>

#### **Impression:**

- An average size prostate of size 25g.
- Insignificant Post void residual bladder volume
- Normal abdominal u/s scan

**Sonographer:** Rosemond Adu-Sarkodie

