

Gilead Medical & Dental Centre

info@gileadmedical.com
House of Balm, C896/3, Kanda Highway, Accra-Ghana
Tel:0302227196
www.gileadmedicalgh.com

Name: Elizabeth Asare-Boateng Age: 51yrs Gender: Female

Clinical History: Left Iliac fossa tenderness R/O an adnexal mass

Type of Investigation: Abdomino-Pelvic Ultrasound Investigation Date: 08/06/2018

ULTRASOUND REPORT

Pelvic

No sonographic evidence of adnexal mass.

The uterus is anteverted, homogeneous in echopattern and of average size measuring (7.8x4.9x3.9) cm. No focal mass noted. Normal endometrial stripe thickness

No fluid collection in the pouch of Douglas.

The ovaries were not visualized (? atrophy)

No adnexal mass noted.

The urinary bladder is normal with no calculus or masses noted.

Abdomen

The gallbladder shows normal outline and wall thickness but contains an echogenic focus at the neck measuring 11.4mm in diameter. No sludge or pericholecystic fluid seen. Absent sonographic Murphy sign.

The liver is of average size, measuring 14.9cm MCL with homogenous echopattern. No focal mass or diffuse lesion noted. No intra or extrahepatic bile duct dilatation.

The visualized portions of the pancreas is of normal sonographic appearance. No pancreatic calcification.

The spleen is normal in size and echopattern measuring 7.9cm. No focal mass or diffuse lesion

Both kidneys are of average size measuring, right (11.1x4.9) and left (11.2x6.3) cm respectively with good corticomedullary differentiation and normal renal sinus. No focal loss of renal parenchyma. No hydronephrosis, calculi or focal mass noted.

No ascites, abdominal mass or lymphadenopathy. There is normal abdominal aorta and IVC.

Impre	ession:		
•	Cholelithiasis Normal Pelvic Ultrasound.		
Sono	grapher: Rosemond Adu-Sarko	odie	