



Gilead Medical & Dental Centre

info@gileadmedical.com

House of Balm, C896/3, Kanda Highway, Accra-Ghana

Tel:0302227196

www.gileadmedicalgh.com

Name: Esther Afua Hanson

Age:62

Gender: Female

Clinical History:

Type of Investigation: Doppler Ultrasound

Investigation Date:05/12/2017

ULTRASOUND REPORT

There is an anechoic fluid collection at the left popliteal fossa measuring about 2.9x2.7x1.3cm (vol=5.9ml)

Noted from the ankle to the foot of both lower limbs are multiple dilated fluid channels with a cobblestone appearance

Also noted at the upper thigh of both lower limbs are multiple (about 3 each) benign-appearing axillary lymph nodes.

The common and superficial femoral, popliteal and tibia arteries are of normal caliber.

The vessel walls are of normal thickness. There are no artero-sclerotic changes. They show normal high resistance triphasic flow waveform.

The flow velocities are normal. The accompanying veins are grossly normal. The soft tissues are grossly normal.

Impression:

- Features Suggestive Left Baker's (Popliteal) Cyst
- Features Suggestive of Bilateral Lymph edema of the foot.
- Normal Arterial Doppler study of both lower limbs

Sonographer: Rosemond Adu-Sarkodie