

## Management of postpartum anaemia

**IV iron** should be considered for severe anaemia as oral iron is known to be slow and unreliable. In a randomised controlled study of 44 women with postpartum anaemia, significantly higher mean haemoglobin and ferritin levels from baseline were achieved for patients on **IV iron sucrose** (200 mg x 2, 48 hours apart) in comparison to those on oral iron (mean Hb day 5: IV vs oral iron, 2.5 vs 0.7 gm/dl - day 14 Hb: 3.8 vs 1.5 gm/dl,  $p = <0.01$  for both periods)<sup>31</sup>. Comparable results for IV iron sucrose were reported in 2 similar trials (mean Hb 2.8 & 3.1, both day 14)<sup>32,33</sup>. These increases in Hb from baseline with IV iron exceed the expected rise after a 2U blood transfusion<sup>32</sup>. The level of life-threatening adverse drug events of IV iron preparations is now very low, varying from 0.6 to 3.3 per million, depending on the iron preparation (FDA data)<sup>34</sup>.

**Erythropoiesis-stimulating agents (ESAs)** should be administered **together with IV iron** in life-threatening anaemia to further accelerate erythropoiesis. A **once weekly EPO dosage of 600 IU/kg subcutaneously (e.g. 40,000 IU for a 66kg patient)** is being increasingly used and found to be satisfactory in critically ill anaemic patients<sup>35,36</sup>. An **EPO dosage of 300 IU/kg x 3 weekly together with IV iron (200mg x 3 weekly)** has also proved efficacious for postpartum anaemia<sup>37</sup>. Augment with **vitamin B-12 and folic acid**.

**Check oxygen saturations:** Give **100% oxygen** if necessary (no contraindications for 48-72 hrs of use). Use **microsampling techniques** to conserve blood (e.g. HemoCue), as well as **paediatric sample tubes**. If bleeding continues consider reinfusing washed drain fluid.

**Hyperbaric oxygen therapy:** Option in life-threatening anaemia<sup>38</sup>. (0151 648 8000 [24 hrs] for suitable and available centres.)

### References:

1. Al RA, Unlubilgin E, Kandemir O, Yalvac S, Cakir L, Haberal A. Intravenous versus oral iron for treatment of anemia in pregnancy: a randomized trial. *Obstet Gynecol* 2005; 106: 1335-40.
2. al-Momen AK, al-Meshari A, al-Nuaim L, Saddique A, Abotalib Z, Khashoggi T, Abbas M. Intravenous iron sucrose complex in the treatment of iron deficiency anemia during pregnancy. *Eur J Obstet Gynecol Reprod Biol* 1996; 69: 121-4.
3. Breyman C, Visca E, Huch R, Huch A. Efficacy and safety of intravenously administered iron sucrose with and without adjuvant recombinant human erythropoietin for the treatment of resistant iron-deficiency anemia during pregnancy. *Am J Obstet Gynecol* 2001; 184: 662-667.
4. Kalu E, Wayne C, Croucher C, Findley I, Manyonda I. Triplet pregnancy in a Jehovah's Witness: recombinant human erythropoietin and iron supplementation for minimising the risks of excessive blood loss. *BJOG* 2002; 109: 723-725.
5. McDonald SJ, Middleton P. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *Cochrane Database Syst Rev* 2008 Apr 23; (2): CD004074.
6. Hofmeyr GJ, Abdel-Aleem H, Abdel-Aleem MA. Uterine massage for preventing postpartum haemorrhage. *Cochrane Database Syst Rev* 2008 Jul 16; (3): CD006431.
7. Davies GA, Tessier JL, Woodman MC, Lipson A, Hahn PM. Maternal hemodynamics after oxytocin bolus compared with infusion in the third stage of labor: a randomized controlled trial. *Obstet Gynecol* 2005; 105: 294-9.
8. Choy CMY, Lau WC, Tam WH, Yuen PM. A randomised controlled trial of intramuscular syntometrine and intravenous oxytocin in the management of the third stage of labour. *BJOG* 2002; 109: 173-177.
9. Blum J, Winikoff B, Raghavan S, Dabash R, Ramadan MC, Dilbaz B, Dao B, Durocher J et al. Treatment of post-partum haemorrhage with sublingual misoprostol versus oxytocin in women receiving prophylactic oxytocin: a double-blind, randomised, non-inferiority trial. *Lancet* 2010; 375: 217-23.
10. O'Brien P P, El-Refaey H, Gordon A, Geary M, Rodeck CH. Rectally administered misoprostol for the treatment of postpartum hemorrhage unresponsive to oxytocin and ergometrine: a descriptive study. *Obstet Gynecol* 1998; 92: 212-214.
11. Adekanmi OA, Purnessur S, Edwards G, Barrington JW. Intrauterine misoprostol for the treatment of severe recurrent atonic secondary postpartum haemorrhage. *BJOG* 2001; 108: 541-542.
12. Condous GS, Arulkumaran S, Symonds I, Chapman R, Sinha A, Razvi K. The "tamponade test" in the management of massive postpartum hemorrhage. *Obstet Gynecol* 2003; 101: 767-772.
13. Danso D, Reginald PW. Internal Uterine Tamponade. In: A Textbook of Postpartum Haemorrhage. Ch 28, p 263-67. Ed Christopher B-Lynch et al. Sapiens Publishing. 2006.
14. Doumouchtisis SK, Papageorgiou AT, Vernier C, Arulkumaran S. Management of postpartum hemorrhage by uterine balloon tamponade: prospective evaluation of effectiveness. *Acta Obstet Gynecol Scand* 2008; 87: 849-55.
15. Yoong W, Ray A, Phillip SA. Balloon tamponade for postpartum vaginal lacerations in a woman refusing blood transfusion. *Int J Gynaecol Obstet* 2009; 106: 261.
16. Miller S, Hamza S, Bray EH, Lester F, Nada K, Gibson R, Fathalla M, Mourad M, Fathy A, Turan JM, Dau KQ, Nasshar I, Elshair I, Hensleigh P. First aid for obstetric hemorrhage: the pilot study of the non-pneumatic anti-shock garment in Egypt. *BJOG* 2006; 113: 424-29.
17. Franchini M, Franchini M, Bergamini V, Salvagno GL, Montagnana M, Lippi G. A critical review on the use of recombinant factor VIIa in life-threatening obstetric postpartum hemorrhage. *Semin Thromb Hemost* 2008; 34: 104-12.
18. Pepas LP, Arif-Adib M, Kadir RA. Factor VIIa in puerperal hemorrhage with disseminated intravascular coagulation. *Obstet Gynecol* 2006; 108: 757-61.
19. Lewis NR, Bruner P, Lemire SJ, Kaufman RM. Failure of recombinant factor VIIa to correct the coagulopathy in a case of severe postpartum hemorrhage. *Transfusion* 2009; 49: 689-95.
20. Weinkove R, Rangarajan S. Fibrinogen concentrate for acquired hypofibrinogenemic states. *Transfus Med* 2008; 18: 151-57.
21. Tanaka KA, Taketomi T, Szlam F, Calatzis A, Levy JH. Improved clot formation by combined administration of activated factor VII (NovoSeven) and fibrinogen (Haemocomplettan P). *Anesth Analg* 2008; 106: 732-8.
22. Bruce D, Nokes TJ. Prothrombin complex concentrate (Beriplex P/N) in severe bleeding: experience in a large tertiary hospital. *Crit Care* 2008; 12: R105.
23. Gai M-y, Wu L-f, Su Q-f, Tatsumoto K. Clinical observation of blood loss reduced by tranexamic acid during and after caesarian section: A multi-center, randomized trial. *Eur J Obstet Gynecol Reprod Biol* 2004; 112: 154-157.
24. Moriarty KT, Premila S, Bulmer PJ. Use of FloSeal haemostatic gel in massive obstetric haemorrhage: a case report. *BJOG* 2008; 115: 793-95.
25. Whiteside JL, Asif RB, Novello RJ. Fibrin Sealant for Management of Complicated Obstetric Lacerations. Case Reports. *Obstet & Gynecol* 2010; 115: 403-404.
26. Danso D, Reginald P. Combined B-Lynch suture with intrauterine balloon catheter triumphs over massive postpartum haemorrhage. *BJOG* 2002; 109: 963.
27. Ghezzi F, Cromi A, Uccella S, Rao L, Bolis P, Surbek D. The Hayman technique: a simple method to treat postpartum haemorrhage. *BJOG* 2007; 114: 362-5.
28. Choji K, Shimizu T. Embolization. In: A Textbook of Postpartum Haemorrhage. Ch 30, p 277-85. Ed Christopher B-Lynch et al. Sapiens Publishing. 2006.
29. Sullivan I, Faulds J, Ralph C. Contamination of salvaged maternal blood by amniotic fluid and fetal red cells during elective Caesarean section. *Br J Anaesth* 2008; 101: 225-9.
30. Catling S. Blood conservation techniques in obstetrics: a UK perspective. *Int J Obstet Anesth* 2007; 16: 241-49.
31. Bhandal N, Russell R. Intravenous versus oral iron therapy for postpartum anaemia. *BJOG* 2006; 113: 1248-52.
32. Wågström E, Akesson A, Van Rooijen M, Larson B, Bremme K. Erythropoietin and intravenous iron therapy in postpartum anaemia. *Acta Obstet Gynecol Scand* 2007; 86: 957-62.
33. Broche DE, Gay C, Armand-Branger S, Grangeasse L, Terzibachian JJ. Acute postpartum anaemia. Clinical practice and interest of intravenous iron. *Gynecol Obstet Fertil* 2004; 32: 613-19.
34. Chertow GM, Mason PD, Vaage-Nilsen O, Ahlmén J. Update on adverse drug events associated with parenteral iron. *Nephrol Dial Transplant* 2006; 21: 378-82.
35. Corwin HL, Gettinger A, Pearl RG, Fink MP, Levy MM, Shapiro MJ, Corwin MJ, Colton T. Efficacy of recombinant human erythropoietin in critically ill patients: a randomized controlled trial. *JAMA* 2002; 288: 2827-35.
36. Georgopoulos D, Matamis D, Routsi C, Michalopoulos A, Maggina N, Dimopoulos G, et al. Recombinant human erythropoietin therapy in critically ill patients: a randomized study. *Crit Care* 2005; 9: R508-15.
37. Breyman C, Richter C, Hüttner C, Huch R, Huch A. Effectiveness of recombinant erythropoietin and iron sucrose vs. iron therapy only, in patients with postpartum anaemia and blunted erythropoiesis. *Eur J Clin Invest* 2000; 30: 154-161.
38. McLoughlin PL, Cope TM, Harrison JC. Hyperbaric oxygen therapy in the management of severe acute anaemia in a Jehovah's Witness. *Anaesthesia* 1999; 54: 891-895.

This document has been reviewed by consultants in obstetrics, gynaecology, anaesthesia and haematology (including experts in haemostasis). It reflects current clinical and scientific knowledge and is subject to change. The strategies are not intended as an exclusive guide to treatment. Good clinical judgement, taking into account individual circumstances, may require adjustments.

**For questions, comments, or information as to the availability of the above medications/treatment options please contact:**  
**Hospital Information Services for Jehovah's Witnesses (054-433-9541, 054-433-9542 (24 hrs); hid.gh@jw.org)**