



Mr. SITSO KWASI KORNU

PID NO: P36180008163

Age: 33 Year(s) Sex: Male

Reference:

Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180108469

Registered On:

10/07/2018 05:00 PM

Collected On:

10/07/2018

Reported On:

10/07/2018 10:15 PM

Investigation	Observed Value	Unit	Biological Reference Interval
Thyroid panel-1 (T3/T4/TSH) (Serum)			
T3 (Total) (CMIA)	0.43	ng/dL	Male: <= 3 Days : 100-740 Male: 3 - 365 Days : 105 - 245 Male: 1 - 5 Years : 105 - 269 Male: 5 - 10 Years : 94 - 241 Both: > 10 Years : 0.52-1.85 : 0.52 - 1.85
T4 (Total) (CMIA)	3.39	µg/dL	4.4-10.8
TSH(Ultrassensitive)	0.73	µIU/mL	0.45-4.5

INTERPRETATION

TSH	T3 / FT3	T4 / FT4	Suggested Interpretation for the Thyroid Function Tests Pattern
Within Range	Decreased	Within Range	• Isolated Low T3-often seen in elderly & associated Non-Thyroidal illness. In elderly the drop in T3 level can be upto 25%.
Raised	Within Range	Within Range	•Isolated High TSHespecially in the range of 4.7 to 15 mIU/ml is commonly associated with Physiological & Biological TSH Variability. •Subclinical Autoimmune Hypothyroidism •Intermittent T4 therapy for hypothyroidism •Recovery phase after Non-Thyroidal illness"
Raised	Decreased	Decreased	•Chronic Autoimmune Thyroiditis •Post thyroidectomy,Post radioiodine •Hypothyroid phase of transient thyroiditis"
Raised or within Range	Raised	Raised or within Range	•Interfering antibodies to thyroid hormones (anti-TPO antibodies) •Intermittent T4 therapy or T4 overdose •Drug interference- Amiodarone, Heparin,Beta blockers,steroids, anti-epileptics"
Decreased	Raised or within Range	Raised or within Range	•Isolated Low TSH -especially in the range of 0.1 to 0.4 often seen in elderly & associated with Non-Thyroidal illness •Subclinical Hyperthyroidism •Thyroxine ingestion"
Decreased	Decreased	Decreased	•Central Hypothyroidism •Non-Thyroidal illness •Recent treatment for Hyperthyroidism (TSH remains suppressed)"
Decreased	Raised	Raised	•Primary Hyperthyroidism (Graves' disease),Multinodular goitre, Toxic nodule •Transient thyroiditis:Postpartum, Silent (lymphocytic), Postviral (granulomatous,subacute, DeQuervain's),Gestational thyrotoxicosis with hyperemesis gravidarum"
Decreased or within Range	Raised	Within Range	•T3 toxicosis •Non-Thyroidal illness

References: 1. Interpretation of thyroid function tests. Dayan et al. THE LANCET • Vol 357 • February 24, 2001
2. Laboratory Evaluation of Thyroid Function, Indian Thyroid Guidelines, JAPI, January 2011,vol. 59

Reports to follow - Kindly await following pending reports :

Investigation :	Status
Cortisol, Serum	Pending

-- End of Report --



Mr. SITSO KWASI KORNU

PID NO: P36180008163
Age: 33 Year(s) Sex: Male

Reference:
Sample Collected At:
GILEAD MEDICAL & DENTAL CENTER
HOUSE NO BALB NO C896/3,KANDA
HIGHWAY NORTH RIDGE,ACCRA-
14911.
014911

VID: 36180108469
Registered On:
10/07/2018 05:00 PM
Collected On:
10/07/2018
Reported On:
10/07/2018 10:15 PM

<u>Investigation</u>	<u>Observed Value</u>	<u>Unit</u>	<u>Biological Reference Interval</u>
----------------------	-----------------------	-------------	--------------------------------------