

Gilead Medical & Dental Centre

info@gileadmedical.com
House of Balm, C896/3, Kanda Highway, Accra-Ghana
Tel:0302227196
www.gileadmedicalgh.com

Name: Mr. Henry Aryee Age: 46yrs Gender: Male

Clinical History:

Type of Investigation: Abdomino-Pelvic Ultrasound Investigation Date:28/12/2017

ULTRASOUND REPORT

Abdomen

The gallbladder shows normal outline and wall thickness but contains an echogenic foci measuring with posterior acoustic shadowing 12.7mm in diameter. No pericholecystic fluid seen. Absent sonographic Murphy sign.

The liver is of average size, measuring 15.0cm MCL with homogenous echopattern. No focal mass, surface nodularity or diffuse lesion noted. No intra or extrahepatic bile duct dilatation.

The visualized portions of the pancreas is of normal sonographic appearance. No pancreatic calcification.

The spleen was not visualized (? Auto-splenectomy)

Both kidneys are of average size measuring, right (11.1x4.2) and left (11.1x4.5) cm respectively with good corticomedullary differentiation and normal renal sinus. No focal loss of renal parenchyma. No hydronephrosis, calculi or focal mass noted.

No ascites, abdominal mass or lymphadenopathy. There is normal abdominal aorta and IVC.

Pelvic

The prostate measures (4.6x3.1x2.7) cm, approximately =20.8cm³ and has homogeneous echo pattern. No focal mass or calcifications noted.

The urinary bladder wall is not thickened. The Urinary bladder appears normal. No focal mass or calcifications were seen in it.

Pre-void bladder volume=317.76cm³

Post void residual bladder volume = 39.7cm³

.

Impression:

- Cholelitiaisis
- An Average Size Prostate of size 20g
 Insignificant Post Void Residual Bladder volume of 39.7 cm³

Sonographer: Rosemond Adu-Sarkodie