

PID NO: P36180015617

Age: 58 Year(s) Sex: Female

Reference:

Sample Collected At: GILEAD MEDICAL & DENTAL CENTER HOUSE NO BALB NO C896/3,KANDA HIGHWAY NORTH RIDGE,ACCRA-14911. 014911 VID: 36180116247

Registered On: 28/09/2018 05:53 PM Collected On: 28/09/2018

Reported On: 09/10/2018 05:30 PM

InvestigationObserved ValueUnitBiological Reference IntervalCA-125 (Cancer Antigen-125)6.41U/mL0-36

(Serum)

Interpretation:

- 1. CA 125 is a glycoprotein normally expressed in coelomic epithelium, which lines body cavities and envelopes the ovaries.
- CA 125 levels are elevated in about 85 percent of women with ovarian cancer (especially serous epithelial tumours), but in only 50 percent of those with stage I disease.
- 3. Multiple benign disorders like Menstruation, pregnancy, fibroids, ovarian cysts, pelvic inflammation, cirrhosis, ascites, pleural and pericardial effusions, endometriosis also are associated with CA 125 elevations.
- 4. Levels above which benign diseases are considered unlikely are 200U/ml in premenopausal & 35 u/ml for postmenopausal women.

Reference: Greg.L.Perkin. et.al. Serum Tumor Markers. American family physicians sep.2003 vol.68 no.6.

Associated Test: HE4 assay is a new test which also can be used for therapeutic monitoring as well as for risk stratification of harboring Epithelial Ovarian Cancer (ROMA value) in early stages.



PID NO: P36180015617

Age: 58 Year(s) Sex: Female

Reference:

Sample Collected At:

GILEAD MEDICAL & DENTAL CENTER HOUSE NO BALB NO C896/3,KANDA HIGHWAY NORTH RIDGE,ACCRA-

14911. **014911**

VID: 36180116247

Registered On: 28/09/2018 05:53 PM Collected On:

28/09/2018

Reported On: 09/10/2018 05:30 PM

PAP SMEAR EXAMINATION



INTERNATIONAL & NATIONAL SUBSPECIALITY PATHOLOGY

Breast Pathology

Dermatopathology

Gastrointestinal Pathology

Genitourinary Pathology

Gynecologic Pathology

Head & Neck Pathology

Hematolymphoid Pathology

Hepatobillary Pathology

Neuropathology

Paediatric & Perinatal Pathology

Renal Pathology

Soft tissue Pathology

Transplant Pathology (Renal & Hepatic)

Senior Consultant & Vice President Operations

Dr Kirti Chadha

Global Reference Laboratory Faculty

Senior Consultants

Dr Anuradha Murthy Dr Vikas Kavishwar Dr Sushma Gurwale Dr Barodawala S.M

Consultants Dr Kunial Lila

Dr Tejal Shah Dr Shital Munde Dr Aditi Raj Dr Ramrao Nilkanthe

Case Summary

CASE NO. 18 ML G - 14038

SPECIMEN CONVENTIONAL PAP SMEAR

DIAGNOSIS Negative For Intraepithelial Lesion or Malignancy (NILM)

ADVICE / COMMENT Reactive cellular changes associated with inflammation.

Kindly repeat PAP smear examination after treatment of

local inflammation.

Clinical Notes -

Gross Examination Received one unstained smear (Evelyn)

MICROSCOPIC EXAMINATION

Specimen Adequacy satisfactory for evaluation; endocervical/transformation

zone component present.

Superficial cells Present Intermediate cells Present

Deep parabasal/ Basal cells

Parabasal cells Present

Metaplastic squamous cells Present

Endocervical cells Present

Others -

Inflammation Severe

ORGANISMS

Doderlein bacilli Trichomonas Vaginilis Fungal organisms Others -

EPITHELIAL CELL Not Detected

ABNORMALITIES

GLANDULAR CELLS SQUAMOUS CELLS -

Note

"Cervical cytology is a screening test and has associated false negative and false positive results. Regular sampling and follow up is recommended".

Processing Method: Manual. Staining: Papanicolaou method

Dr. Aditi Raj

DCP ,M.D (Pathology),

Page 2 of 4 Associate Consultant Pathologist



PID NO: P36180015617

Age: 58 Year(s) Sex: Female

Reference:

Sample Collected At: GILEAD MEDICAL & DENTAL CENTER HOUSE NO BALB NO C896/3,KANDA HIGHWAY NORTH RIDGE,ACCRA-14911.

The smears are reported using the Bethesda System for Reporting Cervical Cytology (2014)

New Cervical Cancer Screening Recommendations from the U.S. Preventive Services Task Force and the American Cancer Society/American Society for Colposcopy and Cervical

014911

VID: 36180116247

Registered On: 28/09/2018 05:53 PM Collected On: 28/09/2018

Reported On: 09/10/2018 05:30 PM

Clinical Application:



INTERNATIONAL & NATIONAL SUBSPECIALITY PATHOLOGY

Breast Pathology

Dermatopathology

Gastrointestinal Pathology

Genitourinary Pathology

Gynecologic Pathology

Head & Neck Pathology

Hematolymphoid Pathology

Hepatobillary Pathology

Neuropathology

Paediatric & Perinatal Pathology

Renal Pathology

Soft tissue Pathology

Transplant Pathology (Renal & Hepatic)

Senior Consultant & Vice President Operations

Dr Kirti Chadha

Global Reference Laboratory Faculty

Senior Consultants

Dr Anuradha Murthy Dr Vikas Kavishwar

Di Vinas Navistiwai

Dr Sushma Gurwale

Dr Barodawala S.M

Consultants

Dr Kunjal Lila Dr Tejal Shah

Dr Shital Munde

Dr Aditi Raj

Dr Ramrao Nilkanthe

			ety for Colposcopy and Cervical ch 15, 2012, issue of Annals of Internal
Population	USPSTF	ACS/ASCCP/ASCP	
Younger than 21 years	Recommends against screening. Grade: D recommendation.	Women should not be screened regardless of the age of sexual initiation or other risk factors.	
21–29 years	Recommends screening with cytology every 3 years. Grade: A recommendation.	Screening with cytology alone every 3 years is recommended.	
30–65 years	Recommends screening with cytology every 3 years or for women who want to lengthen the screening interval, screening with a combination of cytology and HPV testing every 5 years. Grade: A recommendation.	Screening with cytology and HPV testing ("co-testing") every 5 years (preferred) or cytology alone every 3 years (acceptable) is recommended.	
Older than 65 years	Recommends against screening women who have had adequate prior screening and are not otherwise at high risk for cervical cancer. Grade: D recommendation.	Women with evidence of adequate negative prior screening and no history of CIN2+ within the last 20 years should not be screened. Screening should not be resumed for any reason, even if a woman reports having a new sexual partner.	
After hysterectomy	Recommends against screening in women who have had a hysterectomy with removal of the cervix and who do not have a history of a high-grade precancerous	Women of any age following a hysterectomy with removal of the cervix who have no history of CIN2+ should not be screened for vaginal cancer.	

Adith Raj

Dr. Aditi Raj DCP ,M.D (Pathology), Page 3 of 4 Associate Consultant Pathologist



PID NO: P36180015617

Age: 58 Year(s) Sex: Female

Reference:

Sample Collected At: GILEAD MEDICAL & DENTAL CENTER HOUSE NO BALB NO C896/3,KANDA HIGHWAY NORTH RIDGE,ACCRA-14911.

014911

VID: 36180116247

Registered On: 28/09/2018 05:53 PM Collected On: 28/09/2018

Reported On: 09/10/2018 05:30 PM

METROPOLIS
HIST@XPERT
GLOBAL EXPERTISE IN SUB SPECIALTY SOLUTIONS

INTERNATIONAL & NATIONAL SUBSPECIALITY PATHOLOGY

Breast Pathology

Dermatopathology

Gastrointestinal Pathology

Genitourinary Pathology

Gynecologic Pathology

Head & Neck Pathology

Hematolymphoid Pathology

Hepatobillary Pathology

Neuropathology

Paediatric & Perinatal Pathology

Renal Pathology

Soft tissue Pathology

Transplant Pathology (Renal & Hepatic)

Senior Consultant & Vice President Operations

Dr Kirti Chadha

Global Reference Laboratory Faculty

Senior Consultants

Dr Anuradha Murthy

Dr Vikas Kavishwar

Dr Sushma Gurwale

Dr Barodawala S.M

Consultants

Dr Kunjal Lila

Dr Tejal Shah

Dr Shital Munde

Dr Aditi Raj

Dr Ramrao Nilkanthe

	lesion (ie, CIN 2 or 3) or cervical cancer. Grade: D recommendation	Evidence of adequate negative prior screening is not required. Screening should not be resumed for any reason, including if a woman reports having a new sexual partner.
HPV vaccinated	Women who have been vaccinated should continue to be screened.	Recommended screening practices should not change on the basis of HPV vaccination status.

-- End of Report --

Adith Ray

Dr. Aditi Raj DCP ,M.D (Pathology), Page 4 of 4 Associate Consultant Pathologist