

Factor Analysis Examples

PC method vs ML method on stock data

STAT 32950-24620

Spring 2023 (3/30, wk2)

1 / 28

Orthogonal Factor Model

$$X = \mu + LF + \varepsilon$$

- $X = [X_1 \cdots X_p]'$
 - Observable, random, $E(X) = \mu$
- $F = [F_1 \cdots F_m]'$
 - Latent factors, unobservable, random
- $L = L_{p \times m}$
 - Factor loading matrix, parameters, to be estimated

2 / 28

Orthogonal factor model assumptions

Assumptions on common factors

- $E(F) = 0_m$
- $Cov(F) = I_m$

Assumptions on errors (specific factors)

- $E(\varepsilon) = 0_p$,

Assumptions on covariance between common and specific factors

- $Cov(\varepsilon) = \text{diag}\{\psi_1, \dots, \psi_p\} = \Psi$
- $Cov(F, \varepsilon) = E(F\varepsilon') = 0_{m \times p}$
- $Cov(\varepsilon, F) = E(\varepsilon F') = 0_{p \times m}$

3 / 28

Covariance structure in factor model

Key structure assumption of the model:

$$\Sigma = LL' + \Psi$$

where

$$\Sigma = \text{Cov}(X)$$

under the factor relation

$$X = \mu + LF + \varepsilon$$

4 / 28

FA parameter estimation methods

- By Principal Components
- By Maximum Likelihood

The Maximum Likelihood method is usually preferred.

5 / 28

Principal factor estimation method

$$L_m = [\sqrt{\lambda_1}e_1, \dots, \sqrt{\lambda_m}e_m], \quad m \leq p$$

$$\text{When } m = p: \quad \Sigma = L_p L_p'$$

$$\text{When } m < p: \quad \Sigma = L_m L_m' + \dots$$

In orthogonal factor model

$$X = \mu + LF + \varepsilon$$

PC method uses the approximation

$$\Sigma \approx L_m L_m' + \Psi$$

for factor model with $m < p$ factors.

6 / 28

Obtain the e-vector e_i for the L matrix

```
# Using original data (alternative: var. variance=1 later)
summary(princomp(stock), loading=T)
```

```
## Importance of components:
##               Comp.1  Comp.2  Comp.3  Comp.4  (
## Standard deviation  0.0368 0.02635 0.01585 0.01188 0.
## Proportion of Variance 0.5293 0.27133 0.09822 0.05518 0.
## Cumulative Proportion 0.5293 0.80059 0.89881 0.95399 1.
##
## Loadings:
##      Comp.1  Comp.2  Comp.3  Comp.4  Comp.5
## JPMorgan    0.223   0.625   0.326   0.663   0.118
## Citibank     0.307   0.570  -0.250  -0.414  -0.589
## WellsFargo   0.155   0.345           -0.497   0.780
## Shell        0.639  -0.248  -0.642   0.309   0.148
## Exxon        0.651  -0.322   0.646  -0.216
```

7 / 28

Obtain λ_i for the L matrix

```
rtev = princomp(stock)$sdev # =sqrt of e-values of cov(X)
sqrt(eigen(cov(stock))$values) # verify: same as above

## [1] 0.03698 0.02648 0.01593 0.01194 0.01090
rtev

##      Comp.1  Comp.2  Comp.3  Comp.4  Comp.5
## 0.03680 0.02635 0.01585 0.01188 0.01085
```

8 / 28

Two common factors (PC method)

Choose $m = 2$

Take the first 2 PC's

$$Y_i = e_i'X, \quad i = 1, 2.$$

On original data (using covariance matrix)

```
princomp(stock)$loading[,1:2]
```

```
##           Comp.1  Comp.2
## JPMorgan  0.2228  0.6252
## Citibank   0.3073  0.5704
## WellsFargo 0.1548  0.3445
## Shell      0.6390 -0.2479
## Exxon      0.6509 -0.3218
```

9 / 28

Factor loadings for $m=2$ (PC method)

$$L_2 = [\sqrt{\lambda_1}e_1, \sqrt{\lambda_2}e_2]$$

Scale the PC variables by $\sqrt{\lambda_i}$ to get common factor loadings:

```
L = cbind(rtev[1]*princomp(stock)$loading[,1],
          rtev[2]*princomp(stock)$loading[,2]); L
```


```
##           [,1]      [,2]
## JPMorgan  0.008200  0.016475
## Citibank   0.011309  0.015030
## WellsFargo 0.005697  0.009078
## Shell      0.023515 -0.006534
## Exxon      0.023955 -0.008481
```

10 / 28

Plot of variables on two factors (PC method)

```
plot(L,type="n",main="Var. loadings on 2 common factors")
text(L,labels=row.names(L),col=1:5)
```


Var. loadings on 2 common factors



11 / 28

Comparison: Plot of variables on PC1 and PC2

```
plot(princomp(stock)$loading[,1:2],type="n");
text(princomp(stock)$loading[,1:2],
     labels=(colnames(stock)),cex=.8,col=1:5)
```



12 / 28

Covariance structure estimation (PC method)

Model assumption

$$\Sigma = LL' + \Psi$$

PC estimation method:

Let

$$\hat{L} = \hat{L}_m$$

Then

$$\hat{\Sigma} \approx \hat{L}\hat{L}' + \hat{\Psi}$$

13 / 28

Estimate of LL'

Get $\hat{L}\hat{L}'$, the estimate of LL'

```
LLT= L*%t(L)
round(10000*LLT,3)
```

##	JPMorgan	Citibank	WellsFargo	Shell	Exxon
## JPMorgan	3.387	3.404	1.963	0.852	0.567
## Citibank	3.404	3.538	2.009	1.677	1.434
## WellsFargo	1.963	2.009	1.149	0.747	0.595
## Shell	0.852	1.677	0.747	5.957	6.187
## Exxon	0.567	1.434	0.595	6.187	6.458

14 / 28

Estimate of specific factor matrix Φ (PC method)

Get the estimate of $\Psi = Cov(\varepsilon) (\times 10^4)$

```
## get the specific factors
Psi = diag(cov(stock)) - diag(LLT)
round(10000*Psi,3)
```

##	JPMorgan	Citibank	WellsFargo	Shell	Exxon
##	0.946	0.849	1.091	1.268	1.199

Compared with sample variance $s_{ii} = s_i^2 (\times 10^4)$

```
## get sample variance
round(10000*diag(cov(stock)),3)
```

##	JPMorgan	Citibank	WellsFargo	Shell	Exxon
##	4.333	4.387	2.240	7.225	7.657

15 / 28

Estimate of communality h_i^2 (PC method)

Get the estimate of communality $h_i^2 = \ell_{i1}^2 + \dots + \ell_{im}^2 (\times 10^4)$,

the portion of $Var(X_i)$ explained by common factors,

which are diagonal elements of estimated LL' :

```
## get communality h_i^2
round(10000*diag(LLT),3)
```

##	JPMorgan	Citibank	WellsFargo	Shell	Exxon
##	3.387	3.538	1.149	5.957	6.458

16 / 28

%vaiance explained by each common factor (PC)

For $j = 1, \dots, m$.

$$\frac{\hat{\ell}_{1j}^2 + \dots + \hat{\ell}_{pj}^2}{s_{11} + \dots + s_{pp}} = \frac{(\sqrt{\hat{\lambda}_j} e_j)(\sqrt{\hat{\lambda}_j} e_j)'}{\text{tr}(\mathbf{S})} = \frac{\hat{\lambda}_j \|e_j\|^2}{\text{tr}(\mathbf{S})} = \frac{\hat{\lambda}_j}{\sum_{i=1}^p \hat{\lambda}_i}$$

```
eigen(cov(stock))$value[1:2]/sum(eigen(cov(stock))$value)
```

```
## [1] 0.5293 0.2713
```

17 / 28

The approximation $\Sigma \approx L_2 L_2' + \Psi (\times 10^4)$

```
round((LLT + diag(Psi))*10000,2)
```

```
##                JPMorgan Citibank WellsFargo Shell Exxon
## JPMorgan        4.33      3.40      1.96  0.85  0.57
## Citibank        3.40      4.39      2.01  1.68  1.43
## WellsFargo      1.96      2.01      2.24  0.75  0.59
## Shell           0.85      1.68      0.75  7.22  6.19
## Exxon           0.57      1.43      0.59  6.19  7.66
```

Sample $\Sigma (\times 10^4)$

```
round((cov(stock))*10000,2)
```

```
##                JPMorgan Citibank WellsFargo Shell Exxon
## JPMorgan        4.33      2.76      1.59  0.64  0.89
## Citibank        2.76      4.39      1.80  1.81  1.23
## WellsFargo      1.59      1.80      2.24  0.73  0.61
## Shell           0.64      1.81      0.73  7.22  5.08
## Exxon           0.89      1.23      0.61  5.08  7.66
```

18 / 28

Maximum loading with m=5 factors (PC method)

```
L5=
princomp(stock)$loading[,1:5]*%
+diag((princomp(stock)$sdev))
round(L5,4)
```

```
##                [,1]    [,2]    [,3]    [,4]    [,5]
## JPMorgan    0.0082  0.0165  0.0052  0.0079  0.0013
## Citibank    0.0113  0.0150 -0.0040 -0.0049 -0.0064
## WellsFargo  0.0057  0.0091 -0.0006 -0.0059  0.0085
## Shell       0.0235 -0.0065 -0.0102  0.0037  0.0016
## Exxon       0.0240 -0.0085  0.0102 -0.0026 -0.0010
```

19 / 28

Numerical rounding errors

We should have $L_5 L_5' = \Sigma$ in theory, but there are rounding errors...

```
round((L5%*t(L5))*10000,2)
```

```
##                JPMorgan Citibank WellsFargo Shell Exxon
## JPMorgan        4.29      2.73      1.57  0.63  0.88
## Citibank        2.73      4.34      1.78  1.80  1.22
## WellsFargo      1.57      1.78      2.22  0.73  0.60
## Shell           0.63      1.80      0.73  7.15  5.03
## Exxon           0.88      1.22      0.60  5.03  7.58
```

```
round((cov(stock))*10000,2)
```

```
##                JPMorgan Citibank WellsFargo Shell Exxon
## JPMorgan        4.33      2.76      1.59  0.64  0.89
## Citibank        2.76      4.39      1.80  1.81  1.23
## WellsFargo      1.59      1.80      2.24  0.73  0.61
## Shell           0.64      1.81      0.73  7.22  5.08
## Exxon           0.89      1.23      0.61  5.08  7.66
```

20 / 28

Factor model estimation by Maximum Likelihood method

Create normalized data with variance = 1

```
normdata=(as.matrix(stock))%*%diag(1/sqrt(diag(cov(stock))))
dim(normdata)
```

```
## [1] 103 5
```

```
cov(normdata)
```

```
##      [,1] [,2] [,3] [,4] [,5]
## [1,] 1.0000 0.6323 0.5105 0.1146 0.1545
## [2,] 0.6323 1.0000 0.5741 0.3223 0.2127
## [3,] 0.5105 0.5741 1.0000 0.1825 0.1462
## [4,] 0.1146 0.3223 0.1825 1.0000 0.6834
## [5,] 0.1545 0.2127 0.1462 0.6834 1.0000
```

21 / 28

```
ML2 =factanal(normdata,2, rotation="none")
ML2$call
```

```
## factanal(x = normdata, factors = 2, rotation = "none")
```

```
ML2$method
```

```
## [1] "mle"
```

```
ML2$uniquenesses
```

```
## [1] 0.4165 0.2747 0.5420 0.0050 0.5298
```

```
ML2$loadings
```

22 / 28

Proportion of variance explained by 2 common factors (ML)

$$\frac{\hat{\ell}_{1j}^2 + \dots + \hat{\ell}_{pj}^2}{s_{11} + \dots + s_{pp}} = \frac{\hat{\ell}_{1j}^2 + \dots + \hat{\ell}_{pj}^2}{p}, \quad j = 1, \dots, m.$$

```
##
```

```
## Loadings:
```

```
##      Factor1 Factor2
```

```
## [1,] 0.121 0.754
```

```
## [2,] 0.328 0.786
```

```
## [3,] 0.188 0.650
```

```
## [4,] 0.997
```

```
## [5,] 0.685
```

```
##
```

```
##      Factor1 Factor2
```

```
## SS loadings      1.622 1.610
```

```
## Proportion Var   0.324 0.322
```


```
## Cumulative Var   0.324 0.646
```

23 / 28

Plots of variables on two factors

```
Lm = factanal(normdata,2,rotation="none")$loading[,1:2]
plot(Lm,type="n",main="Factor loadings (m=2) by ML")
text(Lm,labels=(row.names(L)),col=1:5)
```

Factor loadings (m=2) by ML



24 / 28

Remarks on ML method for FA

- ML method for FA always estimate correlation matrix

```
factanal(stock,2,rotation="none") #same as using normdata
```

- Large sample test for the number of common factors*

```
ML2$STATISTIC # Bartlett chisq approximation for LR test
```

```
## objective
##      1.974
```

```
ML2$dof # [(p-m)^2 - p - m]/2
```

```
## [1] 1
```

```
ML2$PVAL # p-value
```

```
## objective
##      0.16
```

25 / 28

Factor estimations by ML vs PC

Check residuals $\Sigma - LL' - \Psi$

ML (m=2)

```
Lm = factanal(normdata,2, rotation="none")$loading[,1:2]
```

```
Psim = factanal(normdata,2, rotation="none")$uniq
```

Residual matrix for m = 2, using ML method

```
round(cor(stock) - Lm%*%t(Lm) - diag(Psim),3)
```

```
##                JPMorgan Citibank WellsFargo Shell  Exxon
## JPMorgan      0.000    0.000    -0.003    0  0.052
## Citibank      0.000    0.000     0.002    0 -0.033
## WellsFargo   -0.003    0.002     0.000    0  0.001
## Shell        0.000    0.000     0.000    0  0.000
## Exxon        0.052   -0.033     0.001    0  0.000
```

26 / 28

Check model performance

Residual matrix = $\Sigma - LL' - \Psi$

PC (m=2) on correlation matrix

Residua matrix for m = 2, using PC method

```
normrtev = princomp(stock,cor=T)$sdev
```

```
Ln = cbind(normrtev[1]*princomp(stock,cor=T)$loading[,1],
           normrtev[2]*princomp(stock,cor=T)$loading[,2])
```

```
round(cor(stock) - Ln%*%t(Ln) - diag(rep(1,5)
                                     - diag(Ln%*%t(Ln))),3)
```

```
##                JPMorgan Citibank WellsFargo  Shell  Exxon
## JPMorgan      0.000   -0.099   -0.185 -0.025  0.056
## Citibank     -0.099    0.000   -0.134  0.014 -0.054
## WellsFargo   -0.185   -0.134    0.000  0.003  0.006
## Shell       -0.025    0.014    0.003  0.000 -0.156
## Exxon        0.056   -0.054    0.006 -0.156  0.000
```

27 / 28

Which method is better for this data set?

- Compare proportions of variation explained
- Compare residual matrix element size

ML methods sum-abs and sum-sq of residual(ij), m=2

```
c(sum(abs(cor(stock) - Lm%*%t(Lm) - diag(Psim))),
  sum((cor(stock) - Lm%*%t(Lm) - diag(Psim))^2))
```

```
## [1] 0.180946 0.007612
```

#PC method sum-abs, sum-sq of residuals, m=2, corr mat

```
c(sum(abs(cor(stock) - Ln%*%t(Ln) - diag(rep(1,5)
                                             - diag(Ln%*%t(Ln)))))
  , sum((cor(stock) - Ln%*%t(Ln) - diag(rep(1,5)
                                             - diag(Ln%*%t(Ln))))^2))
```

```
## [1] 1.4630 0.1861
```

28 / 28