

Differential Geometry

Kristian Wichmann

February 11, 2018

1 Regular curves in Euclidean spaces

Let V be a finite-dimensional vector space over \mathbb{R} . Then a C^k *regular curve* in V is a mapping:

$$x : I \rightarrow V \quad (1.1)$$

Here I is an interval, and we require that x is C^k and that $x'(t) \neq 0$ for any $t \in I$ (this last requirement is what makes the curve regular).

Given a C^k regular curve $x : I \rightarrow V$. Then if φ is a C^k bijection from the interval J to I having $\varphi' > 0$, we consider x equivalent to the regular curve $y : J \rightarrow V, y = x \circ \varphi$.

We now assume V has an inner product (\cdot, \cdot) and an induced norm $\|v\| = \sqrt{(v, v)}$. Then for a regular curve x we define an arc length function $s(t)$ by:

$$\frac{ds}{dt} = \|x'(t)\| \quad (1.2)$$

Or equivalently:

$$s(t) = \int \|x'(t)\| dt \quad (1.3)$$

This means that two arc length functions differ only by a constant. Since x' is never zero, $\|x'(t)\| > 0$, and so $s(t)$ is strictly increasing. As an integral of a C^{k-1} function it is itself C^k .

We now define the function $X(s(t)) = x(t)$. Being a composition of two C^k functions, so is X . So we can always parametrize a regular curve by its arc length. The derivative of X is found by the chain rule:

$$\frac{dX}{ds} = \frac{dx}{dt} \frac{dt}{ds} = x'(t(s)) \frac{1}{ds/dt} = \frac{x'(t(s))}{\|x'(t(s))\|} \quad (1.4)$$

Here we've used, that since there is a bijection between I and $s(t)$ we can regard t as a function of s . Taking the norm, we get $\|X'(s)\| = 1$. Squaring

this gives $\|X'(s)\|^2 = (X'(s), X'(s)) = 1$. Differentiate with respect to s to get:

$$2(X''(s), X'(s)) = 0 \quad (1.5)$$

So the second derivate is orthogonal to the tangent $X'(s)$.

1.1 Example: Circular curve

Let $x_0 \in V$ and let $r_1, r_2 \in V$ be perpendicular vectors, each of length R . Then we can make a circular curve centered in x_0 , radius R , and in the plane spanned by r_1 and r_2 by:

$$x(s) = x_0 + r_1 \cos(s/R) + r_2 \sin(s/R) \quad (1.6)$$

The tangent, i.e. the first derivative is:

$$x'(s) = \frac{1}{R}[-r_1 \sin(s/R) + r_2 \cos(s/R)] \quad (1.7)$$

The second derivative is:

$$x''(s) = -\frac{1}{R^2}[r_1 \sin(s/R) + r_2 \cos(s/R)] = -\frac{1}{R^2}(x(s) - x_0) \quad (1.8)$$

This is pointed at the center of the circle and has a magnitude of $\|x''(s)\| = 1/R$.

1.2 Curvature of regular Euclidean curves

Inspired by the circular curve example, we make the following definition:

Definition 1.1. *Given a regular C^2 curve in V parametrized by arc length: $x(s)$ and a point on the curve where $x''(s) \neq 0$.*

Then $n = x''(s)/\|x''(s)\|$ is called the principal normal to the curve at $x(s)$.

$1/\|x''(s)\|$ is known as the radius of curvature at $x(s)$. The circle with center at $x(s) + x''(s)/\|x''(s)\|$, radius equal to the radius of curvature, in the plane spanned by $x'(s)$ and n is known as the osculating circle.

$\kappa = \|x''(s)\|$ is called the curvature of the curve at $x(s)$, even if equal to zero. So when $\kappa \neq 0$ we have $x''(s) = n\kappa$

What if our curve $x(t)$ is not parametrized by arc length s ? Then we can still calculate curvature by using the X function: $x(t) = X(s(t))$. The first derivative of x can be found using the chain rule:

$$x'(t) = \frac{dX}{dt} = \frac{dX}{ds} \frac{ds}{dt} = X'(s) \frac{ds}{dt} \quad (1.9)$$

And the second derivative by the multiplication rule:

$$x''(t) = X''(s) \left(\frac{ds}{dt} \right)^2 + X'(s) \frac{d^2s}{dt^2} \quad (1.10)$$

Remember that $ds/dt = \|x'(t)\|$:

$$x''(t) = X''(s) \|x'(t)\|^2 + X'(s) \frac{d}{dt} \|x'(t)\| \quad (1.11)$$

Consider the last derivative:

$$\frac{d}{dt} \|x'(t)\| = \frac{d}{dt} \sqrt{(x'(t), x'(t))} = \frac{1}{2\sqrt{(x'(t), x'(t))}} 2(x'(t), x'') = \frac{(x'(t), x''(t))}{\|x'(t)\|} \quad (1.12)$$

Rearranging terms and using equation 1.4 we get:

$$X''(s) \|x'(t)\|^2 = x''(t) - \frac{(x'(t), x''(t))}{\|x'(t)\|^2} x'(t) \quad (1.13)$$

Finally we can find the second derivative of X :

$$X''(s) = \frac{x''(t)}{\|x'(t)\|^2} - \frac{(x'(t), x''(t))}{\|x'(t)\|^4} x'(t) \quad (1.14)$$

$$= \frac{1}{\|x'(t)\|^2} \left(x''(t) - \frac{(x'(t), x''(t))}{\|x'(t)\|^2} x'(t) \right) \quad (1.15)$$

This means that the curvature is:

$$\kappa = \|X''(s)\| = \frac{1}{\|x'(t)\|^2} \left\| x''(t) - \frac{(x'(t), x''(t))}{\|x'(t)\|^2} x'(t) \right\| \quad (1.16)$$

1.3 Regular curves in \mathbb{R}^2

Let $x(t)$ be a regular curve in \mathbb{R}^2 . Then we may attach a sign to the curvature κ . Let $x(s)$ be a parametrization of the curve by arc length. Then if $\|x''(s)\| \neq 0$, we know that $x'(s)$ and $x''(s)$ are perpendicular. Because we're in two dimensions, $x''(s)$ is either rotated 90 degrees in positive or negative direction (counter-clockwise or clockwise). This determines the sign of the curvature.

Now consider a regular curve $x(s)$ in \mathbb{R}^2 that is also closed and simple, i.e. one that is periodic, and has no self-intersections apart from this. If the arc length is L we will still let the parametrization be defined for all $s \in \mathbb{R}$, so that:

$$x(s) = x(s') \Leftrightarrow s' - s = nL, n \in \mathbb{Z} \quad (1.17)$$

Since $\|x'(s)\| = 1$ we can parametrize:

$$x'(s) = (\cos(\theta(s)), \sin(\theta(s))), 0 \leq s \leq L \quad (1.18)$$