

HEMODYNAMIC VORTEX ANALYSIS AS A MEANS OF INTRACRANIAL
ANEURYSM RUPTURE PREDICTION

By

Kevin William Sunderland

A DISSERTATION

Submitted in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

In Biomedical Engineering

MICHIGAN TECHNOLOGICAL UNIVERSITY

2019

© 2019 Kevin William Sunderland

This dissertation has been approved in partial fulfillment of the requirements for the Degree of DOCTOR OF PHILOSOPHY in Biomedical Engineering.

Department of Biomedical Engineering

Dissertation Advisor: *Dr. Jingfeng Jiang*

Committee Member: *Dr. Sean Kirkpatrick #1*

Committee Member: *Dr. Gowtham #2*

Committee Member: *Dr. Min Wang #3*

Department Chair: *Dr. Sean Kirkpatrick*

Dedication

To my famliy and friends

who

Contents

List of Figures	xi
List of Tables	xv
Preface	xvii
Acknowledgments	xxi
Definitions	xxiii
List of Abbreviations	xxvii
Abstract	xxxi
1 Introduction	1
1.0.1 Current Rupture Prediction Metrics	2
1.1 Section 1	10
1.1.1 Objective	11
1.1.2 Methodolgy	11
1.2 Aneurysm Geometric Characterisitics	13

1.3	Aneurysm Hemodynamic Characterisitcs	15
1.4	Disturbed Flow on Vascular Endothelium	16
2	Cerebral Vasculature and the Impact of Disturbed Flow	21
2.0.1	Endothelial Cells	22
2.0.2	Impact of Disturbed Flow on Vascular Cells	27
3	Hemodynamic Flow Vortex Identification	33
3.1	Materials and Methods	36
3.1.1	Modeling of "Patient-specific" Vasculature	36
3.1.2	Mesh Generation	37
3.1.3	CFD Simulation	37
3.1.4	Aneurysm Extraction and Voxelization of Aneurismal Velocity Data	39
3.1.5	Vortex Core Extraction and Analysis	40
4	Vortex Analysis to predict IA Initiation	51
	References	57
A	Statistics	109
A.1	Section 1	110
A.2	Section 2	114
A.3	Section 3	117

A.4	Section 4	118
A.5	Section 5	118
A.6	Section 5	120
B	Sample Code	123
B.1	HelloWorld.c	125
C	Letters of Permission	127
D	Cellular Biology	129
D.1	TUNEL-assay	129
D.2	VCAM-1	130

List of Figures

1.1	Zhou 2016 Meta-analysis of the reported low WSS rate of rupture aneurysms and the Odds Ratio for low WSS in predictive modeling	5
1.2	The wall shear stress (N/mm^2) distribution on the aneurysm wall(A) and blood flow pattern inside the aneurysm visualized with streamlines (mm/sec)(B). Wall shear stressors may overlook the disturbed/swirling flow patterns (white arrow) that are generated in the IA sac.	8
1.3	Areas of likely cerebral IA development: Original image from [209].	9
1.4	Schematic representation of our universe	12
1.5	Mathematical functions plotted using TikZ package	12
1.6	Schematic representation of a water molecule	18
2.1	Layers of the vasculature. Original image from [218]	23

2.2 Mechanoreceptors of ECs: ion channels (K^+ , Ca^{2+} , Na^+ , Cl^-), G-proteins, caveolae, tyrosine kinase receptors (TKRs), nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, xanthine oxidase (XO), integrins, and heparan sulfate proteoglycan. Signals are transmitted through the cytoskeleton to the basal or junctional endothelial surface. Integrin mechanosensory complexes consist of platelet endothelial cell adhesion molecule-1 (PECAM-1) and Flk-1. When activated they initiate downstream signaling cascades. Activated integrins trigger multiple complex of non-receptor tyrosine kinases (FAK, c-Src, Shc, paxillin, and p130CAS), adaptor proteins (Grb2, Crk), and guanine nucleotide exchange factors (Sos, C3G), thereby activating Ras family GTPase. Active Ras plays a pivotal role in intracellular transduction of signals as it triggers various parallel downstream cascades of serine kinases; each of these activate downstream signals, ultimately activating mitogen-activated protein kinases (MAPKs). Shear stressors also activate a number of other downstream signaling pathways that impact reactive oxygen species (ROS): NADPH oxidase, activation of protein kinase C (PKC), activation of Rho family small GTPases (mediate the EC remodeling), release of endothelial nitric oxide synthase (eNOS) and activation of phosphoinositide-3 kinase (PI3K)-Akt cascade. These signaling pathways lead to phosphorylation of transcription factors (TFs) such as nuclear factor-kappa($NF-\alpha\beta$) and activator protein-1(AP-1).

2.3	Schematic representation of abdominal aortic aneurysm (AAA) pathophysiology. Inflammation, proteolysis (breakdown of proteins by enzymes), smooth muscle cell (SMC) apoptosis, neovascularization, calcification, and intraluminal thrombosis may be targeted by molecular imaging. ECM indicates extracellular matrix; and MMP, matrix metalloproteinase.	28
2.4	Fancy mathematical plots using TikZ package	31
2.5	Incidence, transmission and reflection	32
3.1	Distribution of random numbers	45
3.2	Fibre optics	47
3.3	A landscape view of a Turboprop engine - these are jet engine derivatives, still gas turbines, that extract work from the hot-exhaust jet to turn a rotating shaft, which is then used to produce thrust by some other means	49
4.1	Layers of the vasculature. Original image from [218]	55
B.1	Two examples illustrating the relationship between the angular histogram and NE: (a) a simple laminar flow case and (b) a rotational flow (eddy) case. In both cases, the right and left plots are the vector flow field and the histogram of angular vector direction, respectively. Vector fields were decimated by a factor of 3 for better visualization.	124

List of Tables

1.1	Recent articles for aneurysm rupture prediction	7
2.1	Disturbed Flow's Impact on Vascular Cells	29
2.2	A portrait table: first column represents the year in which the Nobel prize in physics was awarded; second column indicates the name of the scientist and the third column is the work for which the Nobel prize was awarded	30
3.1	Measured data points representing the relationship between x and y	45
3.2	A landscape table: first column represents the year in which the Nobel prize in physics was awarded; second column indicates the name of the scientist and the third column is an <i>as is</i> Nobel citation	46

Preface

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur.

Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Acknowledgments

I would like to thank all the members of my committee and my advisor Dr. Jingfeng Jiang. Their leadership, support, knowledge and motivation not only helped me throughout my research, but helped drive me to become a better scientist and to never stop learning.

Special thanks are also needed for Dr. Autumn Schumacher, who was willing to take a gamble on a brand new scientist fresh out of their undergraduate education. Her and expertise (and many hours of manuscript editing) were invaluable in getting me to where I am today.

I would also like to thank my friends for their boundless confidence in me which helped push me through my PhD work. Last but not the least, I would of course like to thank my family. All of their love and support helped make this thesis possible.

Definitions

This provides information on how to write your MS thesis or PhD dissertation using the \LaTeX document preparation system in compliance with Michigan Technological University Graduate School requirements.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure

dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

List of Abbreviations

ACA	Anterior Communicating Artery
AFI	Aneurysm Formation Indicator
CFD	Computational Fluid Dynamics
DICOM	Digital Imaging and Communications in Medicine
DVO	Degree of Volume Overlap
ENR	Elastic Net Regression
IA	Intracranial Aneurysm
ICA	Internal Carotid Artery
MCA	Middle Cerebral Artery
MLR	Multiple Logistic Regression
NSC	Nearest Shrunk Centroid
OSI	Oscillatory Shear Index
PC-MRI	Phase Contrast Magnetic Resonance Imaging
ROC	Receiver Operator Characteristic
STA-WSS	Spatiotemporally Averaged Wall Shear Stress
TA-WSS	Temporally Averaged Wall Shear Stress
VMTK	Vascular Modeling Toolkit
VTK	Visualization Toolkit

WSS	Wall Shear Stress
WSSG	Wall Shear Stress Gradient
λ_2	Lambda ₂
ACL	Access Control List
AIB	Add-In Board
ALE	Arbitrary Lagrangian Eulerian
AMANDA	Advanced Maryland Automatic Network Disk Archiver
AMBER	Assisted Model Building with Energy Replacement
AMD	Advanced Micro Devices
AMOLED	Active-Matrix Organic Light Emitting Diode
AMPI	Adaptive Message Passing Interface
ANL	Argonne National Laboratory
API	Application Program Interface
ASCII	American Standard Code for Information Interchange
ATLAS	Automatically Tuned Linear Algebra Software
b_eff	effective bandwidth Benchmark
BIOS	Basic Input/Output Operating System
BLAS	Basic Linear Algebra Subprograms
BOMD	Born-Oppenheimer Molecular Dynamics
BP	Bootstrap Protocol
CCSR	Center for Computer Systems Research

CentOS	Community enterprise Operating System
CFD	Computational Fluid Dynamics
CHARMM	Chemistry at HARvard Macromolecular Mechanics
CHAMBER	CHarmm \leftrightarrow AMBER
CMake	Cross Platform Make
CODINE	Computing in Distributed Networked Environments
CP2K	Car-Parrinello 2000
CPMD	Car-Parrinello Molecular Dynamics
CPU	Central Processing Unit
CSS	Central Security Service
CTM	Chemical Transport Model
CUDA	Compute Unified Device Architecture
CUDPP	CUDA Data-Parallel Primitives Library
DAE	Differential Algebraic Equation
DARPA	Defense Advanced Research Projects Agency
DAE	Delay Differential Equation
DFT	Discrete Fourier Transform
DFT	Density Functional Theory
DGEMM	Double Precision GEneralized Matrix Multiplication
DHCP	Dynamic Host Configuration Protocol
DMCA	Digital Millennial Copyright Act

DOD	Department of Defense
DOE	Department of Energy
DRM	Distributed Resource Manager
DRMAA	Distributed Resource Manager Application API
EFF	Electron Force Field
EVL	Electronic Visualization Laboratory
FCA	Fabric Collectives Accelerator
FEA	Finite Element Analysis
FFT	Fast Fourier Transform
FFTW	Fastest Fourier Transform in the West
FLOPS	Floating Point Operations per Second
FPU	Floating Point Unit
FSI	Fluid Structure Interaction
FTDT	Finite Difference Time Domain
FTP	File Transfer Protocol

Abstract

This provides information on how to write your MS thesis or PhD dissertation using the \LaTeX document preparation system in compliance with Michigan Technological University Graduate School requirements.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure

dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Sed ut perspiciatis unde omnis iste natus error sit voluptatem accusantium doloremque laudantium, totam rem aperiam, eaque ipsa quae ab illo inventore veritatis et quasi architecto beatae vitae dicta sunt explicabo. Nemo enim ipsam voluptatem quia voluptas sit aspernatur aut odit aut fugit, sed quia consequuntur magni dolores eos qui ratione voluptatem sequi nesciunt.

Chapter 1

Introduction

Subarachnoid hemorrhage is a potentially devastating pathologic condition in which bleeding occurs into the space surrounding the brain. One of the prevalent events that result in subarachnoid hemorrhage is the rupture of an intracranial aneurysm (IA). IAs are degenerative, irregular expansions of areas in the cerebral vasculature that occur in an estimated 3-5% of the global population [100, 198, 247], with an estimated 0.15% - 0.7% of the global population suffering the rupture of an IA each year [114]. Mortality rates due to the rupture of IAs are estimated between 45-50%, with survivors suffering significant neurological damage, including physical and cognitive impairment [156, 247]. Yet the disparity between the number of ruptured and unruptured IAs indicate that not all IAs are at high rupture risk. From a clinical perspective, improved medical imaging techniques have led to an increase in the detection

of unruptured IAs, and novel surgical intervention methods have aimed to reduce the instances of IA rupture and their subsequent impacts of patient health [140, 175]. Yet surgical treatments are not without a high healthcare costs: surgical repair (without complications) of IAs can be an estimated \$25,000. Data has also shown that, while treatment risks are relatively low, similar outcomes of morbidity and mortality can occur in the event of treatment complications [45, 153, 164]. Surgical complications also result in a marked increase in costs: between an estimated \$35,000 - \$70,000 depending on possible complications (varies with surgical intervention) [28]. Additionally, continued economic burden is placed upon patients and their families if long term rehabilitation, home care, or in-hospital treatment is required post-surgical complications. This surgical vs rupture risk has led to significant interest in elucidating the processes and conditions which impact IA rupture. Proper identification of IAs at high risk of rupture could significantly benefit risk stratification, improve patient selection and aid treatment planning as to best determine a course of action for the clinical management of IAs.

Current Rupture Prediction Metrics

Research has shown that a wide array of risk factors may impact IA development, growth and rupture potential [67, 141, 188, 220], with said factors generally separated into three categories:

- IA morphological characteristics
- Vascular hemodynamics
- Patient genetic and health factors

Morphological characteristics of IAs are often first considered when determining the severity of an unruptured IA and its potential to rupture. Factors such as IA size (volume), shape, aspect ratio, IA to parent vessel angles, etc. **NEEEEEED REFERENCES** have been identified in a number of studies as a possible metrics to assess IA rupture potential. However, many of these parameters have been shown to generate conflicting results between studies, varying in their strengths toward rupture prediction. As per example, IAs $>10\text{mm}$ in size are often associated with rupture risk, with IAs $>25\text{mm}$ thought to be at the greatest rupture risk leading the majority of IAs at these sizes to undergo surgical intervention[173, 258, 264] even though not all large IAs rupture. If this thought process would be applied clinically, many small IAs ($<4\text{mm}$) may be spared surgical intervention due to the thought that the complications from surgical interventions would be of greater risk than the possibility of small IA rupture. Yet small aneurysms have been shown to be at a non-insignificant risk of rupture [78]. In addition to assessing rupture risk solely on IA size may overestimate the risk of large IAs (and subsequently underestimating small IAs), size characteristics may be of limited usefulness in assessing rupture risk in medium-sized IAs (4-10mm). Additionally, many of the other morphological characteristics of IAs

have had varying degrees of strength toward predicting IA rupture[10, 29, 124]. To overcome the limitations of relying on morphological characteristics for assessment of IA rupture potential, hemodynamic characteristics within the IA sac have been investigated.

Hemodynamic stressors are thought to play a significant role on the initiation, growth and possible rupture of IAs. Tangential fluid stressors along the vascular wall, known as wall shear stress (WSS), and its derivatives are often of investigated when generating models for IA rupture prediction(s)[38, 68, 124, 264]. These near-wall forces have been shown to act as a biological stimulator, eliciting various changes to both vasculature endothelial and smooth muscle cells, inducing changes in gene expression, phenotypical and mechanical properties, and protein expressions which dictate cellular activity [13, 15, 32, 41, 69, 148, 202, 241]. While WSS has been assessed to trigger vascular cell changes, discrepancies exist to which specific hemodynamic characteristics impact IA rupture prediction and their overall predictive strength, similar to IA morphological characteristics. It has been shown that vascular cells maintain healthy physiological characteristics while exposed to a range of WSS between 5 and 20 dynes/cm² yet studies are in conflict if WSS lower [27, 172] or higher [69, 213] from said preferred range is of greater impact on IA rupture risk. It has been suggested that both high and low WSS trigger differing impacts on vascular cells [169], meaning both may play a significant role in IA development and potential rupture. Yet even focusing on one extrema of WSS values and determining their usefulness on prediction

IA rupture risk has proven difficult. In a 2016 meta analysis by Zhou, the impact of low wall shear stress on predicting IA rupture varied between studies (Fig. ??). In addition to WSS extrema as a predictive metric toward IA rupture, fluctuations in WSS gradients and changes in WSS directionality (Oscillatory shear index (OSI)) are thought to cause pathologic changes to the vasculature are often applied to IA rupture prediction models. Its has been theorized that fluctuations in WSS directionality can trigger varying changes to vascular cells, weakening the overall strength of the vasculature and triggering possible IA development or rupture.




Figure 1.1: Zhou 2016 Meta-analysis of the reported low WSS rate of rupture aneurysms and the Odds Ratio for low WSS in predictive modeling

It is worth noting, that the clinical measurement of WSS characteristics *in-vivo* is possible, yet remains difficult especially in smaller blood vessels and small IAs. Improvements in computational fluid dynamics (CFD) can help overcome the limitations

of *in-vivo* WSS measurement and is being adopted as a clinical tool for assessment of rupture potential and planning of IA treatments [222, 240].

While the development of IAs have been related to hemodynamic factors, a number of genetic and patient health-factors have also been associated with structural weakness in the arterial wall indicative of IAs[174, 229]. The increased likelihood of IA development in a person whose family member(s) have also presented with an IA gave early insight towards a possible genetic component to IAs.

A number of patient health factors have also been associated with IA development/rupture and have been incorporated into prediction models. Patient age, gender and smoking habits, which have been linked with changes to the mechanical properties of the vasculature, have been incorporated into IA rupture prediction [10, 29, 166]s. Increases in age and smoking[11, 97, 141, 167] have been linked with a breakdown of the collagen of the vasculature, weakening the overall mechanical strength of the vasculature. There is also a breakdown in the elastin complexes of the vasculature as a function of aging, decreasing the compliance of the vessel (to hemodynamic conditions) and increasing vessel stiffness[79, 252]. Additionally, hypertension has also been linked with an increase in the likelihood of aneurysm development and potential rupture[193, 226]. Hypertension is thought to weaken vessels by increasing mechanical stresses while also causing a subsequent increase in vascular inflammation, leading to possible remodeling of the vascular wall[207].

Combining the assessment of IA morphological characteristics, WSS and genetic/patient medical information have been applied to a number of models aiming to predict the likelihood of IA rupture. Yet the indices chosen and the resultant strength of predictions models can vary significantly between studies (1.1).

Table 1.1
Recent articles for aneurysm rupture prediction

Author-Year	Parameters for Analysis	Aneurysm Location(s)	Ruptured IAs	Unruptured IAs	AUC or Accuracies
Qin et al., 2017[194]	W, H, L, NW, Age, AR, Dmax, HW, BF, WSS, LSA, EL	MCA	36	31	0.931
Bijlenga et al., 2017[25]	S, Age, Hyp, Race, SAH	MCA, ICA, ACA/PcomA/-Post	598	243	0.681-0.756
Detmer et al., 2018[65]	Age, Gender, S, OSIm, NSI	ACA, AcoM, BA, ICA, MCA, PCom, VA	66	183	0.82
Kocur et al., 2019[136]	S, H, W, NS, AR, BF, H/W, SR, A	AcoM, ICA, MCA, Post	146	285	0.55-0.64
Wang et al., 2019[250]	R, DD, NW, W, D, Dia, AR, DW, BF, SR, F, LD, SD,	AcoM	214	147	0.846
Varble et al., 2018[245]	Age, SR, AR, UI, EL, NSI, WSS, OSI, WSSG, RRT, LSA, MWSS, PLc, EL	ACA, ICA, MCA, PcoM, Post	102 (Train) 14 (Test)	311 (Train) 115 (Test)	0.767
Jiang et al., 2018[123]	L, H, AR, SR, UI, EI, D, NSI, NWSSa, WSSm, WSSa, NWSSa, NWSSm, WSSG, LSA, OSI, RRT, Pm, Pa, NPa, NPm	ACA, AcoM, ICA, MCA, PcoM	167	167	0.81
Aneriteor communicating artery (AcoM), Anterior communicating artery (ACA), Internal carotid artery (ICA), Middle cerebral artery (MCA), Posterior communicating artery (PcoM), Posterior inferior cerebral artery (Post), Aneurysm angle (A), Area (A), Aspect Ratio (AR), Bottleneck factor (BF), depth/width (DW), Diameter (D), Energy Loss (EL), Flow angle (F), Height (H), Hypertension (Hyp), Irregular shape (IR), Large daughter artery angle (LD), Lateral angle ratio (LAR), Length (L), Low wall shear area (LSA), Mean diameter (MD), Mean wall shear stress (MWSS), Non-sphericity index (NSI), Normalized pressure average (NPa), Normalized pressure max (NPm), Normalized wall shear stress area (NWSSa), Normalized wall shear stress mean (NWSSm), Oscillatory shear index(max) (OSIm), Pressure average (Pa), Pressure loss(coefficient) (PLc), Pressure max (Pm), Relative residence time (RRT), Size ratio (SR), Small daughter artery angle (SD), Surface (S), Temporally averaged WSS (TAWSS), Undulation Index (UI), Vessel angle (V), Width (W), Wall shear stress (WSS), Wall shear stress gradient (WSSG), Wormsley number (WN), Wall shear stress mean (WSSm).					

These variations between study outcomes suggest that novel assessment methods may help improve the understanding of IA characteristics indicative of rupture. Of recent

attention are the assessments of broader flow patterns occurring within the aneurysm sac and parent vessel for their relationship to aneurysms. While WSS characteristics assess the 'near-wall' flow within the vasculature, this overlooks the remaining flow characteristics within the vasculature that may help improve prediction models (Fig. 1.2).




Figure 1.2: The wall shear stress (N/mm^2) distribution on the aneurysm wall(A) and blood flow pattern inside the aneurysm visualized with streamlines (mm/sec)(B). Wall shear stressors may overlook the disturbed/swirling flow patterns (white arrow) that are generated in the IA sac.

Early investigations into the nature of IAs uncovered the non-random distribution of areas of the vasculature susceptible to IA development: vessel bifurcations and areas of significant vasculature curvature (Fig ??). These areas of likely aneurysm development were shown to correlate with areas of disturbed, swirling hemodynamic flow patterns (vortex/vortices). In much of the vasculature, blood maintains laminar flow characteristics in which fluid travels smoothly in regular paths with little to no mixing. Vortices are antithetical to laminar flow, with areas of significant mixing/recirculation, and causing a significant drop in WSS values. Not only are said vortex patterns present in specific areas of the vasculature, they are also present

within IA sacs. These findings led research toward understanding the impact that these vortices may have on altering the vasculature, triggering IA development and rupture[39, 53, 155]. As hemodynamic forces help regulate the physiological characteristics of the vasculature, vortex flow conditions are thought to impart pathologic cellular changes to vascular cells.




Figure 1.3: Areas of likely cerebral IA development: Original image from [209].

Yet the analysis of disturbed flow is typically done from a visual qualitative assessment of computational fluid dynamic (CFD) based flow patterns or through limited computational analysis of a vortex’s centroid region. These methodologies can only result in a restrictive assessment of the characteristics of vortices, and could limit their usefulness in predicting IA rupture. As per example, a study by Varble et al[244]. was unable to find a strong correlation between vortices and IA rupture, but said study

only focused on identifying the centroid region of vortices. A more comprehensive analysis of vortices may give novel insight towards IAs and could provide a useful tool to help improve the assessment of IA rupture potential.

The **focus of this thesis** is to apply novel computational technique to assess the spatiotemporal characteristics of broader structure of hemodynamic vortex patterns and determine their usefulness in identifying cerebral IAs rupture risk.

While a number of metrics (geometric, hemodynamic, and health factors) have aided in understanding possible mechanisms triggering IA development and rupture, Investigations of hemodynamic vortices in both *in-vivo* and *in-vivo* and have been shown to elicit endothelial cell dysfunction [17, 160, 180].

Section 1

At vix indoctum disputando. Eam cu doctus reprimique, quaeque democritum an eos, sit veniam facete dissentias id. Tale volumus eos te, an eum nulla tincidunt. Mea id recteque theophrastus.

Eirmod malorum vis ei. Choro euismod incorrupte in vim, ludus ornatus vis ex. Hinc wisi impedit eum no, vocent definiebas referrentur in quo. Sanctus vulputate repudiandae usu ut.

Objective

Although there exists a number of studies[34, 245, 277] and methodologies[83, 97] that attempt to assess IAs at a high risk of rupture, inconsistencies between study outcomes leave the development of an ideal predictive model out of reach. In addition, many of these previous studies assess the geometric[2, 130, 245] and/or hemodynamic wall stressors[34, 172, 277] as a means to predict IA rupture, with limited quantitative assessment of the hemodynamic flow conditions within the aneurysm. **The primary objective** of this work is to assess the viability of adapting quantitative analysis of hemodynamic flow patterns, specifically swirling flow pattern(s) (vortex), within IAs to improve the prediction and understanding of IA rupture. In this work, an overview of recent theories concerning

Methodolgy

For the initial focus of this work, image-based computational fluid dynamics models of patient-specific IA geometry will be constructed from 3D phase contrast magnetic resonance imaging (PC-MRI). Computational fluid dynamic (CFD) simulations will be performed on the computational models to generate realistic 3D hemodyanmic velocity and flow pattern data. From said data,




Figure 1.4: Schematic representation of our universe




Figure 1.5: Mathematical functions plotted using TikZ package

Simul noster voluptaria eam ei, sint regione pri ei. Cum no utinam equidem, falli bonorum prodesset an qui. Alterum dissentiet vituperatoribus te eam, eos ea suas oblique. Per ea utinam facilisi. [61, 71, 72] Per iudico probatus complectitur et, cum tollit atomorum rationibus ea.

Aneurysm Geometric Characteristics

All aneurysm geometries were taken from the finalized computational mesh generated for simulations. The aneurysm sac was manually isolated from the parent vessel and the resultant cut plane was capped and identified as the IA ostium using an in-house script written in VMTK. Geometric measurements were either taken directly from the values reported in the Aneurisk dataset, or were calculated using in-house scripts in VMTK.

Aneurysm Surface Area and Volume: Measured directly from the isolated IA geometry before and after (respectively) ostium capping. A number of studies have eluded to an increase in IA size as a risk for both IA growth and rupture. [11, 29, 97, 245]. A meta-analysis performed by Brinjikji et al reported that IA ≤ 10 mm in size (diameter) grew at a rate $< 2.9\%$ per year, while IAs > 10 mm were associated with growth rates of 9.7% per year. This growth was also reported with an associated IA rupture rate: 3.1% per year compared with 0.1% per year for stable (non-growing) aneurysms ($p \leq 0.01$). From a clinical perspective, the overall size of an aneurysm is often a characteristic used to determine course of IA treatment (or lack thereof) [140, 259]. Yet while large IAs are thought to increase the likelihood of rupture, a not-insignificant number of small IAs (< 5 mm diameter) also have been shown to rupture [126, 130, 141]. This disparity between sizes of ruptured IAs suggest that

the assessment of additional factors in tandem with IA size may improve rupture prediction.

Aneurysm Height: The length of the centerline of the IA sac is measured, following the IA shape, as opposed to measuring a straight line from the ostium centroid directly to the highest IA point. The radius of the maximum inscribed sphere at the centerline's furthest point is added to the length measurement to fully measure the IA height. This is a modified version of the typical IA height measurement: a straight line of the maximum stretch from the ostium centroid to the IA dome [78, 158].

Vessel Diameter: The parent artery diameter value is computed at locations close to the aneurysm ostium. For terminal aneurysms, the vessel diameter of the common branch was measured at the point prior to centerline splitting between the daughter arteries, and both daughter arteries' diameter were measured at the point one (common artery) diameter away from the IA ostium cut. The average of the three values was used as the value of the vessel diameter.

Inlet Cross-sectional Area: The beginning of the inlet vessel was cut square in the 3-matic software package, the resultant cross-sectional area of the inlet vessel was calculated.

Aspect Ratio*: A modified calculation of the commonly defined aspect ratio

(aneurysm height/ostium diameter) was used by adapting the length of the center-line of an IA as the IA height ($IA_{height*}$), and the area and circumference of the ostium since ostium diameter is rarely uniform for an IA [189].

$$AspectRatio^* = \frac{IA_{height*}}{4 * (Ostium_{area}/Ostium_{circumference})} \quad (1.1)$$

The aspect ratio of an IA has been shown to be correlated with levels of hemodynamic stressors and has been used as an ease-of-use method to assess conditions within an IA [272].

Aneurysm Hemodynamic Characterisitcs

Wall Shear Stress: The calculation of wall shear stress (WSS) is performed by the ANSYS-FLUENT commercial finite-element solver (ANSYS v17.0). The value is defined as the normal velocity gradient against the (vessel) wall:

$$\tau_w = \mu \frac{\partial v}{\partial n} \quad (1.2)$$

with μ as the fluid dynamic viscosity (0.004 kg/m-s).

The spatial-temporally averaged value of the aneurysm's WSS was calculated alongside its temporally-averaged WSS minimum and temporally-averaged WSS maximum. In a similar manner as IA volume, research differs on whether high [69] or low [273] wall shear stress is a better predictive metric for IA rupture potential. In a study by Meng et. al., both high and low WSS were associated with IA rupture potential, yet causing differing cellular changes [169].

Kinetic Energy Density: The kinetic energy density (KED) within the IA dome was calculated as follows:

$$KED = \frac{\frac{1}{2}\rho \sum v^2}{n} \quad (1.3)$$

Where v is the velocity values, ρ is the mass density of blood, and n is the number of voxels within the IA. The KED at each time-step (along the cardiac phase) was calculated, as well as the Temporally averaged KED (TA-KED) for all cases.

Disturbed Flow on Vascular Endothelium

The vascular endothelial cell (EC) layer forms the innermost lining of blood vessels, directly interacting with hemodynamic stressors and helping to maintain homeostatic functions of the vasculature[51, 125]. The mechanotransduction capabilities of this initial vascular layer help maintain a selective macromolecular barrier, trigger vascular remodeling, regulate vascular smooth muscle cell contraction[242], and

help control vascular inflammatory responses[43]. The degradation of vascular homeostasis, resultant from disturbed hemodynamic flow patterns, has been associated with an array of vascular pathologies: aneurysms[39, 156], atherosclerosis[155], and thrombosis[53, 239]. Due to the life threatening nature of IAs, improved quantitative methods to characterize hemodynamic patterns and to what degree they impart EC pathologic changes, could prove essential to further our understanding of the disease's initiation and progression.

The morphology and cytoskeletal organization of EC have been shown to be susceptible to non-laminar flow conditions[249]. Typically, EC morphology aligns along flow directionality, forming organized parallel actin stress fibers and giving the cells an elongated structure[17, 125, 230]. Disrupted flow patterns resulting in vortex flow and altered WSS, show a differential change in EC characteristics: a rounded morphology with marginally located short actin stress fibers[53, 70, 239]. These changes have been associated with a number of structural-functional changes in vascular cells, such as increased permeability to macromolecules, increased expression of adhesion molecules (ICAM-1, VCAM-1), decreased endothelial cell regeneration and increased smooth muscle cell proliferation/migration.

Additionally, inflammatory processes within vasculature has been shown to be a significant actor in the pathogenesis of IA development and potential rupture [43, 107, 215]. In a typical physiological setting, the vascular EC layer maintains antiatherogenic

characteristics, inhibiting platelet adhesion and aggregation along the vascular wall, as well as limiting cellular pro-inflammatory pathways[4]. In the occurrence of IA pathology, a breakdown of the EC inflammatory-limiting capabilities is noted: small aneurysm shown to have intimal thickening and diffuse macrophage/lymphocyte infiltration, whereas chronic atherosclerotic lesions with embedded macrophages and lymphocytes have been noted in larger aneurysms[142, 238]. Upon leukocyte and macrophage infiltration, the matrix metalloproteinase enzyme is released which digests extracellular matrix proteins leading to additional pathologic damage to the vascular wall[7, 234]. The remodeling of the vascular wall, impart due to inflammatory pathogenic activities, lead to an overall loss vessel mechanical strength and a possible ballooning out of the impacted area

Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros efficiantur. Eos id voluptaria efficiendi comprehensam. [73, 74]




Figure 1.6: Schematic representation of a water molecule

In mel modo dicam vocibus, eruditi consecetuer vim no, cu quaestio instructor eum.
Justo nostrud fuisset ea mea, eam an libris repudiandae vituperatoribus. Est choro
corrumpit definitionem at. Vel sint adhuc vocibus ea, illud epicuri eos no. Sea simul
officiis ea, et qui veri invidunt appellantur. Vix et eros ancillae pertinax. [84, 93,
94, 110, 190] Per iudico probatus complectitur et, cum tollit atomorum rationibus ea.
Per iudico probatus complectitur et, cum tollit atomorum rationibus ea.

Aliquip lobortis ei est, at error viris graeco sed. Vel te elitr detracto, modo graecis
scripserit ex nec. Errem utamur viderer per no, eam ea eripuit referrentur. Pro te di-
cat disputando. Per iudico probatus complectitur et, cum tollit atomorum rationibus
ea. [183, 196, 201, 217]. Per iudico probatus complectitur et, cum tollit atomorum
rationibus ea.

Per iudico probatus complectitur et, cum tollit atomorum rationibus ea. Docendi
eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Per
iudico probatus complectitur et, cum tollit atomorum rationibus ea.

Chapter 2

Cerebral Vasculature and the Impact of Disturbed Flow

The human vasculature is a system aimed at carrying blood and lymph through the body. The arteries of the vascular system help deliver oxygenated blood, nutrients, and components such as inflammatory markers and hormones through the body, while the venous structures help take cellular and tissue waste matter to organs such as the lungs, liver and kidneys for removal as well as carrying deoxygenated blood back to the heart. The lymphatic vessels transport lymph (fluid containing water and blood cells) to help maintain hemodynamic pressures within the body. For the purpose of studying IA, the arterial portion of the vasculature will be focused on in this work as the majority of IAs occur within the arterial system. The innermost layer of arteries,

known as the tunica intima, is made of a monolayer of endothelial cells (EC) supported by a layer of collagen and elastin. This intimal layer comes into direct contact with the hemodynamic flow environment of the lumen (hollow cavity of the arterial system in which blood flows). Underlying the intima layer is the the tunica media, or media, a layer composed of smooth muscle cells, elastic connective tissue and collagen fibers. The main purpose of the media layer is to contract or dilate the arterial vasculature in response to (as signaled by the intima layer) differing hemodynamic conditions as a means to regulate circulation within the body[26, 170]. The outermost layer of arteries, the tunica externa/adventitia, is composed of collagen fibers and elastic tissue helping to maintain the mechanical properties of the vasculature, while the collagen having a secondary purpose of anchoring the vessel to surrounding tissues (improving vessel stability)(Fig. 4.1). In terms of vascular changes concerning aneurysm development and rupture, most focus is on changes to the tunica intima's ECs, and the vascular smooth muscle cells (vSMC) of the tunica media.

Endothelial Cells

ECs are specialized cells that are directly exposed to hemodynamic environment and its circulating components, with a basolateral surface separated from surrounding tissues by a glycoprotein membrane. Due to ECs direct contact with the hemodynamic

The Structure of an Artery Wall




Figure 2.1: Layers of the vasculature. Original image from [218]

environment, their ability to respond to changes in flow conditions is of primary importance. Cytoskeleton actin stress fibers help resist fluid stressors, maintain cellular shape, and aid in cellular mechanotransduction. In laminar flow conditions, actin fibers maintain an oblong orientation, aligning themselves with the direction of flow and forming parallel fibers across the cell to minimize the shear stress exerted by the flowing blood. In addition, ECs act as mechanotransducers, converting mechanical flow stressors into chemical-cellular signals triggering a variety of cellular processes within the vasculature[48]: Cell-cell junctions[75], caveolae[212], integrins[51] and the endothelial glycocalyx[35]. Platelet endothelial cell adhesion molecules (PECAM-1) is shown to create bound complexes with integrins[52, 176], vascular endothelial growth factor receptor 2(VEGFR-2) and VE-cadherin to transmit shear stressors to cellular

responses[46]. Activated integrins trigger multiple complex of non-receptor tyrosine kinases (FAK, c-Src, Shc, paxillin, and p130CAS), adaptor proteins (Grb2, Crk), and guanine nucleotide exchange factors (Sos, C3G), thereby activating Ras family GTPase.2.2. EC glycocalyx, membrane-bound proteoglycans, also take part in sensing and transduction mechanical forces into biochemical signals[56, 171?]. Under physiologic laminar flow conditions, glycocalyx triggers an increase in eNOS expression and subsequent NO production[35, 238, 271]. NO helps regulate vascular dilation/tone and limits vascular inflammation through the inhibition of NF- κ B[42, 165] which in turn limits the expression of intercellular adhesion molecule-1 (ICAM-1), a membrane protein involved in vascular adhesion and migration of leukocytes.

Another role of vascular ECs is the establishment and maintenance of a selective permeability layer within the vasculature[33, 54]. The outward adventitial interstitial pressure (outside the vasculature) is low, roughly 10mmHg, with a much greater intraluminal arterial blood pressure at 130/80mmHg (in humans)[219]. This pressure gradient would create a prominent unidirectional, unchecked mass transport of molecules from the lumen of the vasculature into the surrounding media layer without an additional permeability regulator. The internal EC monolayer helps control macromolecule transport. Typically, maintenance of EC intercellular junctions helps limit macromolecules such as lipoproteins and leukocytes from passing into the intimal layer[125, 178]. The main groups of cell-cell complexes that control permeability are adherens junctions**NEED CITATION** and tight junctions[274], membrane proteins




Figure 2.2: Mechanoreceptors of ECs: ion channels (K^+ , Ca^{2+} , Na^+ , Cl^-), G-proteins, caveolae, tryrosine kinase receptors (TKRs), nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, xanthine oxidase (XO), integrins, and heparan sulfate proteoglycan. Signals are transmitted through the cytoskeleton to the basal or junctional endothelial surface. Integrin mechanosensory complexes consist of platelet endothelial cell adhesion molecule-1 (PECAM-1) and Flk-1. When activated they initiate downstream signaling cascades. Activated integrins trigger multiple complex of non-receptor tyrosine kinases (FAK, c-Src, Shc, paxillin, and p130CAS), adaptor proteins (Grb2, Crk), and guanine nucleotide exchange factors (Sos, C3G), thereby activating Ras family GTPase. Active Ras plays a pivotal role in intracellular transduction of signals as it triggers various parallel downstream cascades of serine kinases; each of these activate downstream signals, ultimately activating mitogen-activated protein kinases (MAPKs). Shear stressors also activate a number of other downstream signaling pathways that impact reactive oxygen species (ROS): NADPH oxidase, activation of protein kinase C (PKC), activation of Rho family small GTPases (mediate the EC remodeling), release of endothelial nitric oxide synthase (eNOS) and activation of phosphoinositide-3 kinase (PI3K)-Akt cascade. These signaling pathways lead to phosphorylation of transcription factors (TFs) such as nuclear factor-kappa(NF- $\alpha\beta$) and activator protein-1(AP-1). Original figure from [48]

connected to the cytoskeleton through transmembrane and cytosolic proteins. The proteins of claudins and occludins make up a significant portion of the tight junction between cells [47, 274]. Tight junctions prevent the passage of molecules and ions through the space between plasma membranes of adjacent cells, so materials must

pass enter the cells (by diffusion or active transport) in order to pass through the tissue helping to maintain osmotic balance within the vasculature. Adherens junctions such as vascular endothelial (VE)-cadherin also help cell-cell adhesion (and subsequent EC permeability) well as maintaining an inhibition of endothelial cell growth. A breakdown in these connections leading to *decreased* regulation of permeability within the EC monolayer as 'gaps' now exist between cells, allowing the greater degree of macromolecule infiltration to subsequent layers of the vasculature. While not directly tasked with the maintenance of EC permeability, the regulation of cellular turnover and/or apoptosis has a secondary impact on vascular permeability[195]. Areas of apoptotic cells, or areas of cellular turnover can change the permeability of the vasculature.

Activated integrins such as β -1 integrin and integrin α 5, transmits additional cellular signaling to proteins such as Src homology domain 2-containing kinase (Shc), which can activate NF- κ B causing subsequent increases cell proliferation, apoptotic signaling through Jun-amino-terminal kinase (JNK) and altered cellular response to cytokines through p38 mitogen-activated protein kinase (MAPK). Activated NF- κ B also triggers downstream activation of a number of genes related to macrophage recruitment and vascular inflammation: MCP1, MMP2, MMP9, IL1 β , and inducible nitric oxide synthase(iNOS).

The tunica media of the vasculature is primarily made up of vSMC and produce

a significant portion of the vascular extracellular matrix proteins (elastin, collagen, proteoglycans, and fibrillin). This multi-layered portion of the arterial system help regulate blood vessel integrity and vascular tone in response to hemodynamic forces and are key components in maintaining contractile functionality. Under physiologic conditions, vSMC maintain a nonproliferative phenotype with an abundance of contractile fibers, intermediate filaments, microtubules and actin. Intermediate filaments such as vimentin and desmin aid in the maintenance of vSMC structure[31], whereas microtubules **DO WHAT**. Actin filaments transmit mechanical signals dispersed within the vSMC cytoplasm and aid in the transmission of cellular signalling[24]. Changes in cellular tension alter actin cytoskeletal organization, regulating cell contraction, migration and cellular viability[33]. The myosin components within vSMC aid in cellular contraction (similar to actin fibers) and are shown to be altered in the event of vascular injury.

Impact of Disturbed Flow on Vascular Cells

Broadly speaking, the change from a healthy artery to aneurysm development and possible rupture follow an (assumed) overall path: alterations to the ECs, increased macrophage infiltration into the vessel wall, breakdown of proteins (proteolysis) in the extracellular matrix[197, 204], loss of internal elastic lamina[203, 253], and decreased collagen synthesis[135, 154]. Continued pathophysiological conditions can lead to

apoptosis of vSMC causing continued weakening of the vascular wall and possible aneurysm formation. In the event of prolonged degradation to the aneurysm wall, an intraluminal thrombosis[66, 132, 150] or aneurysm rupture may occur.




Figure 2.3: Schematic representation of abdominal aortic aneurysm (AAA) pathophysiology. Inflammation, proteolysis (breakdown of proteins by enzymes), smooth muscle cell (SMC) apoptosis, neovascularization, calcification, and intraluminal thrombosis may be targeted by molecular imaging. ECM indicates extracellular matrix; and MMP, matrix metalloproteinase.

As mentioned, laminar flow promotes the oblong orientation of actin stress fibers aligned with the direction of flow across the cell in ECs. Low WSS (< 5 dynes/cm²) disturbed flow conditions generated in in vitro, result in altered cytoskeletal orientation with actin fibers forming non-random accumulation at cell peripheries[17, 149, 261] with ECs taking on a round or polygonal shape. Additionally, changes in structural matrix smooth muscle actin expression is reduced in areas of

disturbed flow, with lower actin expression in aneurysms versus healthy vessels and further reduction in actin expression within ruptured versus unruptured [?]. These alterations to EC morphology

Table 2.1
Disturbed Flow's Impact on Vascular Cells

Cellular Functionality	Resultant Change due to Disturbed Flow	Model Stimulus	Reference
Cellular component/protein			
Endothelial permeability			
VEGF	Increased	<i>In-vivo</i> : CaCl ₂ induction	[9]
VEGFR2	Increased	<i>In-vitro</i> : Orbital Shaker	[152]
VE-cadherin	Decreased	<i>In-vivo</i> : ligated arterial segment	[52]
	Decreased	<i>In-vitro</i> : Magnetic twisting	[235]
Atherosclerosis			
PECAM-1	Increased	<i>In-vivo</i> : ligated arterial segment	[52]
Mechanotransduction			
β 1-Integrin	Increased	<i>In-vitro</i> : PPFC	[262]
Integrin α 5	Increased	<i>In-vitro</i> : Orbital Shaker	[227]
Glycocalyx	Decreased	<i>In-vitro</i> : PPFC <i>In-vivo</i> : Arterial Ligation	[104]
	Decreased	<i>In-vivo</i> : Arterial Ligation	[171]
	Decreased	<i>In-vitro</i> : Curved flow chamber	[56]
Inflammatory Response			
ICAM-1	Increase	<i>In-vitro</i> : PPFC with occlusion	[17]
	Increased	<i>In-vivo</i> : Arterial Ligation	[211]
	Increased	<i>In-vitro</i> : Cone-plate viscometer	[96]
	Increased	<i>In-vitro</i> : PPFC	[16]
VCAM-1	Increased	<i>In-vivo</i> : Arterial Ligation	[262]
	Increased	<i>In-vitro</i> : PPFC	[16]
ECM Remodeling			
MMP2	Increased	<i>In-vitro</i> : PPFC with occlusion	[17]
	Increased	<i>In-vitro</i> : Stenosis flow chamber	[57]
MMP-9	Increased		[56]
Oxidative Stress / Reactive Oxygen Species			
NOX	Increased	<i>In-vivo</i> : genetic knockout/Ang II infusion	[216]
IL-6	Increased	<i>In-vivo</i> : Arterial Ligation	[211]
	Increased	<i>In-vitro</i> : Cone-plate viscometer	[96]
IL-1 β	Increased	<i>In-vitro</i> : Cone-plate viscometer	[186]
IL-5	Increased		[18]
eNOS	Decreased	<i>In-vitro</i> : PPFC with occlusion	[104]
	Decreased	<i>In-vitro</i> : Cone-plate viscometer	[24]
iNOS	Decreased	<i>In-vitro</i> : Orbital Shaker	[161]
Apoptosis			
Caspase 12	Increase	<i>In-vitro</i> : Cone-plate viscometer	[186]
Caspase 3	Increase	<i>In-vitro</i> : Cone-plate viscometer	[109]
Transcription Factors			
NF- κ B	Increased	<i>In-vitro</i> : PPFC	[14]
	Increased	<i>In-vitro</i> : Cone-plate viscometer	[96]
Nrf2	Decreased	<i>In-vitro</i> : Orbital Shaker	[163]
JNK	Increased	<i>In-vivo</i> : Arterial ligation	[251]
	Increased	<i>In-vitro</i> : Oscillating stretch of cells on silicone foundation	[55]

Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros efficiantur. Eos id voluptaria efficiendi comprehensam.

In mel modo dicam vocibus, eruditi consecetuer vim no, cu quaestio instructor eum.
 Justo nostrud fuisset ea mea, eam an libris repudiandae vituperatoribus. Est choro
 corrumpit definitionem at. Vel sint adhuc vocibus ea, illud epicuri eos no. Sea simul
 officiis ea, et qui veri invidunt appellantur. Vix et eros ancillae pertinax.

Aliquip lobortis ei est, at error viris graeco sed. Vel te elitr detracto, modo graecis
 scripserit ex nec. Errem utamur viderer per no, eam ea eripuit referrentur. Pro te
 dicat disputando.

Table 2.2

A portrait table: first column represents the year in which the Nobel prize
 in physics was awarded; second column indicates the name of the scientist
 and the third column is the work for which the Nobel prize was awarded

Year	Scientist(s)	Nobel Work
1901	W. C. Röntgen	X-rays
1902	H. A. Lorentz	Influence of magnetism on radiation
	P. Zeeman	Influence of magnetism on radiation
1903	A. H. Becquerel	Spontaneous radioactivity
	M. Curie	Radiation phenomena discovered by Becquerel
	P. Curie	Radiation phenomena discovered by Becquerel
1904	J. W. Strutt	Argon
1905	P. E. A. von Lenard	Cathode rays
1906	J. J. Thomson	Electrical conductivity of gases
1907	A. A. Michelson	Spectroscopic and metrological investigations
1908	G. Lippmann	Photographic reproduction of colours
1909	K. F. Braun	Wireless telegraphy
	G. Marconi	Wireless telegraphy
1910	J. D. van der Waals	Equation of state of gases and liquids
1911	W. Wien	Laws governing heat radiation
1912	N. G. Dalèn	Automatic regulators for lighting coastal beacons and light buoys

As explained in Table 2.2, Ex offendit elaboraret cum has ex natum honestatis, impedit similique ex duo. Et mei mollis scripta, et vim labores phaedrum, in cum facete saperet. Splendide elaboraret comprehensam qui ne. Putant verterem no vim, mea solum veritus definitiones ei, no labitur propriae deseruisse est. Ius illud everti salutandi id, eu facer pericula principes est.




Figure 2.4: Fancy mathematical plots using TikZ package

Simul noster voluptaria eam ei, sint regione pri ei. Cum no utinam equidem, falli bonorum prodesset an qui. Alterum dissentiet vituperatoribus te eam, eos ea suas oblique. Per ea utinam facilisi. Per iudico probatus complectitur et, cum tollit atomorum rationibus ea.

Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros efficiantur. Eos id voluptaria efficiendi comprehensam.

Simul noster voluptaria eam ei, sint regione pri ei. Cum no utinam equidem, falli bonorum prodesset an qui. Alterum dissentiet vituperatoribus te eam, eos ea suas oblique. Per ea utinam facilisi. Per iudico probatus complectitur et, cum tollit atomorum rationibus ea.




Figure 2.5: Incidence, transmission and reflection

Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros efficiantur. Eos id voluptaria efficiendi comprehensam. Simul noster voluptaria eam ei, sint regione pri ei. Cum no utinam equidem, falli bonorum prodesset an qui.

Chapter 3

Hemodynamic Flow Vortex

Identification

The assessment of disturbed hemodynamic patterns is known to have an impact on the origin and natural history of IAs [30, 255]. From a clinical perspective, phase-contrast magnetic resonance imaging (PC-MRI) or Phase-contrast magnetic resonance angiography (PC-MRA) has been used to assess flow characteristics in the vasculature *in-vivo* [22, 168]. Yet, determining flow details in and around IAs has proven difficult with PC-MRI/PC-MRA. The complex, disturbed patterns of aneurysmal flow results in incoherent velocities (at the sub-grid level) and these specific characteristics cannot be resolved by a typical "averaged" velocity measurement at the relatively large resolution in medical imaging (*at 1-mm scale*). The consequence of

this sub-grid limitation, clinical hemodynamic flow measurements may be impacted by errors and potential flow artifacts which adversely affect the accuracy of PC-MRI/PC-MRA results [168, 224]. In parallel to research efforts of assessing MR flow imaging in and around IAs to determine rupture characteristics, blood flow simulated from "patient-specific" computational fluid dynamics (CFD) simulations [222] have also garnered interest by the clinical and research community [40, 263]. CFD simulation data has an initial advantage over MRI derived data in that a high degree of control of both desired data resolution, as well as data quality (lack of imaging errors and flow artifacts) can be maintained. The development of novel flow measurement techniques can be initially tested and refined on data free from errors which may confound initial findings.

Assessment of hemodynamic conditions from CFD simulations has brought about a number of potential parameters that correlate to IA rupture risk: wall shear stress [13], oscillatory shear index [223], flow impingement [40], and flow stability [32]. As mentioned in the focus of flow stability, Bryne et al. [32] found that aneurysmal flow flow spatial complexity and temporal stability is closely correlated with IA rupture. Yet this methodology only relies on qualitative assessment of the centroid-most region of vortex patterns (vortex core), giving less insight on the broader structural changes to vortices. Expanding upon the idea of vortex analysis to identify, and quantify changes to, the broader structural characteristics of vortices may give additional insight into conditions that can be linked to IA rupture potential.

In an initial study, the development of an alternative technique for vortex analysis (expanding upon vortex core analysis) was investigated to assess the presence, destruction and spatiotemporal characteristics of vortices within an IA [228]. This image processing algorithm expanded upon two established vortex identification methods, the Q -criterion [116] and λ_2 [120] methods, to identify and assess the broader aspects of vortices as opposed to solely identifying the vortex core. Subsequent studies investigated the use of a vortex identification methodology based on the Shannon's entropy (CITATION NEEDED) as an alternative identification metric not wholly reliant upon vortex core identification.

Due to the differences between traditional CFD data resolution (high) and MRI derived flow data (lower resolution), the identification methods were based on velocity data on a rectilinear grid, and the susceptibility of changes to vortex identification outcomes were tested under a range of grid resolutions. Additionally, variations to methodological outputs were tested under a range of chosen threshold values (dependent on the method and will be explained in Section NEED THE SECTION). The vulnerability of a methodology to significant changes in outcomes with minimal changes to methodological threshold values would have the potential to result in broad variations to research findings if thresholds were to be applied to a wide array of studies. Toward this end, the primary focus of this work was to explore analyzing the spatiotemporal characteristics of hemodynamic vortex structures as a possible means to compliment future assessment of IA rupture potential.

Materials and Methods

Modeling of "Patient-specific" Vasculature

Medical imaging scans of ten (10) patient's vasculature structure were arbitrarily selected from an internal database: five patients with a single bifurcation aneurysm, and five with a single sidewall aneurysm. Models were located within the internal carotid artery or the basilar artery. A commercially available image segmentation package (Mimics Innovation Suite, version 17, Materialise Inc. Leuven, Belgium) was used to reconstruct the vascular surface from digital subtraction angiography (DSA) scans, resulting in 'patient-specific' vascular structures. For all cases, the longest available upstream vessel section proximal to the aneurysm was left intact to maintain as much of the patient vessel geometry as possible. Surface irregularities were manually removed using the localized smoothing capability in the commercially available computer aided design (CAD) 3-matic software (Version 9, Materialize Inc., Leuven, Belgium). Additionally, a 1st order Laplacian smoothing filter was used to perform a global smoothing to the vessel structure, reducing surface irregularities while preserving the vascular geometry. Cylindrical flow extensions (6 times the inlet vessel diameter) were added to each model using the open-source Vascular Modeling Toolkit (VMTK) software (version 1.2). The addition of vessel extensions help reduce

the effects of inlet, plug-flow flow on hemodynamic characteristics [191].

Mesh Generation

Processed vascular surface structures were converted into an unstructured, 3D, tetrahedralized volumetric mesh using an open-source mesh generator, Tetgen (version 1.4.2) [214]. The mesh generation process was done by an in-house Python script derived from the VMTK program. Approximately, 1 million computing cells were used per case, with the average mesh size as 0.0022-mm^3 .

CFD Simulation

To compute fluid velocity data in and around the IA, the time-dependent incompressible, 3D Navier-Stokes equations was (THERE NEEDS TO BE AN ADJECTIVE HERE....CHECK RECENT PAPERS) solved using two CFD solvers: a commercial CFD solved (version 14.0, ANSYS-FLUENT Inc., Lebanon, NH) and a research prototype CFD solver (version 4.0, Siemens Medical Solution Inc., IL). Details on the Navier-Stokes equation can be found in the Appendix. In the ANSYS-FLUENT solver, the pressure-velocity coupling was obtained using the SIMPLEC algorithm [246]. The explicit time-marching second-order scheme with a time step of 1×10^{-3}

second (1000 steps per cardiac cycle) was used for computations.

As the Siemens research CFD solver is still under development, limited information on its chosen methods for solving the Navier-Stokes equation will be discussed in this thesis. IA models were defined by water-tight 3D surface triangles were automatically discretized with cubical voxels. A Lattice-Boltzmann Method (LBM) solver was then used to choose adaptively choose the solver time-step, and varied from 1×10^{-3} to 2×10^{-3} . A Siemens Leonardo workstation equipped with a dual quad-core CPU and 8 GB of memory was used to perform CFD simulations. Of note, the exact same vessel geometries (STL files) were used to generate the volumetric meshes (for the ANSYS-FLUENT solver) and voxel discretization (for the LBM solver). The final velocity results obtained from the LBM method were re-sampled to form velocity data onto a rectilinear grid whose voxel size varied from 0.18 to 0.25 mm.

In both solvers, vessels walls were assumed rigid with a no-slip boundary. While blood vessels are not rigid, the assumed rigidity for simulation has been shown to cause a rise in overall wall shear stress values, but have minimal impact on flow pattern characteristics in the arterial system unless the microvasculature is being simulated. Blood was considered an incompressible and Newtonian fluid with a dynamic viscosity of 0.004 kg/m-s and a mass density of 1050 kg/m³. A zero-pressure condition was used for all vessel outlets. For inlet flow rates, two pulsatile waveforms at a rate of 60 bpm were derived from magnetic resonance measurements and were taken from Gwilliam

et al. [99] as patient-specific flow waveforms were not available. Each case had its inlet waveform scaled according to their inlet cross-sectional area, standardizing their mean volumetric flow rate to either 280mL/min for ICA cases or 180mL/min for BAs. This choice of volumetric flow rate(s) were based on measured physiological flow rates available in MR literature [82, 275]. Four (4) cardiac cycles were simulated per case at 20 data points per cardiac cycle with only the final cardiac cycle saved as a means to reduce initial transient flow conditions.

Aneurysm Extraction and Voxelization of Aneurysmal Velocity Data

A published method [121] was used to semi-automatically isolate and extract the IA sac. The isolated IA sac was sealed at the IA opening (ostium) and converted to a binary mask that is spatially-registered with the volumetric velocity data. The mask allows the analysis of only the intra-aneurysmal velocity data. To verify intra-rater reliability of proper sectioning of IA masks, 2 separate users sectioned the IAS and Bland-Altman plots were performed on the resultant mask volumes and ostium areas to determine the similarity between chosen masks. Once no significant differences were ensured between sectioned masks, one user was chosen at random and all resultant masks from that user were implemented in the rest of the study.

Vortex Core Extraction and Analysis

All computational methods for identification and extraction of vortices and spatio-temporal analysis of said vortices were performed using in-house scripts (C++ and Python) that were derived from the open-source VTK/VMTK software package.

This flow assessment was performed through the analysis of vortex critical point (core) lines and it was concluded that "ruptured aneurysms had more complex and more unstable flow patterns than un-ruptured aneurysms." In their work, Bryne used proper orthogonal decomposition [] of time-resolved velocities were used to characterize temporal flow stability. As a brief explanation, vortex core lines, identifying the center-most region of vortex pattern, were identified by the use of a co-linearity conditions between flow instantaneous vorticity $\vec{\omega}$ and velocity \vec{v} vectors.

$$\vec{\omega} \times \vec{v} = 0 \tag{3.1}$$

$$\vec{\omega} = \nabla \times \vec{v}$$

To identify the centroid region of vortices, the eigenvalues of the velocity gradient tensor was calculated. In the event of a pair of complex conjugate eigenvalues was identified, the vorticity vector $\vec{\omega}$ as calculated and tested against the velocity vector to assess whether Equation 3.1 was satisfied. From an identified element, the velocity

component in the direction of vorticity vector was subtracted from the velocity vector (reduced velocities). Element faces that had a point where the reduced velocity is zero was marked, and if two or more faces of an element had a zero reduced velocity, a vortex core line passes through the element.

Vortex core line analysis

$$\begin{aligned}\nabla \vec{u} &= S + \Omega \\ S &= \frac{1}{2} [(\nabla \vec{u}) + (\nabla \vec{u})^T] \\ \Omega &= \left[\frac{1}{2} (\nabla \vec{u}) - (\nabla \vec{u})^T \right]\end{aligned}\tag{3.2}$$

Where $\nabla \vec{u}$ is the calculation of the velocity gradient: S as the rate-of-strain tensor and Ω as the vorticity tensor.

Hunt, Wray and Moin [116] defined a vortex as the spatial region of flow where the Euclidean norm of the vorticity tensor dominates.

$$Q = \frac{1}{2} [|\Omega|^2 - |S|^2] > 0\tag{3.3}$$

Jeong and Hussain identified the vortices as:

$$\lambda_2 = (S^2 + \Omega^2) < 0\tag{3.4}$$

where $\lambda_2 A$ identifies a vortex when the second intermediate eigenvalue of the 3×3 tensor A is symmetric (all three eigenvalues are real).

In our original study, the normalized Q and λ_2 values were tested to identify vortices within IAs.

$$\begin{aligned} Q(x, t) &= \frac{Q(x, t)}{|\vec{u}(x, t)|^2} \\ \lambda_2(x, t) &= \frac{\lambda_2(x, t)}{|\vec{u}(x, t)|^2} \end{aligned} \tag{3.5}$$

Lorem ipsum dolor sit amet, at qui viderer recusabo aliquando, dignissim evertitur ei his. Ignota iuaret fabulas ei vim. Ne utinam inciderint quo. Pri ea congrue postulant conclusionemque. In prima quaeque diceret pri. Enim labores contentiones eos at, duo altera denique nominavi ea, eos inani nominavi consecetuer at. Ut elit dicam elaboraret pro, ius altera voluptaria cu.

Discere dissentiet vel et, soluta nostrum epicurei ad eam, cu has aperiam vituperata. In prima quaeque diceret pri. Enim labores contentiones eos at, duo altera denique nominavi ea, eos inani nominavi consecetuer at. Ut elit dicam elaboraret pro, ius altera voluptaria cu. Eam mazim aliquip cu, recusabo pericula accommodare at mea, facer affert nonumes qui ea. [5, 90]

$$\begin{aligned}
d\nu_\theta &= \frac{N}{V} \left(\frac{m}{2\pi kT} \right)^{3/2} \left[\int_0^{2\pi} \int_0^\infty v^3 e^{-mv^2/2kT} dv d\phi \right] \sin \theta \cos \theta d\theta \\
&= 2\pi \frac{N}{V} \left(\frac{m}{2\pi kT} \right)^{3/2} \left[\int_0^\infty v^3 e^{-mv^2/2kT} dv \right] \sin \theta \cos \theta d\theta
\end{aligned}$$

At vix indoctum disputando. Eam cu doctus reprimique, quaeque democritum an eos, sit veniam facete dissentias id. Tale volumus eos te, an eum nulla tincidunt. Mea id recteque theophrastus.

$$d\nu_\theta = \frac{N}{V} \left(\frac{2kT}{m\pi} \right)^{1/2} \sin \theta \cos \theta d\theta \quad (3.6)$$

Liber liberavisse nec at, movet albucius principes has at. Ea sed persius accusam, clita sententiae adversarium ne sed. Usu no graecis theophrastus delicatissimi, sint aliquam an eam. Mei elit mnesarchum dissentias te, in essent laboramus per. Affert mucius quidam mel ex, per dicam insolens ad.

Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros efficiantur. Eos id voluptaria efficiendi comprehensam. Continuing from Eqn. (3.6)

$$\begin{aligned}
d\nu_v &= \frac{N}{V} \left(\frac{m}{2\pi kT} \right)^{3/2} \left[\int_0^{2\pi} \int_0^{\pi/2} \sin \theta \cos \theta d\theta d\phi \right] v^3 e^{-mv^2/2kT} dv \\
&= 2\pi \frac{N}{V} \left(\frac{m}{2\pi kT} \right)^{3/2} \left[\int_0^{\pi/2} \sin \theta \cos \theta d\theta \right] v^3 e^{-mv^2/2kT} dv
\end{aligned}$$

In mel modo dicam vocibus, eruditi consecetuer vim no, cu quaestio instructor eum. Justo nostrud fuisset ea mea, eam an libris repudiandae vituperatoribus. Est choro corrumpit definitionem at. Vel sint adhuc vocibus ea, illud epicuri eos no. Sea simul officiis ea, et qui veri invidunt appellantur. Vix et eros ancillae pertinax.

In mel modo dicam vocibus, eruditi consecetuer vim no, cu quaestio instructor eum. Justo nostrud fuisset ea mea, eam an libris repudiandae vituperatoribus. Est choro corrumpit definitionem at. Vel sint adhuc vocibus ea, illud epicuri eos no. Sea simul officiis ea, et qui veri invidunt appellantur. Vix et eros ancillae pertinax.

$$d\nu_v = \frac{N}{V} \pi \left(\frac{m}{2\pi kT} \right)^{3/2} v^3 e^{-mv^2/2kT} dv \quad (3.7)$$

Aliquip lobortis ei est, at error viris graeco sed. Vel te elitr detracto, modo graecis scripserit ex nec. Errem utamur viderer per no, eam ea eripuit referrentur. Pro te dicat disputando.




Figure 3.1: Distribution of random numbers

Table 3.1

Measured data points representing the relationship between x and y

x	0	1	2	3	4	5	6	7	8	9	10
y	0	0.94	0.99	-0.52	-1.82	-0.44	3.54	6.69	5.38	0.00	-4.42

Et mei mollis scripta, et vim labores phaedrum, in cum facete saperet. Splendide elaboraret comprehensam qui ne. Putant verterem no vim, mea solum veritus definitiones ei, no labitur propriae deseruisse est. Ius illud everti salutandi id, eu facer pericula principes est.

Table 3.2

A landscape table: first column represents the year in which the Nobel prize in physics was awarded; second column indicates the name of the scientist and the third column is an *as is* Nobel citation

Year	Scientist(s)	Nobel Work
1901	W. C. Röntgen	in recognition of the extraordinary services he has rendered by the discovery of the remarkable rays subsequently named after him
1902	H. A. Lorentz and P. Zeeman	in recognition of the extraordinary service they rendered by their researches into the influence of magnetism upon radiation phenomena
1903	A. H. Becquerel	in recognition of the extraordinary services he has rendered by his discovery of spontaneous radioactivity
	M. Curie and P. Curie	in recognition of the extraordinary services they have rendered by their joint researches on the radiation phenomena discovered by Prof. Henri Becquerel
1904	J. W. Strutt	for his investigations of the densities of the most important gases and for his discover argon in connection with these studies
1905	P. E. A. von Lenard	Cathode rays
1906	J. J. Thomson	Electrical conductivity of gases
1907	A. A. Michelson	Spectroscopic and metrological investigations
1908	G. Lippmann	Photographic reproduction of colours
1909	K. F. Braun and G. Marconi	Wireless telegraphy
1910	J. D. van der Waals	Equation of state of gases and liquids
1911	W. Wien	Laws governing heat radiation
1912	N. G. Dalèn	Automatic regulators for lighting coastal beacons and light buoys

Et mei mollis scripta, et vim labores phaedrum, in cum facete saperet. Splendide elaboraret comprehensam qui ne. Putant verterem no vim, mea solum veritus definitiones ei, no labitur propriae deseruisse est. Ius illud everti salutandi id, eu facer pericula principes est.




Figure 3.2: Fibre optics

Simul noster voluptaria eam ei, sint regione pri ei. Cum no utinam equidem, falli bonorum prodesset an qui. Alterum dissentiet vituperatoribus te eam, eos ea suas oblique. Per ea utinam facilisi. Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te.

Adipisci molestiae vim at, eum everti accommodare eu. Duo ex maiorum consetur. Sea et vivendo concludaturque, rebum conclusionemque pro eu. Mei an everti dolorem. Per id alterum mandamus deseruisse. Copiosae evertitur eum ea, atqui interesset est in. Vim magna munere nostrum an, cu congrue equidem est. Mediocre reformidans ne mel. Et summo nihil mel, an nam postea incorrupte.

In amet verear evertitur qui, ex mea vivendo hendrerit. Ad posse perfecto prodesset usu, cum fugit accumsan no. Tempor nonumes duo ea, oblique fabulas salutatus ne vis. Ne eam scripta dolorem, graece eruditi eum ei. Ei sed brute zril nostro, nostro voluptatum id sea, courtesy of Wikipedia. [113] Adipisci molestiae vim at, eum everti accommodare eu. Duo ex maiorum consetetur. Sea et vivendo concludaturque, rebum conclusionemque pro eu.

Adipisci molestiae vim at, eum everti accommodare eu. Duo ex maiorum consetetur. Sea et vivendo concludaturque, rebum conclusionemque pro eu. Mei an everti dolorem. Per id alterum mandamus deseruisse. Copiosae evertitur eum ea, atqui interest est in. Vim magna munere nostrum an, cu congrue equidem est. Mediocre reformidans ne mel. Et summo nihil mel, an nam postea incorrupte an everti dolorem. Per id alterum mandamus deseruisse. Copiosae evertitur eum ea, atqui interest est in. Vim magna munere nostrum an, cu congrue equidem est. Mediocre reformidans ne mel. Et summo nihil mel, an nam postea incorrupte. Mediocre reformidans ne mel. Et summo nihil mel, an nam postea incorrupte an everti dolorem.

Per id alterum mandamus deseruisse. Copiosae evertitur eum ea, atqui interest est in. Vim magna munere nostrum an, cu congrue equidem est. Mediocre reformidans ne mel. Et summo nihil mel, an nam postea incorrupte.




Figure 3.3: A landscape view of a Turboprop engine - these are jet engine derivatives, still gas turbines, that extract work from the hot-exhaust jet to turn a rotating shaft, which is then used to produce thrust by some other means

Id ius soluta semper audiam, ad eos scriptorem concludaturque, id mel rebum volumus deserunt. Mel libris percipit scriptorem te, his an dicat putent menandri, mazim officiis aliquando mei no. Ne clita veniam disputando vim, postea hendrerit maiestatis qui id. Mei te suscipit quaerendum, an aliquando intellegebat ius, ei simul detraxit dissentiet eam. Zril dolor ut usu.

Everti saperet vis ut. Scripta malisset mel eu, duis antiopam in pro. Sea diceret contentiones ea. Nec eu duis efficiantur, evertitur constituam mediocritatem te vis, pro error regione ad. Sit malorum aliquam at, pericula dissentias mei ei. Cu soluta urbanitas est, albucius vituperatoribus usu et.

Chapter 4

Vortex Analysis to predict IA Initiation

The tangential, frictional stress caused by blood flowing along the vessel wall is known as WSS. The ANSYS-FLUENT software calculates WSS by the normal velocity gradient at the vessel wall:

$$\tau_w = \mu \frac{\partial v}{\partial n} \quad (4.1)$$

where μ is the dynamic viscosity. In this work, areas of high WSS were of interest as it is thought to play a role in the IA initiation [169]. High WSS was defined as values ≥ 20 Pa during peak systole of the MRI waveform.

The WSSG was calculated using in-house VMTK scripts and is derived from three

spatial derivatives of the WSS as follows:

$$WSSG = \sqrt{\left(\frac{\partial \tau_w}{\partial x}\right)^2 + \left(\frac{\partial \tau_w}{\partial y}\right)^2 + \left(\frac{\partial \tau_w}{\partial z}\right)^2} \quad (4.2)$$

with the time-averaged WSSG calculated as

$$WSSG_{av} = \frac{1}{T} \int_0^T |WSSG| dt \quad (4.3)$$

OSI is a nondimensional parameter, computing oscillations in the direction of the WSS vectors over the course of a cardiac cycle:

$$OSI = \frac{1}{2} \left\{ 1 - \frac{|\int_0^T \tau_i dt|}{\int_0^T |\tau_i| dt} \right\} \quad (4.4)$$

where τ_i represents the WSS vector at a given time step across the duration of the cardiac cycle (T). The OSI describes the changes of a WSS vector's alignment with the cardiac cycle's temporally-averaged WSS vector. An OSI of 0 indicates no change in directionality and 0.5 being a complete direction reversal.

The AFI [?] quantifies the variation in angle between the instantaneous WSS vector and time-averaged WSS vector:

$$AFI = \cos(\theta) = \frac{\tau_i \cdot \tau_{av}}{|\tau_i| * |\tau_{av}|} \quad (4.5)$$

For each point along the vessel wall, the minimum AFI calculated during the cardiac cycle was used to indicate the greatest deviation of the WSS vector from its mean direction. A minimum AFI of -1, 0, and 1 indicate deviations of 180°, 90°, and 0° respectively.

The GON index [?] quantifies fluctuations in WSSG directionality over the cardiac cycle.

$$GON = 1 - \frac{|\int_0^T G dt|}{\int_0^T |G| dt} \quad (4.6)$$

T is the period of the cardiac cycle and G is the spatial wall shear stress gradient vector

The human vasculature is a system aimed at carrying blood and lymph through the body. The arteries of the vascular system help deliver oxygenated blood, nutrients, and components such as inflammatory markers and hormones through the body, while the venous structures help take cellular and tissue waste matter to organs such as the lungs, liver and kidneys for removal as well as carrying deoxygenated blood back to the heart. The lymphatic vessels carry lymph (fluid containing water and blood cells) to help maintain hemodynamic pressures within the body. Intracranial aneurysms have only been shown to develop within the arterial portion of the vasculature, and as such will be the part of the vasculature focused upon in this work. The innermost layer of arteries, known as the tunica intima, is made of a monolayer of endothelial cells (EC) supported by a layer of collagen and elastin. This intimal layer comes into

direct contact with the hemodynamic flow environment of the lumen (hollow cavity of the arterial system in which blood flows). This endothelial layer serves multiple purposes: hemodynamic mechanosensors, reacting to fluid forces and transducing said force into biochemical signals triggering cellular cascades controlling vascular tone and homeostasis[35, 51, 75], and acting as a selective permeability layer for macromolecules[21, 95, 177]. Underlying the intima layer is the the tunica media, or media, a layer composed of smooth muscle cells, elastic connective tissue and collagen fibers. The main purpose of the media layer is to contract or dilate the arterial vasculature in response to (as signaled by the intima layer) differing hemodynamic conditions as a means to regulate circulation within the body[26, 170]. The outermost layer of arteries, the tunica externa/adventitia, is composed of collagen fibers and elastic tissue helping to maintain the mechanical properties of the vasculature, while the collagen having a secondary purpose of anchoring the vessel to surrounding tissues (improving vessel stability)(Fig. 4.1).

The Structure of an Artery Wall




Figure 4.1: Layers of the vasculature. Original image from [218]

References

- [1] Cost comparison of surgical and endovascular treatment of unruptured giant intracranial aneurysms. *Neurosurgery* 77, 5 (2015), 733–743.
- [2] ABBOUD, T., RUSTOM, J., BESTER, M., CZORLICH, P., VITTORAZZI, E., PINNSCHMIDT, H. O., WESTPHAL, M., AND REGELSBERGER, J. Morphology of ruptured and unruptured intracranial aneurysms. *World neurosurgery* 99 (2017), 610–617.
- [3] AIRD, W. Spatial and temporal dynamics of the endothelium. *Journal of Thrombosis and Haemostasis* 3, 7 (2005), 1392–1406.
- [4] AL-SOUDI, A., KAAIJ, M., AND TAS, S. Endothelial cells: From innocent bystanders to active participants in immune responses. *Autoimmunity Reviews* 16, 9 (2017), 951 – 962.

- [5] ANDERSON, E., BAI, Z., BISCHOF, C., BLACKFORD, S., DEMMEL, J., DONGARRA, J. J., CROZ, J. D., GREENBAUM, A., HAMMARLING, S., MCKENNEY, A., AND SORENSEN, D. *LAPACK Users' Guide*, 3 ed. Society for Industrial and Applied Mathematics, Philadelphia, PA, 1999.
- [6] ANTIGA, L., AND STEINMAN, D. A. Robust and objective decomposition and mapping of bifurcating vessels. *IEEE transactions on medical imaging* 23, 6 (2004), 704–713.
- [7] AOKI, T., KATAOKA, H., MORIMOTO, M., NOZAKI, K., AND HASHIMOTO, N. Macrophage-derived matrix metalloproteinase-2 and -9 promote the progression of cerebral aneurysms in rats. *Stroke* 38, 162–169.
- [8] AOKI, T., YAMAMOTO, K., FUKUDA, M., SHIMOYONIA, Y., FUKUDA, S., AND NARUMIYA, S. Sustained expression of mcp-1 by low wall shear stress loading concomitant with turbulent flow on endothelial cells of intracranial aneurysm. *Acta Neuropathologica Communications* 4, 1 (2016), 48.
- [9] ASHINA, K., TSUBOSAKA, Y., NAKAMURA, T., OMORI, K., KOBAYASHI, K., HORI, M., OZAKI, H., AND MURATA, T. Histamine induces vascular hyperpermeability by increasing blood flow and endothelial barrier disruption in vivo. *Plos one* 10, 7 (2015).

- [10] BACKES, D., RINKEL, G. J., LABAN, K. G., ALGRA, A., AND VERGOUWEN, M. D. Patient- and aneurysm-specific risk factors for intracranial aneurysm growth. *Stroke* 47, 4 (2016), 951–957.
- [11] BACKES, D., RINKEL, G. J., LABAN, K. G., ALGRA, A., AND VERGOUWEN, M. D. Patient- and aneurysm-specific risk factors for intracranial aneurysm growth. *Stroke* 47, 4 (2016), 951–957.
- [12] BACKES, D., VERGOUWEN, M. D., VELTHUIS, B. K., VAN DER SCHAAF, I. C., BOR, A. S. E., ALGRA, A., AND RINKEL, G. J. Difference in aneurysm characteristics between ruptured and unruptured aneurysms in patients with multiple intracranial aneurysms. *Stroke* 45, 5 (2014), 1299–1303.
- [13] BAEK, H., JAYARAMAN, M., RICHARDSON, P., AND KARNIADAKIS, G. Flow instability and wall shear stress variation in intracranial aneurysms. *Journal of the Royal Society Interface* (2009), rsif20090476.
- [14] BAERISWYL, D. C., PRIONISTI, I., PEACH, T., TSOLKAS, G., CHOOI, K. Y., VARDAKIS, J., MOREL, S., DIAGBOUGA, M. R., BIJLENGA, P., CUHLMANN, S., ET AL. Disturbed flow induces a sustained, stochastic $\text{nf-}\kappa\text{b}$ activation which may support intracranial aneurysm growth in vivo. *Scientific reports* 9, 1 (2019), 1–14.

- [15] BAEYENS, N., BANDYOPADHYAY, C., COON, B. G., YUN, S., AND SCHWARTZ, M. A. Endothelial fluid shear stress sensing in vascular health and disease. *The Journal of clinical investigation* 126, 3 (2016), 821–828.
- [16] BAILEY, K. A., MORENO, E., HAJ, F. G., SIMON, S. I., AND PASSERINI, A. G. Mechanoregulation of p38 activity enhances endoplasmic reticulum stress-mediated inflammation by arterial endothelium. *The FASEB Journal* 33, 11 (2019), 12888–12899.
- [17] BALAGURU, U. M., SUNDARESAN, L., MANIVANNAN, J., MAJUNATHAN, R., MANI, K., SWAMINATHAN, A., VENKATESAN, S., KASIVISWANATHAN, D., AND CHATTERJEE, S. Disturbed flow mediated modulation of shear forces on endothelial plane: A proposed model for studying endothelium around atherosclerotic plaques. *Scientific reports* 6 (2016), 27304.
- [18] BARATCHI, S., KHOSHMANESH, K., WOODMAN, O. L., POTOCHNIK, S., PETER, K., AND MCINTYRE, P. Molecular sensors of blood flow in endothelial cells. *Trends in molecular medicine* 23, 9 (2017), 850–868.
- [19] BARÁTH, K., CASSOT, F., RÜFENACHT, D. A., AND FASEL, J. H. Anatomically shaped internal carotid artery aneurysm in vitro model for flow analysis to evaluate stent effect. *American Journal of Neuroradiology* 25, 10 (2004), 1750–1759.

- [20] BAZILEVS, Y., HSU, M.-C., ZHANG, Y., WANG, W., KVAMSDAL, T., HENTSCHEL, S., AND ISAKSEN, J. Computational vascular fluid–structure interaction: methodology and application to cerebral aneurysms. *Biomechanics and modeling in mechanobiology* 9, 4 (2010), 481–498.
- [21] BENN, A., BREDOW, C., CASANOVA, I., VUKIČEVIĆ, S., AND KNAUS, P. Ve-cadherin facilitates bmp-induced endothelial cell permeability and signaling. *Journal of cell science* 129, 1 (2016), 206–218.
- [22] BENNDORF, G., WELLNHOFER, E., LANKSCH, W., AND FELIX, R. Intraaneurysmal flow: evaluation with doppler guidewires. *American journal of neuroradiology* 17, 7 (1996), 1333–1337.
- [23] BIASETTI, J., HUSSAIN, F., AND GASSER, T. C. Blood flow and coherent vortices in the normal and aneurysmatic aortas: a fluid dynamical approach to intra-luminal thrombus formation. *Journal of The Royal Society Interface* (2011), rsif20110041.
- [24] BIBLI, S.-I., HU, J., LEISEGANG, M. S., WITTIG, J., ZUKUNFT, S., KAPSAKALIDI, A., FISSLTHALER, B., TSILIMIGRAS, D., ZOGRAFOS, G., FILIS, K., ET AL. Shear stress regulates cystathionine γ lyase expression to preserve endothelial redox balance and reduce membrane lipid peroxidation. *Redox biology* 28 (2020), 101379.

- [25] BIJLENGA, P., GONDAR, R., SCHILLING, S., MOREL, S., HIRSCH, S., CUONY, J., CORNIOLA, M.-V., PERREN, F., RÜFENACHT, D., AND SCHALLER, K. Phases score for the management of intracranial aneurysm: a cross-sectional population-based retrospective study. *Stroke* 48, 8 (2017), 2105–2112.
- [26] BOGUNOVIC, N., MEEKEL, J. P., MICHA, D., BLANKENSTEIJN, J. D., HORDIJK, P. L., AND YEUNG, K. K. Impaired smooth muscle cell contractility as a novel concept of abdominal aortic aneurysm pathophysiology. *Scientific reports* 9, 1 (2019), 1–14.
- [27] BOUSSEL, L., RAYZ, V., MCCULLOCH, C., MARTIN, A., ACEVEDO-BOLTON, G., LAWTON, M., HIGASHIDA, R., SMITH, W. S., YOUNG, W. L., AND SALONER, D. Aneurysm growth occurs at region of low wall shear stress: patient-specific correlation of hemodynamics and growth in a longitudinal study. *Stroke* 39, 11 (2008), 2997–3002.
- [28] BRINJIKJI, W., KALLMES, D. F., LANZINO, G., AND CLOFT, H. Hospitalization costs for endovascular and surgical treatment of unruptured cerebral aneurysms in the united states are substantially higher than medicare payments. *American Journal of Neuroradiology* 33, 1 (2012), 49–51.
- [29] BRINJIKJI, W., ZHU, Y.-Q., LANZINO, G., CLOFT, H., MURAD, M., WANG, Z., AND KALLMES, D. Risk factors for growth of intracranial

- aneurysms: A systematic review and meta-analysis. *American Journal of Neuroradiology* (2015).
- [30] BRISMAN, J. L., SONG, J. K., AND NEWELL, D. W. Cerebral aneurysms. *New England journal of medicine* 355, 9 (2006), 928–939.
- [31] BYRNE, G., AND CEBRAL, J. Vortex dynamics in cerebral aneurysms. *arXiv preprint arXiv:1309.7875* (2013).
- [32] BYRNE, G., MUT, F., AND CEBRAL, J. Quantifying the large-scale hemodynamics of intracranial aneurysms. *American Journal of Neuroradiology* 35, 2 (2014), 333–338.
- [33] CAHILL, P. A., AND REDMOND, E. M. Vascular endothelium—gatekeeper of vessel health. *Atherosclerosis* 248 (2016), 97–109.
- [34] CAN, A., AND DU, R. Association of hemodynamic factors with intracranial aneurysm formation and rupture: systematic review and meta-analysis. *Neurosurgery* 78, 4 (2015), 510–520.
- [35] CANCEL, L. M., EBONG, E. E., MENSAH, S., HIRSCHBERG, C., AND TARBELL, J. M. Endothelial glycocalyx, apoptosis and inflammation in an atherosclerotic mouse model. *Atherosclerosis* 252 (2016), 136–146.
- [36] CAR, R., AND PARRINELLO, M. Unified Approach for Molecular Dynamics and Density-Functional Theory. *Physical Review Letters* 55 (1985), 2471.

- [37] CASTRO, M. A., OLIVARES, M. C. A., PUTMAN, C. M., AND CEBRAL, J. R. Wall motion and hemodynamics in intracranial aneurysms. In *Journal of Physics: Conference Series* (2013), vol. 477, IOP Publishing, p. 012004.
- [38] CEBRAL, J., DETMER, F., CHUNG, B., CHOQUE-VELASQUEZ, J., REZAI, B., LEHTO, H., TULAMO, R., HERNESNIEMI, J., NIEMELA, M., YU, A., WILLIAMSON, R., AZIZ, K., SAKUR, S., AMIN-HANJANI, S., CHARBEL, F., TOBE, Y., ROBERTSON, A., AND FRÖSEN, J. Local hemodynamic conditions associated with focal changes in the intracranial aneurysm wall. *American Journal of Neuroradiology* 40, 3 (2019), 510–516.
- [39] CEBRAL, J., OLLIKAINEN, E., CHUNG, B. J., MUT, F., SIPPOLA, V., JAHROMI, B. R., TULAMO, R., HERNESNIEMI, J., NIEMELÄ, M., ROBERTSON, A., AND FRÖSEN, J. Flow conditions in the intracranial aneurysm lumen are associated with inflammation and degenerative changes of the aneurysm wall. *American Journal of Neuroradiology* 38, 1 (2017), 119–126.
- [40] CEBRAL, J. R., MUT, F., WEIR, J., AND PUTMAN, C. M. Association of hemodynamic characteristics and cerebral aneurysm rupture. *American Journal of Neuroradiology* 32, 2 (2011), 264–270.
- [41] CECCHI, E., GIGLIOLI, C., VALENTE, S., LAZZERI, C., GENSINI, G. F., ABBATE, R., AND MANNINI, L. Role of hemodynamic shear stress in cardiovascular disease. *Atherosclerosis* 214, 2 (2011), 249–256.

- [42] CHA, S.-H., HWANG, Y., KIM, K.-N., AND JUN, H.-S. Palmitate induces nitric oxide production and inflammatory cytokine expression in zebrafish. *Fish & shellfish immunology* 79 (2018), 163–167.
- [43] CHALOUHI, N., ALI, M. S., JABBOUR, P. M., TJOUMAKARIS, S. I., GONZALEZ, L. F., ROSENWASSER, R. H., KOCH, W. J., AND DUMONT, A. S. Biology of intracranial aneurysms: role of inflammation. *Journal of Cerebral Blood Flow & Metabolism* 32, 9 (2012), 1659–1676.
- [44] CHALOUHI, N., HOH, B. L., AND HASAN, D. Review of cerebral aneurysm formation, growth, and rupture. *Stroke* 44, 12 (2013), 3613–3622.
- [45] CHALOUHI, N., ZANATY, M., WHITING, A., YANG, S., TJOUMAKARIS, S., HASAN, D., STARKE, R. M., HANN, S., HAMMER, C., KUNG, D., AND ET. AL. Safety and efficacy of the pipeline embolization device in 100 small intracranial aneurysms. *Journal of neurosurgery* 122, 6 (2015), 1498–1502.
- [46] CHATTERJEE, S. Endothelial mechanotransduction, redox signaling and the regulation of vascular inflammatory pathways. *Frontiers in physiology* 9 (2018), 524.
- [47] CHATTOPADHYAY, R., DYUKOVA, E., SINGH, N. K., OHBA, M., MOBLEY, J. A., AND RAO, G. N. Vascular endothelial tight junctions and barrier

- function are disrupted by 15 (s)-hydroxyeicosatetraenoic acid partly via protein kinase c-mediated zona occludens-1 phosphorylation at threonine 770/772. *Journal of Biological Chemistry* 289, 6 (2014), 3148–3163.
- [48] CHATZIZISIS, Y. S., COSKUN, A. U., JONAS, M., EDELMAN, E. R., FELDMAN, C. L., AND STONE, P. H. Role of endothelial shear stress in the natural history of coronary atherosclerosis and vascular remodeling: Molecular, cellular, and vascular behavior. *Journal of the American College of Cardiology* 49, 25 (2007), 2379 – 2393.
- [49] CHEN, C.-N., CHANG, S.-F., LEE, P.-L., CHANG, K., CHEN, L.-J., USAMI, S., CHIEN, S., AND CHIU, J.-J. Neutrophils, lymphocytes, and monocytes exhibit diverse behaviors in transendothelial and subendothelial migrations under coculture with smooth muscle cells in disturbed flow. *Blood* 107, 5 (2006), 1933–1942.
- [50] CHEN, Z., AND TZIMA, E. Pecam-1 is necessary for flow-induced vascular remodeling. *Arteriosclerosis, thrombosis, and vascular biology* 29, 7 (2009), 1067–1073.
- [51] CHIEN, S. Mechanotransduction and endothelial cell homeostasis: the wisdom of the cell. *American Journal of Physiology-Heart and Circulatory Physiology* 292, 3 (2007), H1209–H1224.

- [52] CHISTIYAKOV, D. A., OREKHOV, A. N., AND BOBRYSHV, Y. V. Effects of shear stress on endothelial cells: go with the flow. *Acta physiologica* 219, 2 (2017), 382–408.
- [53] CHIU, J.-J., AND CHIEN, S. Effects of disturbed flow on vascular endothelium: pathophysiological basis and clinical perspectives. *Physiological reviews* 91, 1 (2011), 327–387.
- [54] CLAESSEON-WELSH, L. Vascular permeabilitythe essentials. *Upsala journal of medical sciences* 120, 3 (2015), 135–143.
- [55] CODELIA, V. A., SUN, G., AND IRVINE, K. D. Regulation of yap by mechanical strain through jnk and hippo signaling. *Current biology* 24, 17 (2014), 2012–2017.
- [56] COOPER, S., EMMOTT, A., McDONALD, K. K., CAMPEAU, M.-A., AND LEASK, R. L. Increased mmp activity in curved geometries disrupts the endothelial cell glycocalyx creating a proinflammatory environment. *PloS one* 13, 8 (2018).
- [57] COOPER, S., McDONALD, K., BURKAT, D., AND LEASK, R. L. Stenosis hemodynamics disrupt the endothelial cell glycocalyx by mmp activity creating a proinflammatory environment. *Annals of biomedical engineering* 45, 9 (2017), 2234–2243.

- [58] CORNELISSEN, B., SCHNEIDERS, J., POTTERS, W., VAN DEN BERG, R., VELTHUIS, B., RINKEL, G., SLUMP, C., VANBAVEL, E., MAJOIE, C., AND MARQUERING, H. Hemodynamic differences in intracranial aneurysms before and after rupture. *American Journal of Neuroradiology* 36, 10 (2015), 1927–1933.
- [59] THE CPMD CONSORTIUM. *CPMD (v3.15.1): An ab initio Electronic Structure and Molecular Dynamics Program*, 2011.
- [60] DELLEY, B. An All-Electron Numerical Method for Solving the Local Density Functional for Polyatomic Molecules. *Journal of Chemical Physics* 92 (1990), 508.
- [61] DELLEY, B. Fast Calculation of Electrostatics in Crystals and Large Molecules. *Journal of Physical Chemistry* 100 (1996), 6107.
- [62] DEMARTINI, L. C., VIELMO, H. A., AND MÖLLER, S. Numeric and experimental analysis of the turbulent flow through a channel with baffle plates. *Journal of the Brazilian Society of Mechanical Sciences and Engineering* 26, 2 (2004), 153–159.
- [63] DEMPÈRE-MARCO, L., OUBEL, E., CASTRO, M., PUTMAN, C., FRANGI, A., AND CEBRAL, J. Cfd analysis incorporating the influence of wall motion:

application to intracranial aneurysms. In *International Conference on Medical Image Computing and Computer-Assisted Intervention* (2006), Springer, pp. 438–445.

- [64] DEPLANO, V., KNAPP, Y., BERTRAND, E., AND GAILLARD, E. Flow behaviour in an asymmetric compliant experimental model for abdominal aortic aneurysm. *Journal of biomechanics* 40, 11 (2007), 2406–2413.
- [65] DETMER, F. J., FAJARDO-JIMÉNEZ, D., MUT, F., JUCHLER, N., HIRSCH, S., PEREIRA, V. M., BIJLENGA, P., AND CEBRAL, J. R. External validation of cerebral aneurysm rupture probability model with data from two patient cohorts. *Acta neurochirurgica* 160, 12 (2018), 2425–2434.
- [66] DI ACHILLE, P., TELLIDES, G., AND HUMPHREY, J. Hemodynamics-driven deposition of intraluminal thrombus in abdominal aortic aneurysms. *International journal for numerical methods in biomedical engineering* 33, 5 (2017), e2828.
- [67] DIAGBOUGA, M. R., MOREL, S., BIJLENGA, P., AND KWAK, B. R. Role of hemodynamics in initiation/growth of intracranial aneurysms. *European journal of clinical investigation* 48, 9 (2018), e12992.
- [68] DODDASOMAYAJULA, R., CHUNG, B. J., MUT, F., JIMENEZ, C. M., HAMZEI-SICHANI, F., PUTMAN, C. M., AND CEBRAL, J. R. Hemodynamic characteristics of ruptured and unruptured multiple aneurysms at mirror and

- ipsilateral locations. *American Journal of Neuroradiology* 38, 12 (2017), 2301–2307.
- [69] DOLAN, J. M., KOLEGA, J., AND MENG, H. High wall shear stress and spatial gradients in vascular pathology: a review. *Annals of biomedical engineering* 41, 7 (2013), 1411–1427.
- [70] DOLAN, J. M., MENG, H., SINGH, S., PALUCH, R., AND KOLEGA, J. High fluid shear stress and spatial shear stress gradients affect endothelial proliferation, survival, and alignment. *Annals of biomedical engineering* 39, 6 (2011), 1620–1631.
- [71] DONGARRA, J. J. LINPACK Working Note 3: Fortran BLAS Timing. *Argonne National Laboratory Report, ANL-80-24* (1980).
- [72] DONGARRA, J. J., BUNCH, J., MOLER, C., AND STEWART, G. W. *LINPACK User's Guide*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 1979.
- [73] DONGARRA, J. J., CROZ, J. D., HAMMARLING, S., AND DUFF, I. S. A Set of Level 3 Basic Linear Algebra Subprograms. *Association for Computing Machinery Transactions on Mathematical Software* 16 (1990), 1.
- [74] DONGARRA, J. J., CROZ, J. D., HAMMARLING, S., AND HANSON, R. An Extended Set of FORTRAN Basic Linear Algebra Subprograms. *Association*

- for *Computing Machinery Transactions on Mathematical Software* 14 (1988), 1.
- [75] DORLAND, Y. L., AND HUVENEERS, S. Cell-cell junctional mechanotransduction in endothelial remodeling. *Cellular and Molecular Life Sciences* 74, 2 (2017), 279–292.
- [76] DOVESI, R., ORLANDO, R., CIVALLERI, B., ROETTI, C., SAUNDERS, V. R., AND ZICOVICH-WILSON, C. M. CRYSTAL: A Computational Tool for the Ab Initio Study of the Electronic Properties of Crystals. *Zeitschrift für Kristallographie* 220 (2005), 571.
- [77] DOVESI, R., SAUNDERS, V. R., ROETTI, C., ORLANDO, R., ZICOVICH-WILSON, C. M., PASCALE, F., CIVALLERI, B., DOLL, K., HARRISON, N. M., BUSH, I. J., D’ARCO, P., AND LLUNELL, M. *CRYSTAL 09 User’s Manual*. University of Torino, Italy, 2009.
- [78] DUAN, Z., LI, Y., GUAN, S., MA, C., HAN, Y., REN, X., WE, L., LI, W., LO, J., AND YANG, Z. Morphological parameters and anatomical locations associated with rupture status of small intracranial aneurysms. *Scientific reports* 8 (2018).
- [79] DUCA, L., BLAISE, S., ROMIER, B., LAFFARGUE, M., GAYRAL, S., EL BTAOURI, H., KAWECKI, C., GUILLOT, A., MARTINY, L., DEBELLE,

- L., ET AL. Matrix ageing and vascular impacts: focus on elastin fragmentation. *Cardiovascular research* 110, 3 (2016), 298–308.
- [80] EFRON, B., HASTIE, T., JOHNSTONE, I., TIBSHIRANI, R., AND ET. AL. Least angle regression. *The Annals of statistics* 32, 2 (2004), 407–499.
- [81] ELAD, D., AND EINAV, S. Physical and flow properties of blood. *Standard handbook of biomedical engineering and design* (2004), 3–1.
- [82] ENZMANN, D. R., ROSS, M. R., MARKS, M. P., AND PELC, N. J. Blood flow in major cerebral arteries measured by phase-contrast cine mr. *American journal of neuroradiology* 15, 1 (1994), 123–129.
- [83] ETMINAN, N., BROWN, R. D., BESEOGLU, K., JUVELA, S., RAYMOND, J., MORITA, A., TORNER, J. C., DERDEYN, C. P., RAABE, A., MOCCO, J., AND ET. AL. The unruptured intracranial aneurysm treatment score a multidisciplinary consensus. *Neurology* 85, 10 (2015), 881–889.
- [84] FALGOUT, R. D., AND YANG, U. M. HYPRE: A Library of High Performance Preconditioners. In *Proceedings of the International Conference on Computational Science - Part III* (London, UK, 2002), ICCS '02, Springer-Verlag, p. 632.
- [85] FELICIANI, G., POTTERS, W. V., VAN OOIJ, P., SCHNEIDERS, J. J., NEDERVEEN, A. J., VAN BAVEL, E., MAJOIE, C. B., AND MARQUERING, H. A. Multiscale 3-d+ t intracranial aneurysmal flow vortex detection. *IEEE Trans. Biomed. Engineering* 62, 5 (2015), 1355–1362.

- [86] FINCH, H. A comparison of methods for group prediction with high dimensional data. *Journal of Modern Applied Statistical Methods* 13, 2 (2014), 5.
- [87] FORD, M., HOI, Y., PICCINELLI, M., ANTIGA, L., AND STEINMAN, D. An objective approach to digital removal of saccular aneurysms: technique and applications. *The British Journal of Radiology* 82, special_issue_1 (2009), S55–S61.
- [88] FORD, M. D., ALPERIN, N., LEE, S. H., HOLDSWORTH, D. W., AND STEINMAN, D. A. Characterization of volumetric flow rate waveforms in the normal internal carotid and vertebral arteries. *Physiological measurement* 26, 4 (2005), 477.
- [89] FORD, M. D., NIKOLOV, H. N., MILNER, J. S., LOWNIE, S. P., DEMONT, E. M., KALATA, W., LOTH, F., HOLDSWORTH, D. W., AND STEINMAN, D. A. Piv-measured versus cfd-predicted flow dynamics in anatomically realistic cerebral aneurysm models. *Journal of biomechanical engineering* 130, 2 (2008), 021015.
- [90] FRIGO, M., AND JOHNSON, S. G. The Design and Implementation of FFTW3. In *Proceedings of the IEEE* (2005), vol. 93, p. 216.
- [91] FUNG, J. C. H. Residence time of inertial particles in a vortex. *Journal of Geophysical Research: Oceans* 105, C6 (2000), 14261–14272.

- [92] GABRIEL, S. A., DING, Y., AND FENG, Y. Quantifying the influence of oscillatory flow disturbances on blood flow. *Journal of Theoretical Biology* 430 (2017), 195 – 206.
- [93] GALE, J. D. Empirical Potential Derivation for Ionic Materials. *Philosophical Magazine B* 73 (1996), 3.
- [94] GALE, J. D. GULP - A Computer Program for the Symmetry Adapted Simulation of Solids. *Journal of Chemical Society, Faraday Transactions* 93 (1997), 629.
- [95] GHIM, M., ALPRESA, P., YANG, S.-W., BRAAKMAN, S. T., GRAY, S. G., SHERWIN, S. J., VAN REEUWIJK, M., AND WEINBERG, P. D. Visualization of three pathways for macromolecule transport across cultured endothelium and their modification by flow. *American Journal of Physiology-Heart and Circulatory Physiology* 313, 5 (2017), H959–H973.
- [96] GO, Y.-M., SON, D. J., PARK, H., ORR, M., HAO, L., TAKABE, W., KUMAR, S., KANG, D. W., KIM, C. W., JO, H., ET AL. Disturbed flow enhances inflammatory signaling and atherogenesis by increasing thioredoxin-1 level in endothelial cell nuclei. *PloS one* 9, 9 (2014).
- [97] GREVING, J. P., WERMER, M. J., JR, M. J. B., MORIT, A., JUVELA, S., YONEKURA, M., ISHIBASHI, T., TORNER, J. C., NAKAYAMA, T., RINKEL, G. J., AND ET. AL. Development of the phases score for prediction of risk of

- rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *The Lancet Neurology* 13, 1 (2014), 59–66.
- [98] GSAM KIM, Y., PAR, Y., AND LIM, S. 3d simulations of blood flow dynamics in compliant vessels: normal, aneurysmal, and stenotic arteries. *Communications in Computational Physics* 19, 5 (2016), 1167–1190.
- [99] GWILLIAM, M. N., HOGGARD, N., CAPENER, D., SINGH, P., MARZO, A., VERMA, P. K., AND WILKINSON, I. D. Mr derived volumetric flow rate waveforms at locations within the common carotid, internal carotid, and basilar arteries. *Journal of Cerebral Blood Flow & Metabolism* 29, 12 (2009), 1975–1982.
- [100] HACKENBERG, K. A., HÄNGGI, D., AND ETMINAN, N. Unruptured intracranial aneurysms: Contemporary data and management. *Stroke* 49, 9 (2018), 2268–2275.
- [101] HAIMES, R., AND KENWRIGHT, D. On the velocity gradient tensor and fluid feature extraction. In *14th Computational Fluid Dynamics Conference* (1999), p. 3288.
- [102] HANCZAR, B., HU, J., SIM, C., WEINSTEIN, J., BITTNER, M., AND RDOUGHERTY, E. Small-sample precision of roc-related estimates. *Bioinformatics* 26, 6 (2010), 822–830.

- [103] HANLEY, J. A., AND MCNEIL, B. J. The meaning and use of the area under a receiver operating characteristic (roc) curve. *Radiology* 143, 1 (1982), 29–36.
- [104] HARDING, I. C., MITRA, R., MENSAH, S. A., HERMAN, I. M., AND EBONG, E. E. Pro-atherosclerotic disturbed flow disrupts caveolin-1 expression, localization, and function via glycocalyx degradation. *Journal of translational medicine* 16, 1 (2018), 1–20.
- [105] HARRELL, F. E., LEE, K. L., AND MARK, D. B. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Statistics in medicine* 15, 4 (1996), 361–387.
- [106] HASAN, D. M., NADAREYSHVILI, A. I., HOPPE, A. L., MAHANE, K. B., KUNG, D. K., AND RAGHAVAN, M. L. Cerebral aneurysm sac growth as the etiology of recurrence after successful coil embolization. *Stroke* 43, 3 (2012), 866–868.
- [107] HASHIMOTO, T., MENG, H., AND YOUNG, W. Intracranial aneurysms: links among inflammation, hemodynamics and vascular remodeling. *Neurol Res* 28 (2006), 372–380.
- [108] HELMKE, B. P. Molecular control of cytoskeletal mechanics by hemodynamic forces. *Physiology* 20, 1 (2005), 43–53.
- [109] HEO, K.-S., LE, N.-T., CUSHMAN, H. J., GIANCORSIO, C. J., CHANG, E., WOO, C.-H., SULLIVAN, M. A., TAUNTON, J., YEH, E. T., FUJIWARA,

- K., ET AL. Disturbed flow-activated p90rsk kinase accelerates atherosclerosis by inhibiting senp2 function. *The Journal of clinical investigation* 125, 3 (2015), 1299–1310.
- [110] HESS, B., KUTZNER, C., VAN DER SPOEL, D., AND LINDAHL, E. GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *Journal of Chemical Theory and Computation* 4 (2008), 235.
- [111] HOLDSWORTH, D., NORLEY, C., FRAYNE, R., STEINMAN, D., AND RUTT, B. Characterization of common carotid artery blood-flow waveforms in normal human subjects. *Physiological measurement* 20, 3 (1999), 219.
- [112] HOPPE, A. L., RAGHAVA, M. L., AND HASAN, D. M. Comparison of the association of sac growth and coil compaction with recurrence in coil embolized cerebral aneurysms. *PloS one* 10, 4 (2015), e0123017.
- [113] [HTTP://WIKIPEDIA.ORG/](http://WIKIPEDIA.ORG/). *Wikipedia*. Wikipedia, The Internet, 2012.
- [114] HUGHES, J. D., BOND, K. M., MEKARY, R. A., DEWAN, M. C., RATTANI, A., BATICULON, R., KATO, Y., AZEVEDO-FILHO, H., MORCOS, J. J., AND PARK, K. B. Estimating the global incidence of aneurysmal subarachnoid hemorrhage: a systematic review for central nervous system vascular lesions and meta-analysis of ruptured aneurysms. *World neurosurgery* 115 (2018), 430–447.

- [115] HUMPHREY, W., DALKE, A., AND SCHULTEN, K. VMD - Visual Molecular Dynamics. *Journal of Molecular Graphics* 14 (1996), 33.
- [116] HUNT, J. C., WRAY, A. A., AND MOIN, P. Eddies, streams, and convergence zones in turbulent flows.
- [117] HUO, Y., CHOY, J. S., SVENDSEN, M., SINHA, A. K., AND KASSAB, G. S. Effects of vessel compliance on flow pattern in porcine epicardial right coronary arterial tree. *Journal of biomechanics* 42, 5 (2009), 594–602.
- [118] ILDIKO, F. E., AND FRIEDMAN, J. H. A statistical view of some chemometrics regression tools. *Technometrics* 35, 2 (1993), 109–135.
- [119] INVESTIGATORS, U. J. The natural course of unruptured cerebral aneurysms in a japanese cohort. *New England Journal of Medicine* 366, 26 (2012), 2474–2482.
- [120] JEONG, J., AND HUSSAIN, F. On the identification of a vortex. *Journal of fluid mechanics* 285 (1995), 69–94.
- [121] JIANG, J., AND STROTHER, C. M. Interactive decomposition and mapping of saccular cerebral aneurysms using harmonic functions: its first application with patient-specific computational fluid dynamics (cfd) simulations. *IEEE transactions on medical imaging* 32, 2 (2013), 153–164.
- [122] JIANG, M., MACHIRAJU, R., AND THOMPSON, D. Detection and visualization of vortices. *The visualization handbook* 295 (2005).

- [123] JIANG, P., LIU, Q., WU, J., CHEN, X., LI, M., LI, Z., YANG, S., GUO, R., GAO, B., CAO, Y., ET AL. A novel scoring system for rupture risk stratification of intracranial aneurysms: a hemodynamic and morphological study. *Frontiers in neuroscience* 12 (2018), 596.
- [124] JING, L., FAN, J., WANG, Y., LI, H., WANG, S., YANG, X., AND ZHANG, Y. Morphologic and hemodynamic analysis in the patients with multiple intracranial aneurysms: Ruptured versus unruptured. *PLOS ONE* 10, 7 (07 2015), 1–12.
- [125] JR, M. A. G., AND GARCÍA-CARDEÑA, G. Endothelial cell dysfunction and the pathobiology of atherosclerosis. *Circulation research* 118, 4 (2016), 620–636.
- [126] JR, T. R. F., BENITEZ, R., VEZNEDAROGLU, E., SHARAN, A., MITCHELL, W., SILVA, M., AND ROSENWASSER, R. H. A review of size and location of ruptured intracranial aneurysms. *Neurosurgery* 49, 6 (2001), 1322–1326.
- [127] JUVELA, S. Risk factors for multiple intracranial aneurysms. *Stroke* 31, 2 (2000), 392–397.
- [128] JUVELA, S., POUSSA, K., LEHTO, H., AND PORRAS, M. Natural history of unruptured intracranial aneurysms: a long-term follow-up study. *Stroke* 44, 9 (2013), 2414–2421.

- [129] KARINO, T., AND GOLDSMITH, H. Flow behaviour of blood cells and rigid spheres in an annular vortex. *Phil. Trans. R. Soc. Lond. B* 279, 967 (1977), 413–445.
- [130] KASHIWAZAKI, D., AND KURODA, S. Size ratio can highly predict rupture risk in intracranial small (≤ 5 mm) aneurysms. *Stroke* 44, 8 (2013), 2169–2173.
- [131] KAUFMANN, B. A., SANDERS, J. M., DAVIS, C., XIE, A., ALDRED, P., SAREMBOCK, I. J., AND LIDNER, J. R. Molecular imaging of inflammation in atherosclerosis with targeted ultrasound detection of vascular cell adhesion molecule-1. *Circulation* 116, 3 (2007), 276–284.
- [132] KELSEY, L. J., POWELL, J. T., NORMAN, P. E., MILLER, K., AND DOYLE, B. J. A comparison of hemodynamic metrics and intraluminal thrombus burden in a common iliac artery aneurysm. *International journal for numerical methods in biomedical engineering* 33, 5 (2017), e2821.
- [133] KHAN, M., VALEN-SENDSTAD, K., AND STEINMAN, D. Narrowing the expertise gap for predicting intracranial aneurysm hemodynamics: impact of solver numerics versus mesh and time-step resolution. *American Journal of Neuroradiology* (2015).
- [134] KIM, M.-C., NAM, J. H., AND LEE, C.-S. Near-wall deposition probability of blood elements as a new hemodynamic wall parameter. *Annals of Biomedical Engineering* 34, 6 (Jun 2006), 958–970.

- [135] KLAUS, V., TANIOS-SCHMIES, F., REEPS, C., TRENNER, M., MATEVOSIAN, E., ECKSTEIN, H.-H., AND PELISEK, J. Association of matrix metalloproteinase levels with collagen degradation in the context of abdominal aortic aneurysm. *European Journal of Vascular and Endovascular Surgery* 53, 4 (2017), 549–558.
- [136] KOCUR, D., PRZYBYŁKO, N., NIEDBAŁA, M., AND RUDNIK, A. Alternative definitions of cerebral aneurysm morphologic parameters have an impact on rupture risk determination. *World neurosurgery* 126 (2019), e157–e164.
- [137] KÖHLER, B., GASTEIGER, R., PREIM, U., THEISEL, H., GUTBERLET, M., AND PREIM, B. Semi-automatic vortex extraction in 4d pc-mri cardiac blood flow data using line predicates. *IEEE Transactions on Visualization and Computer Graphics* 19, 12 (2013), 2773–2782.
- [138] KOLÁŘ, V. Vortex identification: New requirements and limitations. *International journal of heat and fluid flow* 28, 4 (2007), 638–652.
- [139] KOLKAJI, A. *Molecular Graphics Modelling* 17 (1999), 176.
- [140] KOMOTAR, R. J., MOCCO, J., AND SOLOMON, R. A. Guidelines for the surgical treatment of unruptured intracranial aneurysms: the first annual j. lawrence pool memorial research symposiumcontroversies in the management of cerebral aneurysms. *Neurosurgery* 62, 1 (2008), 183–194.

- [141] KORJA, M., LEHTO, H., AND JUVELA, S. Lifelong rupture risk of intracranial aneurysms depends on risk factors. *Stroke* 45, 7 (2014), 1958–1963.
- [142] KOSIERKIEWICZ, T., FACTOR, S., AND DICKSON, D. Immunocytochemical studies of atherosclerotic lesions of cerebral berry aneurysms. *J Neuropathol Exp Neurol* 53, 4 (1994), 399–406.
- [143] KOTOWSKI, M., NAGGARA, O., DARSAUT, T. E., NOLET, S., GEVRY, G., KOUZNETSOV, E., AND RAYMOND, J. Safety and occlusion rates of surgical treatment of unruptured intracranial aneurysms: a systematic review and meta-analysis of the literature from 1990 to 2011. *Journal of Neurology, Neurosurgery & Psychiatry* 84, 1 (2013), 42–48.
- [144] KRESSE, G., AND HAFNER, J. Ab Initio Molecular Dynamics for Liquid Metals. *Physical Review B* 47 (1993), 558.
- [145] KRESSE, G., AND HAFNER, J. Ab Initio Molecular-Dynamics Simulation of the Liquid-Metal-Amorphous-Semiconductor Transition in Germanium. *Physical Review B* 49 (1994), 14251.
- [146] KU, D., GIDDEN, D., PHILLIPS, D., AND JR, D. S. Hemodynamics of the normal human carotid bifurcation: in vitro and in vivo studies. *Ultrasound in medicine & biology* 11, 1 (1985), 13–26.
- [147] KU, D., GIDDENS, D. P., ZARINS, C. K., AND GLAGOV, S. Pulsatile flow and atherosclerosis in the human carotid bifurcation. positive correlation

- between plaque location and low oscillating shear stress. *Arteriosclerosis: An Official Journal of the American Heart Association, Inc.* 5, 3 (1985), 293–302.
- [148] KULCSÁR, Z., UGRON, A., BERENTEI, Z., PAÁL, G., SZIKORA, I., AND ET AL. Hemodynamics of cerebral aneurysm initiation: the role of wall shear stress and spatial wall shear stress gradient. *American Journal of neuroradiology* (2011).
- [149] LAY, A. J., COLEMAN, P. R., FORMAZ-PRESTON, A., TING, K. K., ROEDIGER, B., WENINGER, W., SCHWARTZ, M. A., VADAS, M. A., AND GAMBLE, J. R. Arhgap18: A flow-responsive gene that regulates endothelial cell alignment and protects against atherosclerosis. *Journal of the American Heart Association* 8, 2 (2019), e010057.
- [150] LEACH, J. R., KAO, E., ZHU, C., SALONER, D., AND HOPE, M. D. On the relative impact of intraluminal thrombus heterogeneity on abdominal aortic aneurysm mechanics. *Journal of biomechanical engineering* 141, 11 (2019).
- [151] LEE, G., EOM, K., LEE, C., KIM, D., AND KANG, S. Rupture of very small intracranial aneurysms: Incidence and clinical characteristics. *J Cerebrovasc Endovasc Neurosurg* 17(3) (2015), 217–222.
- [152] LI, T., WANG, H., LI, X., GE, H., SUN, H., AND MA, D. Predictive significance of vegfa variations in intracranial aneurysm. *Int J Clin Exp Med* 10, 9 (2017), 13802–13807.

- [153] LIU, J., JING, L., WANG, C., ZHANG, Y., AND YANG, X. Recanalization, regrowth, and delayed rupture of a previously coiled unruptured anterior communicating artery aneurysm: a longitudinal hemodynamic analysis. *World neurosurgery* 89 (2016), 726–e5.
- [154] LIU, P., SONG, Y., ZHOU, Y., LIU, Y., QIU, T., AN, Q., SONG, J., LI, P., SHI, Y., LI, S., ET AL. Cyclic mechanical stretch induced smooth muscle cell changes in cerebral aneurysm progress by reducing collagen type iv and collagen type vi levels. *Cellular Physiology and Biochemistry* 45, 3 (2018), 1051–1060.
- [155] LIU, X., SUN, A., FAN, Y., AND DENG, X. Physiological significance of helical flow in the arterial system and its potential clinical applications. *Annals of Biomedical Engineering* 43, 1 (Jan 2015), 3–15.
- [156] LONGO, M., GRANATA, F., RACCHIUSA, S., MORMINA, E., GRASSO, G., LONGO, G. M., GARUFI, G., SALPIETRO, F. M., AND ALAFACI, C. Role of hemodynamic forces in unruptured intracranial aneurysms: An overview of a complex scenario. *World Neurosurgery* 105 (2017), 632 – 642.
- [157] LORENSEN, W. E., AND CLINE, H. E. Marching cubes: A high resolution 3d surface construction algorithm. In *ACM siggraph computer graphics* (1987), vol. 21, ACM, pp. 163–169.

- [158] MA, D., TREMMEL, M., PALUCH, R. A., LEVY, E. L. I., MENG, H., AND MOCCO, J. Size ratio for clinical assessment of intracranial aneurysm rupture risk. *Neurological research* 32, 5 (2010), 482–486.
- [159] MA, J., WANG, C., SHENE, C.-K., AND JIANG, J. A graph-based interface for visualanalytics of 3d streamlines and pathlines. *IEEE transactions on visualization and computer graphics* 20, 8 (2014), 1127–1140.
- [160] MANNINO, R. G., MYERS, D. R., AHN, B., WANG, Y., ROLLINS, M., GOLE, H., LIN, A. S., GULDBERG, R. E., GIDDENS, D. P., TIMMINS, L. H., AND ET. AL. Do-it-yourself in vitro vasculature that recapitulates in vivo geometries for investigating endothelial-blood cell interactions. *Scientific reports* 5 (2015), 12401.
- [161] MAO, X., XIE, L., GREENBERG, R. B., GREENBERG, J. B., PENG, B., MIELING, I., JIN, K., AND GREENBERG, D. A. Flow-induced regulation of brain endothelial cells in vitro. *Vascular pharmacology* 62, 2 (2014), 82–87.
- [162] MARKL, M., WEGENT, F., ZECH, T., BAUER, S., STRECKER, C., SCHUMACHER, M., WEILLER, C., HENNIG, J., AND HARLOFF, A. In vivo wall shear stress distribution in the carotid artery: effect of bifurcation geometry, internal carotid artery stenosis, and recanalization therapy. *Circulation: Cardiovascular Imaging* 3, 6 (2010), 647–655.

- [163] MARTIN, D., LI, Y., YANG, J., WANG, G., MARGARITI, A., JIANG, Z., YU, H., ZAMPETAKI, A., HU, Y., XU, Q., ET AL. Unspliced x-box-binding protein 1 (xbp1) protects endothelial cells from oxidative stress through interaction with histone deacetylase 3. *Journal of Biological Chemistry* 289, 44 (2014), 30625–30634.
- [164] MASCITELLI, J. R., OERMANN, E. K., LEACY, R. A. D., MOYLE, H., MOCCO, J., AND PATEL, A. B. Predictors of treatment failure following coil embolization of intracranial aneurysms. *Journal of Clinical Neuroscience* 22, 8 (2015), 1275–1281.
- [165] McDONALD, K. K., COOPER, S., DANIELZAK, L., AND LEASK, R. L. Glycocalyx degradation induces a proinflammatory phenotype and increased leukocyte adhesion in cultured endothelial cells under flow. *PloS one* 11, 12 (2016).
- [166] McDOWELL, M. M., ZHAO, Y., KELLNER, C. P., BARTON, S. M., SUSSMAN, E., CLAASSEN, J., DUCRUET, A. F., AND CONNOLLY, E. S. Demographic and clinical predictors of multiple intracranial aneurysms in patients with subarachnoid hemorrhage. *Journal of Neurosurgery JNS* 128, 4 (2018), 961 – 968.
- [167] McDOWELL, M. M., ZHAO, Y., KELLNER, C. P., BARTON, S. M., SUSSMAN, E., CLAASSEN, J., DUCRUET, A. F., AND CONNOLLY, E. S. Demographic and clinical predictors of multiple intracranial aneurysms in patients

- with subarachnoid hemorrhage. *Journal of neurosurgery* 128, 4 (2018), 961–968.
- [168] MECKEL, S., STALDER, A. F., SANTINI, F., RADÜ, E.-W., RÜFENACHT, D. A., MARKL, M., AND WETZEL, S. G. In vivo visualization and analysis of 3-d hemodynamics in cerebral aneurysms with flow-sensitized 4-d mr imaging at 3 t. *Neuroradiology* 50, 6 (2008), 473–484.
- [169] MENG, H., TUTINO, V. M., XIANG, J., AND SIDDIQUI, A. High wss or low wss? complex interactions of hemodynamics with intracranial aneurysm initiation, growth, and rupture: Toward a unifying hypothesis. *American Journal of Neuroradiology* 35, 7 (2014), 1254–1262.
- [170] MICHEL, J.-B., JONDEAU, G., AND MILEWICZ, D. M. From genetics to response to injury: vascular smooth muscle cells in aneurysms and dissections of the ascending aorta. *Cardiovascular research* 114, 4 (2018), 578–589.
- [171] MITRA, R., QIAO, J., MADHAVAN, S., ONEIL, G. L., RITCHIE, B., KULKARNI, P., SRIDHAR, S., VAN DE VEN, A. L., KEMMERLING, E. M. C., FERRIS, C., ET AL. The comparative effects of high fat diet or disturbed blood flow on glycocalyx integrity and vascular inflammation. *Translational medicine communications* 3, 1 (2018), 1–15.

- [172] MIURA, Y., ISHIDA, F., UMEDA, Y., TANEMURA, H., SUZUKI, H., MATSUSHIMA, S., SHIMOSAKA, S., AND TAKI, W. Low wall shear stress is independently associated with the rupture status of middle cerebral artery aneurysms. *Stroke* 44, 2 (2013), 519–521.
- [173] MOCCO, J., BROWN JR, R. D., TORNER, J. C., CAPUANO, A. W., FARGEN, K. M., RAGHAVAN, M. L., PIEPGRAS, D. G., MEISSNER, I., III, J. H., AND OF UNRUPTURED INTRACRANIAL ANEURYSMS INVESTIGATORS, I. S. Aneurysm morphology and prediction of rupture: an international study of unruptured intracranial aneurysms analysis. *Neurosurgery* 82, 4 (2018), 491–496.
- [174] MOHAN, D., MUNTEANU, V., COMAN, T., AND CIUREA, A. Genetic factors involves in intracranial aneurysms—actualities. *Journal of medicine and life* 8, 3 (2015), 336.
- [175] MOLYNEUX, A., KERR, R., GROUP, I. S. A. T. I. C., ET AL. International subarachnoid aneurysm trial (isat) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomized trial. *Journal of stroke and cerebrovascular diseases* 11, 6 (2002), 304–314.
- [176] MORIGUCHI, T., AND SUMPIO, B. E. Pecam-1 phosphorylation and tissue factor expression in huvecs exposed to uniform and disturbed pulsatile flow and chemical stimuli. *Journal of vascular surgery* 61, 2 (2015), 481–488.

- [177] MUNDI, S., MASSARO, M., SCODITTI, E., CARLUCCIO, M. A., VAN HINSBERGH, V. W., IRUELA-ARISPE, M. L., AND DE CATERINA, R. Endothelial permeability, ldl deposition, and cardiovascular risk factorsa review. *Cardiovascular research* 114, 1 (2018), 35–52.
- [178] MUNDI, S., MASSARO, M., SCODITTI, E., CARLUCCIO, M. A., VAN HINSBERGH, V. W. M., IRUELA-ARISPE, M. L., AND DE CATERINA, R. Endothelial permeability, LDL deposition, and cardiovascular risk factorsa review. *Cardiovascular Research* 114, 1 (12 2017), 35–52.
- [179] NOBARI, S., MONGRAIN, R., LEASK, R., AND CARTIER, R. The effect of aortic wall and aortic leaflet stiffening on coronary hemodynamic: a fluid–structure interaction study. *Medical & biological engineering & computing* 51, 8 (2013), 923–936.
- [180] NOWICKI, K. W., HOSAKA, K., HE, Y., MCFETRIDGE, P. S., SCOTT, E. W., AND HOH, B. L. Novel high-throughput in vitro model for identifying hemodynamic-induced inflammatory mediators of cerebral aneurysm formation. *Hypertension* 64, 6 (2014), 1306–1313.
- [181] OELTZE-JAFRA, S., CEBRAL, J. R., JANIG, G., AND PREIM, B. Cluster analysis of vortical flow in simulations of cerebral aneurysm hemodynamics. *IEEE transactions on visualization and computer graphics* 22, 1 (2016), 757–766.

- [182] OLLIKAINEN, E., TULAMO, R., LEHTI, S., LEE-RUECKERT, M., HERNES-
NIEMI, J., NIEMEL, M., YL-HERTTUALA, S., KOVANEN, P. T., AND FRSEN,
J. Smooth muscle cell foam cell formation, apolipoproteins, and abca1 in in-
tracranial aneurysms: Implications for lipid accumulation as a promoter of
aneurysm wall rupture. *Journal of Neuropathology and Experimental Neurology*
75, 7 (2016), 689–699.

- [183] ORDEJÓ, P., DRABOLD, D. A., GRUMBACH, M. P., AND MARTIN,
R. M. Unconstrained Minimization Approach for Electronic Computations
That Scales Linearly with System Size. *Physical Review B* 48 (1993), 14646.

- [184] OTANI, T., NAKAMURA, M., FUJINAKA, T., HIRATA, M., KURODA, J.,
SHIBANO, K., AND WADA, S. Computational fluid dynamics of blood flow
in coil-embolized aneurysms: effect of packing density on flow stagnation in an
idealized geometry. *Medical & biological engineering & computing* 51, 8 (2013),
901–910.

- [185] OUBEL, E., CRAENE, M. D., PUTMAN, C. M., CEBRAL, J. R., AND
FRANGI, A. F. Analysis of intracranial aneurysm wall motion and its effects
on hemodynamic patterns. In *Medical Imaging 2007: Physiology, Function,
and Structure from Medical Images* (2007), vol. 6511, International Society for
Optics and Photonics, p. 65112A.

- [186] PAN, L., HONG, Z., YU, L., GAO, Y., ZHANG, R., FENG, H., SU, L., AND WANG, G. Shear stress induces human aortic endothelial cell apoptosis via interleukin-1 receptor-associated kinase 2-induced endoplasmic reticulum stress. *Molecular medicine reports* 16, 5 (2017), 7205–7212.
- [187] PAPAIOANNOU, T. G., AND STEFANADIS, C. Vascular wall shear stress: basic principles and methods. *Hellenic J Cardiol* 46, 1 (2005), 9–15.
- [188] PENN, D. L., WITTE, S. R., KOMOTAR, R. J., AND JR, E. S. C. The role of vascular remodeling and inflammation in the pathogenesis of intracranial aneurysms. *Journal of clinical neuroscience* 21, 1 (2014), 28–32.
- [189] PICCINELLI, M., STEINMAN, D. A., HOI, Y., TONG, F., VENEZIANI, A., AND ANTIGA, L. Automatic neck plane detection and 3d geometric characterization of aneurysmal sacs. *Annals of Biomedical Engineering* 40, 10 (2012), 2188–2211.
- [190] PLIMPTON, S. J. Fast Parallel Algorithms for Short-Range Molecular Dynamics. *Journal of Computational Physics* 117 (1995), 1.
- [191] POETHKE, J., SPULER, A., PETZ, C., HEGE, H.-C., GOUBERGRITS, L., AFFELD, K., AND KERTZSCHER, U. Cerebral aneurysm hemodynamics and a length of parent vessel. In *World Congress on Medical Physics and Biomedical Engineering, September 7-12, 2009, Munich, Germany* (2009), Springer, pp. 1608–1611.

- [192] POTTERS, W. V., MARQUERING, H. A., VANBAVEL, E., AND NEDERVEEN, A. J. Measuring wall shear stress using velocity-encoded mri. *Current Cardiovascular Imaging Reports* 7, 4 (2014), 9257.
- [193] QIAN, Z., KANG, H., TANG, K., JIANG, C., WU, Z., LI, Y., AND LIU, A. Assessment of risk of aneurysmal rupture in patients with normotensives, controlled hypertension, and uncontrolled hypertension. *Journal of Stroke and Cerebrovascular Diseases* 25, 7 (2016), 1746–1752.
- [194] QIN, H., YANG, Q., ZHUANG, Q., LONG, J., YANG, F., AND ZHANG, H. Morphological and hemodynamic parameters for middle cerebral artery bifurcation aneurysm rupture risk assessment. *Journal of Korean Neurosurgical Society* 60, 5 (2017), 504.
- [195] QIU, J., ZHENG, Y., HU, J., LIAO, D., GREGERSEN, H., DENG, X., FAN, Y., AND WANG, G. Biomechanical regulation of vascular smooth muscle cell functions: from in vitro to in vivo understanding. *Journal of The Royal Society Interface* 11, 90 (2014), 20130852.
- [196] R DEVELOPMENT CORE TEAM. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2011.
- [197] RAMELLA, M., BERNARDI, P., FUSARO, L., MANFREDI, M., CASELLA, F., PORTA, C. M., NICOLAI, L., GALEAZZI, E., BOLDORINI, R., SETTEMBRINI,

- A. M., ET AL. Relevance of inflammation and matrix remodeling in abdominal aortic aneurysm (aaa) and popliteal artery aneurysm (paa) progression. *American journal of translational research* 10, 10 (2018), 3265.
- [198] REVILLA-PACHECO, F., ESCALANTE-SEYFFERT, M. C., HERRADA-PINEDA, T., MANRIQUE-GUZMAN, S., PEREZ-ZUNIGA, I., RANGEL-SUAREZ, S., RUBALCAVA-ORTEGA, J., AND LOYO-VARELA, M. Prevalence of incidental clinoid segment saccular aneurysms. *World neurosurgery* 115 (2018), e244–e251.
- [199] RINNE, J., HERNESNIEMI, J., PURANEN, M., AND SAARI, T. Multiple intracranial aneurysms in a defined population: prospective angiographic and clinical study. *Neurosurgery* 35, 5 (1994), 803–808.
- [200] ROCHA, A. R. *Theoretical and Computational Aspects of Electronic Transport at the Nanoscale*. PhD thesis, University of Dublin, Trinity College, 2007.
- [201] RUNGGER, I., AND SANVITO, S. Algorithm for the Construction of Self-Energies for Electronic Transport Calculations Based on Singularity Elimination and Singular Value Decomposition. *Physical Review B* 78 (2008), 035407.
- [202] SATO, M., NAGAYAMA, K., KATAOKA, N., SASAKI, M., AND HANE, K. Local mechanical properties measured by atomic force microscopy for cultured bovine endothelial cells exposed to shear stress. *Journal of biomechanics* 33, 1 (2000), 127–135.

- [203] SAVASTANO, L. E., BHAMBRI, A., WILKINSON, D. A., AND PANDEY, A. S. Biology of cerebral aneurysm formation, growth, and rupture. In *Intracranial Aneurysms*. Elsevier, 2018, pp. 17–32.
- [204] SAWYER, D. M., PACE, L. A., PASCALE, C. L., KUTCHIN, A. C., ONEILL, B. E., STARKE, R. M., AND DUMONT, A. S. Lymphocytes influence intracranial aneurysm formation and rupture: role of extracellular matrix remodeling and phenotypic modulation of vascular smooth muscle cells. *Journal of neuroinflammation* 13, 1 (2016), 185.
- [205] SCHAFHITZEL, T., VOLLRATH, J. E., GOIS, J. P., WEISKOPF, D., CASTELO, A., AND ERTL, T. Topology-preserving λ_2 -based vortex core line detection for flow visualization. In *Computer Graphics Forum* (2008), vol. 27, Wiley Online Library, pp. 1023–1030.
- [206] SCHAFTENAAR, G., AND NOORDIK, J. H. *Journal of Computer-Aided Molecular Design* 14 (2000), 123.
- [207] SCHIFFRIN, E. L. Novel mechanisms of hypertension and vascular dysfunction. *Nature Reviews Nephrology* 14, 2 (2018), 73–74.
- [208] SFORZA, D. M., PUTMAN, C. M., AND CEBRAL, J. R. Hemodynamics of cerebral aneurysms. *Annual review of fluid mechanics* 41 (2009), 91–107.

- [209] SHABADI, V., SHABADI, S., AND CHANDRAMOHAN, J. Aneurysmal subarachnoid hemorrhage and its practical implications in a peripheral set-up. *International Journal of Advanced Research* 5 (02 2017), 821–826.
- [210] SHANNON, C. E. A mathematical theory of communication. *ACM SIGMOBILE Mobile Computing and Communications Review* 5, 1 (2001), 3–55.
- [211] SHEINBERG, D. L., MCCARTHY, D. J., ELWARDANY, O., BRYANT, J.-P., LUTHER, E., CHEN, S. H., THOMPSON, J. W., AND STARKE, R. M. Endothelial dysfunction in cerebral aneurysms. *Neurosurgical focus* 47, 1 (2019), E3.
- [212] SHIHATA, W. A., MICHELL, D. L., ANDREWS, K. L., AND CHIN-DUSTING, J. P. F. Caveolae: A role in endothelial inflammation and mechanotransduction? *Frontiers in Physiology* 7 (2016), 628.
- [213] SHOJIMA, M., OSHIMA, M., TAKAGI, K., TORII, R., HAYAKAWA, M., KATADA, K., MORITA, A., AND KIRINO, T. Magnitude and role of wall shear stress on cerebral aneurysm: computational fluid dynamic study of 20 middle cerebral artery aneurysms. *Stroke* 35, 11 (2004), 2500–2505.
- [214] SI, H., AND GÄRTNER, K. 3d boundary recovery by constrained delaunay tetrahedralization. *International Journal for Numerical Methods in Engineering* 85, 11 (2011), 1341–1364.

- [215] SIGNORELLI, F., SELA, S., GESUALDO, L., CHEVREL, S., TOLLET, F.,
 PAILLER-MATTEI, C., TACCONI, L., TURJMAN, F., VACCA, A., AND
 SCHUL, D. B. Hemodynamic stress, inflammation, and intracranial aneurysm
 development and rupture: A systematic review. *World Neurosurgery* 115
 (2018), 234 – 244.

- [216] SIU, K. L., LI, Q., ZHANG, Y., GUO, J., YOUN, J. Y., DU, J., AND CAI,
 H. Nox isoforms in the development of abdominal aortic aneurysm. *Redox
 biology* 11 (2017), 118–125.

- [217] SOLER, J. M., ARTACHO, E., GALE, J. D., GARCÍA, A., JUNQUERA, J.,
 ORDEJÓN, P., AND SÁNCHEZ-PORTAL, D. The SIESTA Method for Ab Ini-
 tio Order-N Materials Simulation. *Journal of Physics: Condensed Matter* 14
 (2002), 2745.

- [218] STAFF, B. Medical gallery of blausen medical 2014, 2014. File:Blausen 0055
 ArteryWallStructure.png.

- [219] STANGER, O. H., PEPPER, J. R., AND SVENSSON, L. G. *Surgical Manage-
 ment of Aortic Pathology: Current Fundamentals for the Clinical Management
 of Aortic Disease*. Springer, 2019.

- [220] STARKE, R. M., RAPER, D. M., DING, D., CHALOUHI, N., OWENS, G. K.,
 HASAN, D. M., MEDEL, R., AND DUMONT, A. S. Tumor necrosis factor-
 α modulates cerebral aneurysm formation and rupture. *Translational stroke*

research 5, 2 (2014), 269–277.

- [221] STEINER, T., JUVELA, S., UNTERBERG, A., JUNG, C., FORSTING, M., AND RINKEL, G. European stroke organization guidelines for the management of intracranial aneurysms and subarachnoid haemorrhage. *Cerebrovascular diseases (Basel, Switzerland)* 35 (02 2013), 93–112.
- [222] STEINMAN, D. A. Image-based computational fluid dynamics modeling in realistic arterial geometries. *Annals of biomedical engineering* 30, 4 (2002), 483–497.
- [223] STEINMAN, D. A., MILNER, J. S., NORLEY, C. J., LOWNIE, S. P., AND HOLDSWORTH, D. W. Image-based computational simulation of flow dynamics in a giant intracranial aneurysm. *American Journal of Neuroradiology* 24, 4 (2003), 559–566.
- [224] STRÄTER, A., HUBER, A., RUDOLPH, J., BERNDT, M., RASPER, M., RUMMENY, E. J., AND NADJIRI, J. 4d-flow mri: Technique and applications. In *RöFo-Fortschritte auf dem Gebiet der Röntgenstrahlen und der bildgebenden Verfahren* (2018), vol. 190, © Georg Thieme Verlag KG, pp. 1025–1035.
- [225] SUJUDI, D., AND HAIMES, R. Identification of swirling flow in 3-d vector fields. In *12th Computational Fluid Dynamics Conference* (1995), p. 1715.
- [226] SUN, H. T., SZE, K. Y., TANG, A. Y. S., TSANG, A. C. O., YU, A. C. H., AND CHOW, K. W. Effects of aspect ratio, wall thickness and hypertension in

- the patient-specific computational modeling of cerebral aneurysms using fluid-structure interaction analysis. *Engineering Applications of Computational Fluid Mechanics* 13, 1 (2019), 229–244.
- [227] SUN, X., FU, Y., GU, M., ZHANG, L., LI, D., LI, H., CHIEN, S., SHYY, J. Y.-J., AND ZHU, Y. Activation of integrin $\alpha 5$ mediated by flow requires its translocation to membrane lipid rafts in vascular endothelial cells. *Proceedings of the National Academy of Sciences* 113, 3 (2016), 769–774.
- [228] SUNDERLAND, K., HAFFERMAN, C., CHINTALAPANI, G., AND JIANG, J. Vortex analysis of intra-aneurysmal flow in cerebral aneurysms. *Computational and Mathematical Methods in Medicine* 2016 (2016).
- [229] THEODOTOU, C. B., SNELLING, B. M., SUR, S., HAUSSEN, D. C., PETERSON, E. C., AND ELHAMMADY, M. S. Genetic associations of intracranial aneurysm formation and sub-arachnoid hemorrhage. *Asian journal of neurosurgery* 12, 3 (2017), 374.
- [230] THOMAS, A., OU-YANG, H. D., LOWE-KRENTZ, L., MUZYKANTOV, V. R., AND LIU, Y. Biomimetic channel modeling local vascular dynamics of pro-inflammatory endothelial changes. *Biomicrofluidics* 10, 1 (2016), 014101.
- [231] THOMPSON, G. B., BROWN, R. D., AMIN-HANJANI, S., BRODERICK, J. P., COCKROFT, K. M., CONNOLLY, E. S., DUCKWILER, G. R., HARRIS, C. C.,

- HOWARD, V. J., JOHNSTON, S. C. C., AND ET. AL. Guidelines for the management of patients with unruptured intracranial aneurysms: a guideline for healthcare professionals from the american heart association/american stroke association. *Stroke* (2015), STR-0000000000000070.
- [232] TIBSHIRANI, R. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society. Series B (Methodological)* (1996), 267–288.
- [233] TIBSHIRANI, R., HASTIE, T., NARASIMHAN, B., AND CHU, G. Diagnosis of multiple cancer types by shrunken centroids of gene expression. *Proceedings of the National Academy of Sciences* 99, 10 (2002), 6567–6572.
- [234] TRONIC, F., MALLAT, Z., LEHOUX, S., WASSEF, M., ESPOSITO, B., AND TEDGUI, A. Role of matrix metalloproteinases in blood flow-induced arterial enlargement: interaction with no. *Arterioscler Thromb Vasc Biol* 20, 12 (2000), E120–E126.
- [235] TSUJI, M., ISHIKAWA, T., ISHIDA, F., FURUKAWA, K., MIURA, Y., SHIBA, M., SANO, T., TANEMURA, H., UMEDA, Y., SHIMOSAKA, S., ET AL. Stagnation and complex flow in ruptured cerebral aneurysms: a possible association with hemostatic pattern. *Journal of neurosurgery* 126, 5 (2017), 1566–1572.
- [236] TURJMA, A. S., TURJMAN, F., AND EDELMAN, E. R. Role of fluid dynamics and inflammation in intracranial aneurysm formation. *Circulation* 129, 3 (2014), 373–382.

- [237] TZIMA, E., IRANI-TEHRANI, M., KIOSSES, W. B., DEJANA, E., SCHULTZ, D. A., ENGELHARDT, B., CAO, G., DELISSER, H., AND SCHWARTZ, M. A. A mechanosensory complex that mediates the endothelial cell response to fluid shear stress. *Nature* 437, 7057 (2005), 426.
- [238] UHANA FRÖSEN, TULAMO, R., PAETAU, A., LAAKSAMO, E., KORJA, M., LAAKSO, A., MIKANIELMELÄ, AND HERNESNIEMI, J. Saccular intracranial aneurysm: pathology and mechanisms. *Acta Neuropathologica* 123, 6 (Jun 2012), 773–786.
- [239] UZARSKI, J. S., SCOTT, E. W., AND MCFETRIDGE, P. S. Adaptation of endothelial cells to physiologically-modeled, variable shear stress. *PloS one* 8, 2 (2013), e57004.
- [240] VALEN-SENDSTAD, K., AND STEINMAN, D. Mind the gap: impact of computational fluid dynamics solution strategy on prediction of intracranial aneurysm hemodynamics and rupture status indicators. *American Journal of Neuroradiology* (2013).
- [241] VAN DER MEER, A. D., POOT, A. A., FEIJEN, J., AND VERMES, I. Analyzing shear stress-induced alignment of actin filaments in endothelial cells with a microfluidic assay. *Biomicrofluidics* 4, 1 (2010), 011103.

- [242] VANHOUTTE, P. M., SHIMOKAWA, H., TANG, E. H., AND FELETOU, M.
Endothelial dysfunction and vascular disease. *Acta physiologica* 196, 2 (2009), 193–222.
- [243] VANROSSOMME, A., EKER, O. F., THIRAN, J.-P., COURBEBAILLISSE, G.,
AND BOUDJELTIA, K. Z. Intracranial aneurysms: Wall motion analysis for
prediction of rupture. *American journal of neuroradiology* 36, 10 (2015), 1796–
1802.
- [244] VARBLE, N., TRYLESINSKI, G., XIANG, J., SNYDER, K., AND MENG, H.
Identification of vortex structures in a cohort of 204 intracranial aneurysms.
Journal of The Royal Society Interface 14, 130 (2017), 20170021.
- [245] VARBLE, N., TUTINO, V., YU, J., SONIG, A., SIDDIQUI, A., DAVIES, J.,
AND MENG, H. Shared and distinct rupture discriminants of small and large
intracranial. *Stroke* 49 (2018), 856–864.
- [246] VERSTEEG, H., AND MALALASEKERA, W. An introduction to computational
fluid dynamics: the finite volume method.
- [247] VILLABLANCA, J. P., DUCKWILE, G. R., JAHAN, R., TATESHIMA, S., MAR-
TIN, N. A., FRAZEE, J., GONZALEZ, N. R., SAYRE, J., AND VINUELA, F. V.
Natural history of asymptomatic unruptured cerebral aneurysms evaluated at ct
angiography: growth and rupture incidence and correlation with epidemiologic
risk factors. *Radiology* 269, 1 (2013), 258–265.

- [248] VLAK, M. H., ALGRA, A., BRANDENBURG, R., AND RINKEL, G. J. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *The Lancet Neurology* 10, 7 (2011), 626–636.
- [249] WANG, C., BAKER, B. M., CHEN, C. S., AND SCHWARTZ, M. A. Endothelial cell sensing of flow direction significance. *Arteriosclerosis, thrombosis, and vascular biology* 33, 9 (2013), 2130–2136.
- [250] WANG, G.-X., WANG, S., LIU, L.-L., GONG, M.-F., ZHANG, D., YANG, C.-Y., AND WEN, L. A simple scoring model for prediction of rupture risk of anterior communicating artery aneurysms. *Frontiers in Neurology* 10 (2019), 520.
- [251] WANG, L., LUO, J.-Y., LI, B., TIAN, X. Y., CHEN, L.-J., HUANG, Y., LIU, J., DENG, D., LAU, C. W., WAN, S., ET AL. Integrin-yap/taz-jnk cascade mediates atheroprotective effect of unidirectional shear flow. *Nature* 540, 7634 (2016), 579–582.
- [252] WANG, M., MONTICONE, R. E., AND MCGRAW, K. R. Proinflammatory arterial stiffness syndrome: a signature of large arterial aging. *Journal of vascular research* 55, 4 (2018), 210–223.
- [253] WANG, S., DAI, D., PARAMESWARAN, P. K., KADIRVEL, R., DING, Y.-H., ROBERTSON, A. M., AND KALLMES, D. F. Rabbit aneurysm models mimic

- histologic wall types identified in human intracranial aneurysms. *Journal of neurointerventional surgery* 10, 4 (2018), 411–415.
- [254] WATTON, P., SELIMOVIC, A., RABERGER, N. B., HUANG, P., HOLZAPFEL, G., AND VENTIKOS, Y. Modelling evolution and the evolving mechanical environment of saccular cerebral aneurysms. *Biomechanics and modeling in mechanobiology* 10, 1 (2011), 109–132.
- [255] WEIR, B. Unruptured intracranial aneurysms: a review. *Journal of neurosurgery* 96, 1 (2002), 3–42.
- [256] WEIR, B., DISNEY, L., AND KARRISON, T. Sizes of ruptured and unruptured aneurysms in relation to their sites and the ages of patients. *Journal of neurosurgery* 96, 1 (2002), 64–70.
- [257] WEN, J., DING, G., JIANG, W., WANG, Q., AND ZHENG, T. Numerical simulation of compliant artery bypass grafts using fluid–structure interaction framework. *Asaio Journal* 60, 5 (2014), 533–540.
- [258] WERMER, M. J., VAN DER SCHAAF, I. C., ALGRA, A., AND RINKEL, G. J. Risk of rupture of unruptured intracranial aneurysms in relation to patient and aneurysm characteristics: an updated meta-analysis. *Stroke* 38, 4 (2007), 1404–1410.
- [259] WILLIAMS, L. N., AND BROWN, R. D. Management of unruptured intracranial aneurysms. *Neurology: Clinical Practice* 3, 2 (2013), 99–108.

- [260] WOLF, F., VOGT, F., SCHMITZ-RODE, T., JOCKENHOEVEL, S., AND MELA, P. Bioengineered vascular constructs as living models for in vitro cardiovascular research. *Drug discovery today* 21, 9 (2016), 1446–1455.
- [261] WONG, A. K., LLANOS, P., BORODA, N., ROSENBERG, S. R., AND RAB-BANY, S. Y. A parallel-plate flow chamber for mechanical characterization of endothelial cells exposed to laminar shear stress. *Cellular and molecular bioengineering* 9, 1 (2016), 127–138.
- [262] XANTHIS, I., SOUILHOL, C., SERBANOVIC-CANIC, J., RODDIE, H., KALLI, A. C., FRAGIADAKI, M., WONG, R., SHAH, D. R., ASKARI, J. A., CAN-HAM, L., ET AL. β 1 integrin is a sensor of blood flow direction. *Journal of cell science* 132, 11 (2019), jcs229542.
- [263] XIANG, J., NATARAJAN, S. K., TREMMEL, M., MA, D., MOCCO, J., HOP-KINS, L. N., SIDDIQUI, A. H., LEVY, E. I., AND MENG, H. Hemodynamic-morphologic discriminants for intracranial aneurysm rupture. *Stroke* 42, 1 (2011), 144–152.
- [264] XIANG, J., NATARAJAN, S. K., TREMMEL, M., MA, D., MOCCO, J., HOP-KINS, L. N., SIDDIQUI, A. H., LEVY, E. I., AND MENG, H. Hemodynamic-morphologic discriminants for intracranial aneurysm rupture. *Stroke* 42, 1 (2011), 144–152.

- [265] XIANG, J., NATARAJAN, S. K., TREMMEL, M., MA, D., MOCCO, J., HOPKINS, L. N., SIDDIQUI, A. H., LEVY, E. I., AND MENG, H. Hemodynamic–morphologic discriminants for intracranial aneurysm rupture. *Stroke* 42, 1 (2011), 144–152.
- [266] XIANG, J., SIDDIQUI, A., AND MENG, H. The effect of inlet waveforms on computational hemodynamics of patient-specific intracranial aneurysms. *Journal of biomechanics* 47, 16 (2014), 3882–3890.
- [267] XIONG, G., FIGUEROA, C. A., XIAO, N., AND TAYLOR, C. A. Simulation of blood flow in deformable vessels using subject-specific geometry and spatially varying wall properties. *International journal for numerical methods in biomedical engineering* 27, 7 (2011), 1000–1016.
- [268] XU, L., GU, L., AND LIU, H. Exploring potential association between flow instability and rupture in patients with matched-pairs of ruptured–unruptured intracranial aneurysms. *Biomedical engineering online* 15, 2 (2016), 166.
- [269] XU, L., LEE, T.-Y., AND SHEN, H.-W. An information-theoretic framework for flow visualization. *IEEE Transactions on Visualization and Computer Graphics* 16, 6 (2010), 1216–1224.
- [270] YASUDA, R., STROTHER, C. M., TAKI, W., SHINKI, K., ROYALTY, K., PULFER, K., AND KARMONIK, C. Aneurysm volume-to-ostium area ratio: a parameter useful for discriminating the rupture status of intracranial

- aneurysms. *Neurosurgery* 68, 2 (2011), 310–318.
- [271] ZENG, Y. Endothelial glycocalyx as a critical signalling platform integrating the extracellular haemodynamic forces and chemical signalling. *Journal of cellular and molecular medicine* 21, 8 (2017), 1457–1462.
- [272] ZENG, Z., DURKA, M. J., KALLMES, D. F., DING, Y., AND ROBERTSON, A. M. Can aspect ratio be used to categorize intra-aneurysmal hemodynamics? a study of elastase induced aneurysms in rabbit. *Journal of biomechanics* 44, 16 (2011), 2809–2816.
- [273] ZHANG, Y., JING, L., ZHANG, Y., LIU, J., AND YANG, X. Low wall shear stress is associated with the rupture of intracranial aneurysm with known rupture point: case report and literature review. *BMC Neurol* 16 (2016).
- [274] ZHANG, Y., AND YANG, W.-X. Tight junction between endothelial cells: the interaction between nanoparticles and blood vessels. *Beilstein journal of nanotechnology* 7, 1 (2016), 675–684.
- [275] ZHAO, M., AMIN-HANJANI, S., RULAND, S., CURCIO, A., OSTERGREN, L., AND CHARBEL, F. Regional cerebral blood flow using quantitative mr angiography. *American Journal of Neuroradiology* 28, 8 (2007), 1470–1473.
- [276] ZHOU, G., ZHU, Y., YIN, Y., SU, M., AND LI, M. Association of wall shear stress with intracranial aneurysm rupture: Systematic review and meta-analysis. *Scientific Reports* 7 (12 2017).

- [277] ZHOU, G., ZHU, Y., YIN, Y., SU, M., AND LI, M. Association of wall shear stress with intracranial aneurysm rupture: systematic review and meta-analysis. *Scientific reports* 7, 1 (2017), 5331.
- [278] ZOU, H., AND HASTIE, T. Regularization and variable selection via the elastic net. *J.R. Statist. Soc. B* (2005), 301–320.

Appendix A

Statistics

In this type of predictive modeling, there exists an input-output dataset $(X, Y) \in X \times Y$ with an unknown probability distribution P . The goal of predictive modeling is to find a function $f_n : X \rightarrow Y$, that is determined using a training set $(X_1, Y_1, \dots, (X_n, Y_n))$ of n random pairs distributed as (X, Y) . A desirable solution of f_n is one that, given a new data-point $x \in X$, the resultant $f_n(x)$ is an accurate prediction of the true output $y \in Y$. This desired outcomes not only relies on the chosen function's predictive accuracy, but also of the selecting of relevant variables that are capable of achieving desired predictions. For desired models, it is often preferred to find the prediction function that achieves the desired accuracy while using the minimal amount of variables required: i.e a *parsimonious* model. Brute-force methods of testing all variable combinations becomes increasingly unviable, especially when the

number of variables in a dataset is larger than the number of n data points (cases) available for analysis: often refereed to the "large p , small n paradigm". One type of methodology to determine a desired model is through the use of sparsity-based regularization methods [118, 232, 233, 278]

Section 1

Multiple logistic regression (MLR) analysis looks both to estimate the odds of a dichotomous outcome occurring, and to determine the impact of an individual variable (covariate) in relation to the other covariates in a model. The probability of an outcome occurring in MLR can be calculated as such:

$$\hat{p} = \frac{\exp(b_0 + b_1X_1 + b_2X_2 + \dots + b_pX_p)}{1 + \exp(b_0 + b_1X_1 + b_2X_2 + \dots + b_pX_p)} \quad (\text{A.1})$$

\hat{p} being the probability of the desired outcome, X_1 through X_p as the individual dependent variables applied to the model, and b_1 to b_p being each variable's (respective) regression coefficients. To determine the expected log odds ratios of the model's variables, the *logit* function of the above equation can be calculated:

$$\begin{aligned}
\text{logit}[\hat{p}] &= \ln\left[\frac{\hat{p}}{1-\hat{p}}\right] \\
&= \ln\left[\frac{\frac{\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}{1+\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}}{1-\frac{\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}{1+\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}}\right] \\
&= \ln\left[\frac{\frac{\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}{1+\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}}{\frac{1}{1+\exp(b_0+b_1X_1+b_2X_2+\dots+b_pX_p)}}\right] \tag{A.2} \\
&= \ln[\exp(b_0 + b_1X_1 + b_2X_2 + \dots + b_pX_p)] \\
&= b_0 + b_1X_1 + b_2X_2 + \dots + b_pX_p
\end{aligned}$$

Taking the *logit* of the desired outcome's probability, transforms the occurrence of the event given Xs into a simplified linear function.

For each variable added to a regression model, the resultant R^2 (coefficient of multiple determination) may increase, indicating an improved fit of the data. However applying a large number of variables to a predictive model may result in over-fitting without a significantly large dataset: large p , small n paradigm. In such an event, the R^2 values, regression coefficients, and any statistical significance (p -values) determined may be misleading. To reduce the initial choices of variables in assessed predictive models, the correlation between variables were determined. The correlation of data can be determine by:

$$r_{jk} = \frac{s_{jk}}{s_j s_k} = \frac{\sum_{i=1}^n (x_{ij} - \bar{x}_j)(x_{ik} - \bar{x}_k)}{\sqrt{\sum_{i=1}^n (x_{ij} - \bar{x}_j)^2} \sqrt{\sum_{i=1}^n (x_{ik} - \bar{x}_k)^2}} \tag{A.3}$$

with r as the Pearson correlation coefficient between variables x_j and x_k , n as the sample size, and \bar{x} is a variable sample mean. Correlations between the variables are often displayed via a correlation table:

$$R = \begin{bmatrix} 1 & r_{12} & r_{13} & \dots & r_{1p} \\ r_{21} & 1 & r_{23} & \dots & r_{2p} \\ r_{31} & r_{32} & 1 & \dots & r_{3p} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ r_{p1} & r_{p2} & r_{p3} & \dots & 1 \end{bmatrix}$$

Initial correlation analysis of all available geometric and hemodynamic variables was performed to eliminate highly correlated variables from analysis: i.e aneurysm volume and surface area are highly correlated so surface area was removed from analysis.

From the remaining variables, stepwise MLR was implemented to determine the parsimonious model. In stepwise regression, a linear regression is first performed for each variable X one at a time, and the variable with the highest R^2 is kept for the model. Next, a multiple regression step is performed with the kept variable and each remaining variable. The variable with the largest increase in R^2 , if the p value of the R^2 is below a desired cutoff (<0.05), is added to the model. The calculation of the p value of an increase in R^2 resulting from the increasing of X variable(s) from a to

b is as follows:

$$p_{ab} = \frac{(R_b^2 - R_a^2)/(b - a)}{(1 - R_b^2)/(n - b - 1)} \quad (\text{A.4})$$

with the total sample size n .

Each time a new variable is added to the model, the impact of removing any of the other variables (already added to the model) on outcomes is tested. The chosen (removed) variable is excluded from the model if it does not make R^2 significantly worse. This process is continued till adding any new variables does not increase R^2 and removing any X variables does not significantly decrease R^2 .

In the event that all of the independent variables in the model are completely uncorrelated with each other, the interpretation of coefficients are as such:

$$OR = \exp(b_1)^z \quad (\text{A.5})$$

Where z is the number of unit changes for a variable X , and OR is the odds ratio resultant from said change. When the variables are not uncorrelated, the $OR = \exp^z b_1$ is expressed as the change of unit z for a variable *adjusted in relation to the impacts of the other variables in the model*. This stresses the need to assess collinearity between variables prior to model assessment.

in Section ??.

Section 2

Limitations may arise in applying multiple logistic regression analysis to data sets with a large number of variables in relation to the number of samples.

The Nearest Shrunken Centroid (NSC) method is a statistical classification methodology developed by Tibshirani et al.[233] as a means to improve predictions in high-dimensional data. Additionally, a 2014 study by Finch [86] compared a number of methods for statistical group prediction. The NSC method was found to be robust in terms of accuracy and identification of predictor variables over other methods when dealing with high-dimensional datasets. Initially developed for predictions within genetic data, NSC aims to shrink class (e.g. rupture status) centroids towards the *overall* centroid of the entire data-set after standardizing by the within-class deviation for each variable. This standardization of the resultant centroid gives higher impact to variables whose expression is more stable within samples of the same class. Due to the number of geometric and hemodynamic variables that may impact IA rupture, the NSC method was investigated as a useful methodology to help predict rupture potential within our data-set.

In our study, x_{ijk} represents the value of quantified variables $i = 1, \dots, p$ in aneurysm $j = 1, \dots, n_k$ of class k . The mean for variable i in each class k is calculated

$$\bar{x}_{ik} = \sum_{j=1}^{n_k} \frac{x_{ijk}}{n_k} \quad (\text{A.6})$$

and the *overall* mean for variable i is calculated.

$$\bar{x}_i = \sum_{k=1}^K \sum_{j=1}^{n_k} \frac{x_{ijk}}{n_k} \quad (\text{A.7})$$

A t -statistic value, d_{ik} , for each variable is calculated, comparing class k to the overall mean:

$$d_{ik} = \frac{\bar{x}_{ik} - \bar{x}_i}{m_k \cdot (s_i + s_0)} \quad (\text{A.8})$$

with

$$s_i^2 = \frac{1}{n - K} \sum_{k=1}^K \sum_{j=1}^{n_k} (x_{ij} - \bar{x}_{ik})^2 \quad (\text{A.9})$$

and

$$m_k = \sqrt{1/n_k + 1/n}, \text{ where } n = \sum_{k=1}^K n_k \quad (\text{A.10})$$

The s_0 in (A.8) is used as a *regularization parameter* to help protect from large d_{ik} values occurring from variables at low expression levels. The value of s_0 is determined

as the median value of s_i over the set of variables.

With the inclusion of d_{ik} , the class centroid can be rewritten as:

$$\bar{x}_{ik} = \bar{x}_i + m_k(s_i + S_0)d_{ik} \quad (\text{A.11})$$

The NSC method shrinks each d_{ik} toward zero, creating a new value d'_{ik} which generates a new shrunken centroid value

$$\bar{x}'_{ik} = \bar{x}_i + m_k(s_i + s_0)d'_{ik} \quad (\text{A.12})$$

The value of shrinkage is determined through *soft thresholding*, where the absolute value of d_{ik} is reduced by Δ and is given the value of 0 if the result is < 0 , with

$$d'_{ik} = \text{sign}(d_{ik})(|d_{ik}| - \Delta)_+ \quad (\text{A.13})$$

This combination of shrinkage and *soft thresholding* can result in many 'noisy' \bar{x}_{ik} being close to the value of the overall mean \bar{x}_i . If Δ shrinks d_{ik} to zero for all k , then the centroid for variable i is the same for all classes and is excluded from prediction analysis.

The value of Δ is determined through cross-validation: fitting the model for many values of Δ and determining the level of error per chosen Δ . The Δ resulting in the smallest error was chosen for our prediction.

Section 3

Elastic Net Regularization (ENR) overcomes some of the limitations of the LASSO selection method, primarily being able to accurately handle data sets with a high number of variables in relation to the sample size [80, 232]. Additionally, the ENR method is able to handle data sets with groups of highly correlated variables.

ENR solves two optimization problems:

$$\begin{aligned} \tilde{\beta} = \arg \min_{\beta} & \sum_{i=1}^N (y_i - (X\beta)_i)^2 \\ \text{subject to} & \sum_{j=1}^p |\beta_j| \leq t_1 \text{ and } \sum_{j=1}^p \beta_j^2 \leq t_2 \end{aligned} \tag{A.14}$$

where a penalty is placed on the L_1 norm ($\sum_{j=1}^p |\beta_j|$) and the L_2 norm ($\sum_{j=1}^p \beta_j^2 \leq t_2$) of the regression coefficients. The purpose of these penalties are as follows: L_1 performs variable selection by setting some coefficients to 0, and L_2 works toward group selection by shrinking the coefficients of correlated variables toward each other. Re-writing equation A.14 in the Lagrangian form using two tuning parameters (λ_1 and λ_2)

is as follows:

$$\tilde{\beta} = \arg \min_{\beta} \left(\sum_{i=1}^N (y_i - (X\beta)_i)^2 + \lambda_1 \sum_{j=1}^p |\beta_j| + \lambda_2 \sum_{j=1}^p \beta_j^2 \right) \quad (\text{A.15})$$

The choice of tuning parameter values is performed by analyzing an array of λ_2 values (0, 0.01, 0.1, 1, 10, and 100). For each value in the array, the LARS-EN algorithm calculates the resultant λ_1 value. The *lambda*₁ value that yields the smallest *k*-fold cross validation error, and its *lambda*₂ value used to generate it, are used as the tuning parameters for the ENR method.

Section 4

Section 5

To assess the diagnostic ability of predictive model(s), a receiver operating characteristic curve (ROC) is often deployed (REFERENCES). The ROC curve assesses a model's predictive true positive rate (TPR) against its false positive rate (FPR) as a means to determine overall predictive strength (HANLEY). From a statistical perspective, ROC analysis can be considered as a plot of the power (probability of a test correctly rejecting the null hypothesis when an alternative hypothesis is true)

$$\begin{aligned}
TPR &= \frac{\Sigma TruePositive}{\Sigma ConditionPositive} \\
FPR &= \frac{\Sigma FalsePpositive}{\Sigma ConditionNegative} \\
FNR &= \frac{\Sigma FalseNegative}{\Sigma ConditionPositive} \\
Specificity &= \frac{\Sigma TrueNegative}{\Sigma ConditionNegative}
\end{aligned} \tag{A.16}$$

When dealing with a binary classification, as per this study, the predictive test measure for each instance is denoted by a continuous random variable (x). Given a desired threshold (T), each instance is positive if $x > T$ and negative if $x < T$. Setting the probability distribution functions of the positive and negative values of x to $f_p(x)$ and $f_n(x)$ respectively, the . Given this, TPR is calculated as:

$$TPR(T) = \int_T^{\infty} f_p(x) dx \tag{A.17}$$

and the FNR as:

$$FPR(T) = 1 - \int_T^{\infty} f_n(x) dx \tag{A.18}$$

The ROC curve is generated by plotting $TPR(T)$ against $FPR(T)$ parametrically, varying across T, or as a plot of:

$$ROC(T) = 1 - f_p(f_n^{-1}(1 - T)) \quad (A.19)$$

over T from $[0,1]$ where $f_p^{-1}(1-T) = \inf$

Comparing the resultant ROC curves across multiple models provides the selection of the desired model based off of varying predictive accuracies. To quantify the predictive accuracy, the area under the curve (AUC) of the ROC curve is calculated, as it equals the probability of a classifier ranking a positive instance higher than a negative instance (both chosen at random).

$$\begin{aligned} A &= \int_{-\infty}^{\infty} TPR(T)FPR'(T)dT \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} I(T' > T)f_1(T')f_0(T)dT'dT = P(X_1 > X_0) \end{aligned} \quad (A.20)$$

The initial integral has reversed boundaries due to larger T values having a lower value on the x-axis.

Section 5

Docendi eligendi sit et, pri ea dicam eligendi percipitur, has soleat dolores convenire te. Sed altera placerat an, id verterem abhorreant interesset mea. Eum at ceteros

efficiantur. Eos id voluptaria efficiendi comprehensam.

In mel modo dicam vocibus, eruditi consecutuer vim no, cu quaestio instructor eum.

Justo nostrud fuisset ea mea, eam an libris repudiandae vituperatoribus. Est choro
corrumpit definitionem at. Vel sint adhuc vocibus ea, illud epicuri eos no. Sea simul
officiis ea, et qui veri invidunt appellantur. Vix et eros ancillae pertinax.

Appendix B

Sample Code

The method for vortex identification for this study is a modification from previous work[228]. The calculation of vortex cores was based on in-house C++/Python codes derived from the open-source Vascular Modelling ToolKit (VMTK) [6]. Prior to any calculations, velocity data is first re-sampled onto a rectilinear grid whose voxel size is 0.2mm.

In the first step, the classic λ_2 method by Jeong and Hussain [120] was used to define the negative λ_2 region (*i.e* $\lambda_2 < 0$). Then, in the second step, vortex core lines were estimated by the method proposed by Sujudi and Haines [225]. In essence, in the negative λ_2 region, a local velocity vector \bar{v} lies along a vortex core line if the following two conditions hold: (1) the 3×3 spatial gradient matrix of \bar{v} has two complex eigenvalues and one real eigenvalue and (2) the 3×3 spatial gradient matrix of \bar{v} has




Figure B.1: Two examples illustrating the relationship between the angular histogram and NE: (a) a simple laminar flow case and (b) a rotational flow (eddy) case. In both cases, the right and left plots are the vector flow field and the histogram of angular vector direction, respectively. Vector fields were decimated by a factor of 3 for better visualization.

an eigenvector $\vec{\alpha}$ corresponding to the above-mentioned real eigenvalue. Now, if we define a new scalar value K as follows,

$$K(x, y, z) = \begin{cases} |\dot{dot}(\bar{v}, \vec{\alpha})|, & \text{if } \lambda_2 < 0 \\ 0, & \text{Otherwise} \end{cases} \quad (\text{B.1})$$

where $|\cdot|$ is an absolute operator. Of note, in Eqn. 2, both the \bar{v} and $\vec{\alpha}$ are normalized and therefore, the scalar field K defined above is bounded between 0 and 1. If the $K(x, y, z)$ is close to 1 then the location (x, y, z) is within the proximity of the vortex core line as suggested by Sujudi and Haimes [225].

In the third step, we calculated local normalized entropy (NE) of velocity directions [210] following work in the flow visualization literature (e.g. [159, 269]). The NE is close to 0 if the velocity direction closely concentrates one value out of N possible values (see Fig. B.1(a); $NE=0.05$). In contrast, the entropy measure NE becomes

0.95 if the probability of velocity directions is almost equally likely, as shown in Fig B.1(b). Given an arbitrary voxel located at (x, y, z) within the dome of an IA, we selected a fixed volume of interest (VOI; $N_x \times N_y \times N_z$; $N_x = N_y = N_z = 11$ in this study) centered at the voxel. One additional metric $H(x, y, z)$ can be obtained by combining $K(x, y, z)$ together with the $NE(x, y, z)$ as follows,

$$H(x, y, z) = K(x, y, z) * NE(x, y, z) \quad (\text{B.2})$$

$H(x, y, z)$ is a scalar field representing the likelihood of residing within a vortex core region for a location (x, y, z) . H also has a normalized range between 0 and 1. Thus, based on a fixed threshold, the vortex core region in this study can be obtained using the classic Marching-cube method [157]. in this study, 0.30 was used as the threshold for all data sets.

HelloWorld.c

```
// HelloWorld.c
// C program to display 'Hello, World!' in the terminal.
//
// Compilation:
// gcc -g -Wall HelloWorld.c -o HelloWorld.x
//
// Execution:
```

```
// ./HelloWorld.x

// Standard headers
#include <stdio.h>

// main() begins
int main() {

    // Print the message
    printf("\n Hello, World!\n\n");

    // Indicate the termination of main()
    return 0;
}
// main() ends
```

Appendix C

Letters of Permission

Include letters of permission from journal editors and/or other sources from which you may have used materials (images, information, etc.) in this this work.

These materials may also be submitted separately to the Graduate School as a single, well-organized PDF file.

Appendix D

Cellular Biology

TUNEL-assay

Terminal deoxynucleotidyl transferase dUTP-biotin nick end labeling (TUNEL) is an assay for detecting DNA fragmentation: an aspect of cellular damage and apoptosis. TUNEL uses the enzyme terminal deoxynucleotidyl transferase (TdT) to attach labeled deoxyuridine triphosphate (dUTP) onto the 3'-hydroxyl termini of internucleosomal DNA fragmentation. Modification of dUTP through the addition of fluorophores or haptens, such as biotin, allow for DNA fragments to be detected directly using a fluorescently-modified nucleotide and fluorescence microscopy or flow cytometry.

VCAM-1

VCAM-1 is a member of the immunoglobulin superfamily (cell surface and soluble proteins involved in the recognition and/or binding of cells) and encodes a cell surface sialoglycoprotein (sialic acid and glycoprotein combination) expressed by cytokine-activated endothelium. This membrane protein acts as a ligand for leukocyte-endothelial cell adhesion, signal transduction, and may play a role in the development of atherosclerotic and/or inflammatory based pathologies. Molecules containing VCAM-1 counterreceptors (VLA-4 on monocytes and lymphocytes) can adhere to VCAM-1 activated cells[131]. Bound leukocytes may undergo polarized motility into the vascular wall, disrupting the cellular and matrix components of the vasculature, and degrading endothelial cell permeability.