

5c. Mathematical notation

The table below sets out the notation that may be used in A Level Mathematics A. Students will be expected to understand this notation without need for further explanation.

1	Set Notation	
1.1	\in	is an element of
1.2	\notin	is not an element of
1.3	\subseteq	is a subset of
1.4	\subset	is a proper subset of
1.5	$\{x_1, x_2, \dots\}$	the set with elements x_1, x_2, \dots
1.6	$\{x: \dots\}$	the set of all x such that ...
1.7	$n(A)$	the number of elements in set A
1.8	\emptyset	the empty set
1.9	\mathcal{E}	the universal set
1.10	A'	the complement of the set A
1.11	\mathbb{N}	the set of natural numbers, $\{1, 2, 3, \dots\}$
1.12	\mathbb{Z}	the set of integers, $\{0, \pm 1, \pm 2, \pm 3, \dots\}$
1.13	\mathbb{Z}^+	the set of positive integers, $\{1, 2, 3, \dots\}$
1.14	\mathbb{Z}_0^+	the set of non-negative integers, $\{0, 1, 2, 3, \dots\}$
1.15	\mathbb{R}	the set of real numbers
1.16	\mathbb{Q}	the set of rational numbers, $\left\{\frac{p}{q}: p \in \mathbb{Z}, q \in \mathbb{Z}^+\right\}$
1.17	\cup	union
1.18	\cap	intersection
1.19	(x, y)	the ordered pair x, y
1.20	$[a, b]$	the closed interval $\{x \in \mathbb{R}: a \leq x \leq b\}$
1.21	$[a, b)$	the interval $\{x \in \mathbb{R}: a \leq x < b\}$
1.22	$(a, b]$	the interval $\{x \in \mathbb{R}: a < x \leq b\}$
1.23	(a, b)	the open interval $\{x \in \mathbb{R}: a < x < b\}$
2	Miscellaneous Symbols	
2.1	$=$	is equal to
2.2	\neq	is not equal to

2.3	\equiv	is identical to or is congruent to
2.4	\approx	is approximately equal to
2.5	∞	infinity
2.6	\propto	is proportional to
2.7	\therefore	therefore
2.8	\because	because
2.9	$<$	is less than
2.10	\leq, \leq	is less than or equal to, is not greater than
2.11	$>$	is greater than
2.12	\geq, \geq	is greater than or equal to, is not less than
2.13	$p \Rightarrow q$	p implies q (if p then q)
2.14	$p \Leftarrow q$	p is implied by q (if q then p)
2.15	$p \Leftrightarrow q$	p implies and is implied by q (p is equivalent to q)
2.16	a	first term for an arithmetic or geometric sequence
2.17	l	last term for an arithmetic sequence
2.18	d	common difference for an arithmetic sequence
2.19	r	common ratio for a geometric sequence
2.20	S_n	sum to n terms of a sequence
2.21	S_∞	sum to infinity of a sequence
3	Operations	
3.1	$a + b$	a plus b
3.2	$a - b$	a minus b
3.3	$a \times b, ab, a.b$	a multiplied by b
3.4	$a \div b, \frac{a}{b}$	a divided by b
3.5	$\sum_{i=1}^n a_i$	$a_1 + a_2 + \dots + a_n$
3.6	$\prod_{i=1}^n a_i$	$a_1 \times a_2 \times \dots \times a_n$
3.7	\sqrt{a}	the non-negative square root of a
3.8	$ a $	the modulus of a
3.9	$n!$	n factorial: $n! = n \times (n-1) \times \dots \times 2 \times 1, n \in \mathbb{N}; 0! = 1$

3.10	$\binom{n}{r} {}^nC_r, {}^nC_r$	the binomial coefficient $\frac{n!}{r!(n-r)!}$ for $n, r \in \mathbb{Z}_0^+, r \leq n$ or $\frac{n(n-1)\dots(n-r+1)}{r!}$ for $n \in \mathbb{Q}, r \in \mathbb{Z}_0^+$
4	Functions	
4.1	$f(x)$	the value of the function f at x
4.2	$f: x \mapsto y$	the function f maps the element x to the element y
4.3	f^{-1}	the inverse function of the function f
4.4	gf	the composite function of f and g which is defined by $gf(x) = g(f(x))$
4.5	$\lim_{x \rightarrow a} f(x)$	the limit of $f(x)$ as x tends to a
4.6	$\Delta x, \delta x$	an increment of x
4.7	$\frac{dy}{dx}$	the derivative of y with respect to x
4.8	$\frac{d^n y}{dx^n}$	the n th derivative of y with respect to x
4.9	$f'(x), f''(x), \dots, f^{(n)}(x)$	the first, second, ..., n th derivatives of $f(x)$ with respect to x
4.10	\dot{x}, \ddot{x}, \dots	the first, second, ... derivatives of x with respect to t
4.11	$\int y \, dx$	the indefinite integral of y with respect to x
4.12	$\int_a^b y \, dx$	the definite integral of y with respect to x between the limits $x = a$ and $x = b$
5	Exponential and Logarithmic Functions	
5.1	e	base of natural logarithms
5.2	$e^x, \exp x$	exponential function of x
5.3	$\log_a x$	logarithm to the base a of x
5.4	$\ln x, \log_e x$	natural logarithm of x
6	Trigonometric Functions	
6.1	$\left. \begin{matrix} \sin, \cos, \tan \\ \operatorname{cosec}, \sec, \cot \end{matrix} \right\}$	the trigonometric functions
6.2	$\left. \begin{matrix} \sin^{-1}, \cos^{-1}, \tan^{-1} \\ \arcsin, \arccos, \arctan \end{matrix} \right\}$	the inverse trigonometric functions
6.3	$^\circ$	degrees
6.4	rad	radians

9	Vectors	
9.1	$\mathbf{a}, \underline{a}, \hat{a}$	the vector $\mathbf{a}, \underline{a}, \hat{a}$; these alternatives apply throughout section 9
9.2	\overrightarrow{AB}	the vector represented in magnitude and direction by the directed line segment AB
9.3	$\hat{\mathbf{a}}$	a unit vector in the direction of \mathbf{a}
9.4	$\mathbf{i}, \mathbf{j}, \mathbf{k}$	unit vectors in the directions of the cartesian coordinate axes
9.5	$ \mathbf{a} , a$	the magnitude of \mathbf{a}
9.6	$ \overrightarrow{AB} , AB$	the magnitude of \overrightarrow{AB}
9.7	$\begin{pmatrix} a \\ b \end{pmatrix}, a\mathbf{i} + b\mathbf{j}$	column vector and corresponding unit vector notation
9.8	\mathbf{r}	position vector
9.9	\mathbf{s}	displacement vector
9.10	\mathbf{v}	velocity vector
9.11	\mathbf{a}	acceleration vector
11	Probability and Statistics	
11.1	$A, B, C, \text{ etc.}$	events
11.2	$A \cup B$	union of the events A and B
11.3	$A \cap B$	intersection of the events A and B
11.4	$P(A)$	probability of the event A
11.5	A'	complement of the event A
11.6	$P(A B)$	probability of the event A conditional on the event B
11.7	$X, Y, R, \text{ etc.}$	random variables
11.8	$x, y, r, \text{ etc.}$	values of the random variables X, Y, R etc.
11.9	x_1, x_2, \dots	values of observations
11.10	f_1, f_2, \dots	frequencies with which the observations x_1, x_2, \dots occur
11.11	$p(x), P(X=x)$	probability function of the discrete random variable X
11.12	p_1, p_2, \dots	probabilities of the values x_1, x_2, \dots of the discrete random variable X
11.13	$E(X)$	expectation of the random variable X
11.14	$\text{Var}(X)$	variance of the random variable X

11.15	\sim	has the distribution
11.16	$B(n, p)$	binomial distribution with parameters n and p , where n is the number of trials and p is the probability of success in a trial
11.17	q	$q = 1 - p$ for binomial distribution
11.18	$N(\mu, \sigma^2)$	Normal distribution with mean μ and variance σ^2
11.19	$Z \sim N(0, 1)$	standard Normal distribution
11.20	ϕ	probability density function of the standardised Normal variable with distribution $N(0, 1)$
11.21	Φ	corresponding cumulative distribution function
11.22	μ	population mean
11.23	σ^2	population variance
11.24	σ	population standard deviation
11.25	\bar{x}	sample mean
11.26	s^2	sample variance
11.27	s	sample standard deviation
11.28	H_0	Null hypothesis
11.29	H_1	Alternative hypothesis
11.30	r	product moment correlation coefficient for a sample
11.31	ρ	product moment correlation coefficient for a population
12	Mechanics	
12.1	kg	kilograms
12.2	m	metres
12.3	km	kilometres
12.4	m/s, m s ⁻¹	metres per second (velocity)
12.5	m/s ² , m s ⁻²	metres per second per second (acceleration)
12.6	F	Force or resultant force
12.7	N	Newton
12.8	N m	Newton metre (moment of a force)
12.9	t	time
12.10	s	displacement
12.11	u	initial velocity
12.12	v	velocity or final velocity

12.13	a	acceleration
12.14	g	acceleration due to gravity
12.15	μ	coefficient of friction

5d. Mathematical formulae and identities

Learners must be able to use the following formulae and identities for A Level mathematics, without these formulae and identities being provided, either in these forms or in equivalent forms. These formulae and identities may only be provided where they are the starting point for a proof or as a result to be proved.

Pure Mathematics

Quadratic Equations

$$ax^2 + bx + c = 0 \text{ has roots } \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Laws of Indices

$$a^x a^y \equiv a^{x+y}$$

$$a^x \div a^y \equiv a^{x-y}$$

$$(a^x)^y \equiv a^{xy}$$

Laws of Logarithms

$$x = a^n \Leftrightarrow n = \log_a x \text{ for } a > 0 \text{ and } x > 0$$

$$\log_a x + \log_a y \equiv \log_a (xy)$$

$$\log_a x - \log_a y \equiv \log_a \left(\frac{x}{y} \right)$$

$$k \log_a x \equiv \log_a (x^k)$$

Coordinate Geometry

A straight line graph, gradient m passing through (x_1, y_1) has equation

$$y - y_1 = m(x - x_1)$$

Straight lines with gradients m_1 and m_2 are perpendicular when $m_1 m_2 = -1$

Sequences

General term of an arithmetic progression:

$$u_n = a + (n - 1)d$$

General term of a geometric progression:

$$u_n = ar^{n-1}$$

Trigonometry

In the triangle ABC

$$\text{Sine rule: } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$\text{Cosine rule: } a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{Area} = \frac{1}{2}ab \sin C$$

$$\cos^2 A + \sin^2 A \equiv 1$$

$$\sec^2 A \equiv 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A \equiv 1 + \cot^2 A$$

$$\sin 2A \equiv 2 \sin A \cos A$$

$$\cos 2A \equiv \cos^2 A - \sin^2 A$$

$$\tan 2A \equiv \frac{2 \tan A}{1 - \tan^2 A}$$

Mensuration

Circumference and Area of circle, radius r and diameter d :

$$C = 2\pi r = \pi d \quad A = \pi r^2$$

Pythagoras' Theorem: In any right-angled triangle where a , b and c are the lengths of the sides and c is the hypotenuse:

$$c^2 = a^2 + b^2$$

Area of a trapezium = $\frac{1}{2}(a + b)h$, where a and b are the lengths of the parallel sides and h is their perpendicular separation.

Volume of a prism = area of cross section \times length

For a circle of radius r , where an angle at the centre of θ radians subtends an arc of length s and encloses an associated sector of area A :

$$s = r\theta \quad A = \frac{1}{2}r^2\theta$$

Calculus and Differential Equations

Differentiation

Function

$$x^n$$

$$\sin kx$$

$$\cos kx$$

$$e^{kx}$$

$$\ln x$$

$$f(x) + g(x)$$

$$f(x)g(x)$$

$$f(g(x))$$

Derivative

$$nx^{n-1}$$

$$k \cos kx$$

$$-k \sin kx$$

$$ke^{kx}$$

$$\frac{1}{x}$$

$$f'(x) + g'(x)$$

$$f'(x)g(x) + f(x)g'(x)$$

$$f'(g(x))g'(x)$$

Integration

Function

Integral

$$x^n \quad \frac{1}{n+1}x^{n+1} + c, \quad n \neq -1$$

$$\cos kx \quad \frac{1}{k} \sin kx + c$$

$$\sin kx \quad -\frac{1}{k} \cos kx + c$$

$$e^{kx} \quad \frac{1}{k} e^{kx} + c$$

$$\frac{1}{x} \quad \ln|x| + c, \quad x \neq 0$$

$$f'(x) + g'(x) \quad f(x) + g(x) + c$$

$$f'(g(x))g'(x) \quad f(g(x)) + c$$

$$\text{Area under a curve} = \int_a^b y \, dx \quad (y \geq 0)$$

Vectors

$$|x\mathbf{i} + y\mathbf{j}| = \sqrt{x^2 + y^2}$$

$$|x\mathbf{i} + y\mathbf{j} + z\mathbf{k}| = \sqrt{x^2 + y^2 + z^2}$$

Mechanics

Forces and Equilibrium

$$\text{Weight} = \text{mass} \times g$$

$$\text{Friction: } F \leq \mu R$$

$$\text{Newton's second law in the form: } F = ma$$

Kinematics

For motion in a straight line with variable acceleration:

$$v = \frac{dr}{dt} \quad a = \frac{dv}{dt} = \frac{d^2r}{dt^2}$$

$$r = \int v \, dt \quad v = \int a \, dt$$

$$v = \frac{ds}{dt} \quad a = \frac{dv}{dt} = \frac{d^2s}{dt^2}$$

$$r = \int v \, dt \quad v = \int a \, dt$$

Statistics

$$\text{The mean of a set of data: } \bar{x} = \frac{\sum x}{n} = \frac{\sum fx}{\sum f}$$

$$\text{The standard Normal variable: } Z = \frac{X - \mu}{\sigma} \quad \text{where } X \sim N(\mu, \sigma^2)$$

Learners will be given the following formulae sheet in each question paper.

Formulae A Level Mathematics A (H240)

Arithmetic series

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \text{ for } |r| < 1$$

Binomial series

$$(a+b)^n = a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_r a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N}),$$

$$\text{where } {}^nC_r = {}_nC_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Differentiation

$$f(x)$$

$$\tan kx$$

$$\sec x$$

$$\cot x$$

$$\operatorname{cosec} x$$

$$f'(x)$$

$$k \sec^2 kx$$

$$\sec x \tan x$$

$$-\operatorname{cosec}^2 x$$

$$-\operatorname{cosec} x \cot x$$

$$\text{Quotient Rule } y = \frac{u}{v}, \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Integration

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$$

$$\int f'(x)(f(x))^n dx = \frac{1}{n+1}(f(x))^{n+1} + c$$

Integration by parts $\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$

Small angle approximations

$\sin \theta \approx \theta$, $\cos \theta \approx 1 - \frac{1}{2}\theta^2$, $\tan \theta \approx \theta$ where θ is measured in radians

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad (A \pm B \neq (k + \frac{1}{2})\pi)$$

Numerical methods

Trapezium rule: $\int_a^b y dx \approx \frac{1}{2}h \{(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})\}$, where $h = \frac{b-a}{n}$

The Newton-Raphson iteration for solving $f(x) = 0$: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A)P(B|A) = P(B)P(A|B) \quad \text{or} \quad P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Standard deviation

$$\sqrt{\frac{\sum(x - \bar{x})^2}{n}} = \sqrt{\frac{\sum x^2}{n} - \bar{x}^2} \quad \text{or} \quad \sqrt{\frac{\sum f(x - \bar{x})^2}{\sum f}} = \sqrt{\frac{\sum fx^2}{\sum f} - \bar{x}^2}$$

The binomial distribution

If $X \sim B(n, p)$ then $P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}$, Mean of X is np , Variance of X is $np(1-p)$

Hypothesis test for the mean of a normal distribution

If $X \sim N(\mu, \sigma^2)$ then $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ and $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$

Percentage points of the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p , the table gives the value of z such that $P(Z \leq z) = p$.

p	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
z	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Kinematics

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$s = \frac{1}{2}(u + v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2}at^2$$

Motion in two dimensions

$$\mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$$

$$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$$

$$\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$$