

AN EXTENSION TO THE TANGENT SEQUENCE MARTINGALE INEQUALITY

STEPHEN MONTGOMERY-SMITH AND SHIH-CHI SHEN

ABSTRACT. For each $1 < p < \infty$, there exists a positive constant c_p , depending only on p , such that the following holds. Let (d_k) , (e_k) be real-valued martingale difference sequences. If for all bounded nonnegative predictable sequence (s_k) and all positive integers k we have

$$E[s_k \vee |e_k|] \leq E[s_k \vee |d_k|]$$

then for all positive integers n we have

$$\left\| \sum_{k=1}^n e_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p.$$

1. INTRODUCTION

Let (Ω, \mathcal{F}, P) be a probability space, and let (\mathcal{F}_k) be a filtration on (Ω, \mathcal{F}, P) . (We will suppose that $\mathcal{F}_0 = \{\emptyset, \Omega\}$.) If an adapted sequence (d_k) is a real-valued martingale difference sequence, Burkholder's inequality [2] shows that for any $1 < p < \infty$ that there exists a positive constant c_p depending only on p such that for all $\varepsilon_k \in \{1, -1\}$ and all positive integers n that

$$\left\| \sum_{k=1}^n \varepsilon_k d_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p.$$

More generally, if (v_k) is a predictable sequence bounded in absolute value by 1, then

$$\left\| \sum_{k=1}^n v_k d_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p.$$

A different approach to this inequality was proposed by Kwapień and Woyczynski [8] (see also [9]). Two adapted sequence (f_k) and (g_k) are said to be tangent if for each $k \geq 1$, we have that the law of f_k conditionally on \mathcal{F}_{k-1} is the same as the law of g_k conditionally on \mathcal{F}_{k-1} , that is,

$$P(f_k < \lambda | \mathcal{F}_{k-1}) = P(g_k < \lambda | \mathcal{F}_{k-1})$$

for all real numbers λ . Answering a conjecture of Kwapień and Woyczinski [8], it was proved by Hitczenko [4] (see also [14]) that for $1 < p < \infty$ that there exists a positive constant c_p , depending only on p , such that if (d_k) and (e_k) are martingale difference sequences and $(d_k), (e_k)$ are tangent, then for all positive integers n

$$(1) \quad \left\| \sum_{k=1}^n e_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p.$$

The purpose of this paper is to provide a common generalization to these two results.

Theorem 1. *For each $1 < p < \infty$, there exists a positive constant c_p , depending only on p , such that the following holds. Let $(d_k), (e_k)$ be real-valued martingale difference sequences. If for all bounded nonnegative predictable sequence (s_k) and all positive integers k we have*

$$(2) \quad E[s_k \vee |e_k|] \leq E[s_k \vee |d_k|]$$

then for all positive integers n we have equation (1).

This is essentially equivalent to another result, which concerns martingales in a specific situation. We will consider the probability space $[0, 1]^{\mathbf{N}}$ equipped with the product Lebesgue measure \mathcal{L} , and consider the filtration (\mathcal{L}_k) , where \mathcal{L}_k is the minimal σ -field for which the first k coordinate functions of $[0, 1]^{\mathbf{N}}$ are measurable. Then two sequences (d_k) and (e_k) are tangent if

$$e_k(x_1, \dots, x_k) = d_k(x_1, \dots, x_{k-1}, \phi_k(x_1, \dots, x_k))$$

where $(\phi_k : [0, 1]^k \rightarrow [0, 1])$ is a sequence of measurable functions such that $\phi_k(x_1, \dots, x_{k-1}, \cdot)$ is a measure preserving map for almost all x_1, \dots, x_{k-1} .

We will consider a more general situation. Suppose we have a sequence of linear operators $(T_k(x_1, \dots, x_{k-1}))$, depending measurably upon $(x_k) \in [0, 1]^{\mathbf{N}}$, that are bounded operators on both $L_1[0, 1]$ and $L_\infty[0, 1]$ with norm 1. Then consider the condition

$$(3) \quad e_k(x_1, \dots, x_{k-1}, \cdot) = [T_k(x_1, \dots, x_{k-1})]d_k(x_1, \dots, x_{k-1}, \cdot).$$

Theorem 2. *For each $1 < p < \infty$, there exists a positive constant c_p , depending only on p , such that the following holds. If $(d_k), (e_k)$ and (T_k) be as above satisfying (3), then for all positive integers n we have equation (1).*

We will also need the following intermediate result. For any random variable f , let $f^\#$ be the decreasing rearrangement of $|f|$, that is,

$$f^\#(t) = \sup\{s \in \mathbf{R} : P(|f| < s) < t\}.$$

Theorem 3. *For each $1 < p < \infty$, there exists a positive constant c_p , depending only on p , such that the following holds. Let $(d_k), (e_k)$ be martingale difference sequences on $[0, 1]^N$ with respect to (\mathcal{L}_k) . Suppose that for each positive integer k*

$$\int_0^t (e_k(x_1, \dots, x_{k-1}, \cdot))^{\#}(s) ds \leq \int_0^t (d_k(x_1, \dots, x_{k-1}, \cdot))^{\#}(s) ds$$

for all $t \in [0, 1]$ and almost all x_1, \dots, x_{k-1} . Then for all positive integers n we have equation (1).

2. THE DISCRETE TYPE CASE

In this section we will prove Theorems 1, 2 and 3 in a special discrete situation, which we now describe. For any positive integer N , let Σ be the σ -field generated by the partition $\{[\frac{i-1}{N}, \frac{i}{N}); i = 1, 2, \dots, N\}$. Define a filtration on $[0, 1]^N$ by (\mathcal{F}_k) by $\mathcal{F}_k = \mathcal{L}_{k-1} \times \Sigma$. Suppose $(d_k), (e_k)$ are (\mathcal{F}_k) -adapted. Then for each k and for each x_1, \dots, x_{k-1} , we see that $d_k(x_1, \dots, x_{k-1}, \cdot)$ and $e_k(x_1, \dots, x_{k-1}, \cdot)$ are Σ -measurable simple functions on $[0, 1]$. Therefore d_k and e_k can be written as N -dimensional vectors and $T_k(x_1, \dots, x_{k-1})$ can be represented by a $N \times N$ matrix, that is,

$$(4) \quad \begin{bmatrix} e_k(1) \\ e_k(2) \\ \vdots \\ e_k(N) \end{bmatrix} = \begin{bmatrix} a_k(1, 1), & \dots, & a_k(1, N) \\ a_k(2, 1), & \dots, & a_k(2, N) \\ \vdots & & \vdots \\ a_k(N, 1), & \dots, & a_k(N, N) \end{bmatrix} \begin{bmatrix} d_k(1) \\ d_k(2) \\ \vdots \\ d_k(N) \end{bmatrix}$$

where

$$d_k(i) = d_k(x_1, \dots, x_{k-1}, i) = d_k(x_1, \dots, x_k) \text{ if } x_k \in [\frac{i-1}{N}, \frac{i}{N})$$

$$e_k(i) = e_k(x_1, \dots, x_{k-1}, i) = e_k(x_1, \dots, x_k) \text{ if } x_k \in [\frac{i-1}{N}, \frac{i}{N})$$

$$T_k = T_k(x_1, \dots, x_{k-1}) = [(a_k(x_1, \dots, x_{k-1}))(i, j)]_{N \times N} = [a_k(i, j)]_{N \times N}$$

The condition of being martingale difference sequences implies that

$$\sum_{i=1}^N d_k(i) = \sum_{i=1}^N e_k(i) = 0$$

We will prove Theorem 2 to in this discrete setting.

Theorem 4. *Theorem 2 holds in the case that (d_k) and (e_k) are adapted to the filtration (\mathcal{F}_k) described above.*

In this discrete case, the boundedness of $\|T_k\|_{L_1}$ and $\|T_k\|_{L_\infty}$ by 1 is equivalent to the condition that $\sum_{j=1}^N |a_k(i, j)| \leq 1$ for all i and

$\sum_{i=1}^N |a_k(i, j)| \leq 1$ for all j . We claim that without loss of generality, we can assume that every row sum and column sum of T_k is 0, that is,

$$\sum_{j=1}^N a_k(i, j) = \sum_{i=1}^N a_k(i, j) = 0$$

for all i and j . Suppose the i^{th} row sum $\sum_{j=1}^N a_k(i, j) = R_k(i)$. Let T'_k be the linear operator defined by

$$T'_k = \left[a_k(i, j) - \frac{R_k(i)}{N} \right]_{N \times N}$$

It is clear that every row sum of T'_k is 0 and

$$\begin{aligned} (T'_k d_k)(i) &= \sum_{j=1}^N \left(a_k(i, j) - \frac{R_k(i)}{N} \right) d_k(j) \\ &= \sum_{j=1}^N a_k(i, j) d_k(j) - \frac{R_k(i)}{N} \sum_{j=1}^N d_k(j) \\ &= e_k(i) \end{aligned}$$

Now we can assume that every row sum of T_k is 0. Similarly suppose the j^{th} column sum $\sum_{i=1}^N a_k(i, j) = C_k(j)$. Let T''_k be the linear operator defined by

$$T''_k = \left[a_k(i, j) - \frac{C_k(j)}{N} \right]_{N \times N}$$

Again it is clear that every row sum and column sum of T''_k is 0 and

$$\begin{aligned} (T''_k d_k)(i) &= \sum_{j=1}^N \left(a_k(i, j) - \frac{C_k(j)}{N} \right) d_k(j) \\ &= \sum_{j=1}^N a_k(i, j) d_k(j) - \frac{1}{N} \sum_{j=1}^N C_k(j) d_k(j) \\ &= e_k(i) \end{aligned}$$

since

$$\sum_{i=1}^N e_k(i) = \sum_{j=1}^N C_k(j) d_k(j) = 0$$

After adjusting T_k , it is easy to check that the norms of T_k may be enlarged up to 4. Of course, we can pick up $T_k/4$ instead and absorb the 4 into the constant c_p .

A nonnegative real matrix is said to be *doubly stochastic* if each of its row and column sum is 1. A sub-doubly stochastic matrix means that

each of its row and column sum is less than or equal to 1. Therefore we can change the assumption in Theorem 4 to be that: “for almost all x_1, \dots, x_{k-1} , every row sum and column sum of the matrix from T_k is 0, and the matrix from $|T_k|$ is sub-doubly stochastic for each positive integer k ”

One of the fundamental results in the theory of doubly stochastic matrices was introduced by Birkhoff (see for example [11, p. 117]).

Theorem 5. *If M is a doubly stochastic matrix, then*

$$M = \sum_{i=1}^S \theta_i P_i$$

where P_i are permutation matrices, and the θ_i are nonnegative numbers satisfying $\sum_{i=1}^S \theta_i = 1$.

Lemma 1. *If M is a $n \times n$ sub-doubly stochastic matrix, then there exists a $2n \times 2n$ doubly stochastic matrix such that its upper left $n \times n$ sub-matrix is M .*

Proof. Suppose that $R(i)$ is the i^{th} row sum of M , $C(j)$ is the j^{th} column sum and S is the sum of all entries. Let

$$A = \begin{bmatrix} \frac{1-R(1)}{n}, & \dots, & \frac{1-R(1)}{n} \\ \vdots & & \vdots \\ \frac{1-R(n)}{n}, & \dots, & \frac{1-R(n)}{n} \end{bmatrix}_{n \times n}$$

$$B = \begin{bmatrix} \frac{1-C(1)}{n}, & \dots, & \frac{1-C(n)}{n} \\ \vdots & & \vdots \\ \frac{1-C(1)}{n}, & \dots, & \frac{1-C(n)}{n} \end{bmatrix}_{n \times n}$$

$$C = \text{Diag} \left[\frac{S}{n}, \dots, \frac{S}{n} \right]_{n \times n}$$

Then define

$$M' = \begin{bmatrix} M & A \\ B & C \end{bmatrix}_{2n \times 2n}$$

It is easy to check that M' is a doubly stochastic matrix. \square

Lemma 2. *If M is a sub-doubly stochastic matrix, then there exists a sub-doubly stochastic matrix N such that $M + N$ is doubly stochastic.*

Proof. Let M' be the $2n \times 2n$ doubly stochastic matrix such that its upper left $n \times n$ sub-matrix is M . By Theorem 5,

$$M' = \sum_{i=1}^S \theta_i P'_i$$

where P'_i are $2n \times 2n$ permutation matrices and $\sum_{i=1}^S \theta_i = 1$. Suppose that P_i is the upper left $n \times n$ sub-permutation matrix of P'_i , then

$$M = \sum_{i=1}^S \theta_i P_i$$

Let Q_i be a $n \times n$ sub-permutation matrix such that $P_i + Q_i$ is a permutation matrix, say R_i . Define

$$N = \sum_{i=1}^S \theta_i Q_i$$

thus

$$M + N = \sum_{i=1}^S \theta_i R_i$$

which is a doubly stochastic matrix. \square

Theorem 6. *Let M be an $n \times n$ matrix. If every row sum and column sum of M is 0 and $|M|$ is sub-doubly stochastic, then*

$$M = \sum_{i=1}^S \theta_i P_i$$

where P_i are permutation matrices, $\sum_{i=1}^S \theta_i = 0$ and $\sum_{i=1}^S |\theta_i| = 1$

Proof. Let

$$A = \frac{|M| + M}{2}$$

$$B = \frac{|M| - M}{2}$$

so A and B are nonnegative, and $2A$ and $2B$ are sub-doubly stochastic. By Lemma 2, there exists a sub-doubly stochastic matrix C such that $2(A + C)$ is a doubly stochastic. But A and B have the same row sums and column sums, and hence $2(B + C)$ is also a doubly stochastic. By applying Theorem 5, we have

$$2(A + C) = \sum_{i=1}^m \lambda_i Q_i$$

$$2(B + C) = \sum_{i=1}^{m'} \lambda'_i Q'_i$$

where Q_i, Q'_i are permutation matrices, and the λ_i, λ'_i are nonnegative numbers satisfying $\sum_{i=1}^m \lambda_i = \sum_{i=1}^{m'} \lambda'_i = 1$. Then the result follows because

$$M = (A + C) - (B + C) = \sum_{i=1}^m \frac{\lambda_i}{2} Q_i - \sum_{i=1}^{m'} \frac{\lambda'_i}{2} Q'_i$$

□

Proof of Theorem 4. From Theorem 6, we know that for each $k \geq 1$ and almost all x_1, \dots, x_{k-1}

$$T_k(x_1, \dots, x_{k-1}) = \sum_{i_k=1}^{S_k} \theta_{k,i_k}(x_1, \dots, x_{k-1}) \cdot P_{k,i_k}(x_1, \dots, x_{k-1})$$

where P_{k,i_k} are permutation matrices, $\sum_{i_k=1}^{S_k} \theta_{k,i_k} = 0$, and $\sum_{i_k=1}^{S_k} |\theta_{k,i_k}| = 1$. Let

$$(5) \quad h_{k,i_k}(x_1, \dots, x_{k-1}, \cdot) = [P_{k,i_k}(x_1, \dots, x_{k-1})] d_k(x_1, \dots, x_{k-1}, \cdot).$$

Then

$$\begin{aligned} e_k &= \left[\sum_{i_k=1}^{S_k} \theta_{k,i_k} P_{k,i_k} \right] d_k \\ &= \sum_{i_k=1}^{S_k} |\theta_{k,i_k}| (\varepsilon_{k,i_k} h_{k,i_k}) \end{aligned}$$

where $\varepsilon_{k,i_k} = \text{sgn}(\theta_{k,i_k})$.

Now we need to consider the probability space $\Omega_1 \times \Omega_2$, where $\Omega_1 = \Omega_2 = [0, 1]^{\mathbf{N}}$. We consider all of the previous random variables considered as random variables on this new probability space, depending only upon the first coordinate ω_1 . We define a filtration (\mathcal{G}_k) where $\mathcal{G}_k = \mathcal{F}_k \otimes \mathcal{F}_k$.

We define a sequence of random variables (I_k) so that for each $\omega_1 \in \Omega_1$, the random variable $I_k(\omega_1, \cdot)$ takes the value i with probability $|\theta_{k,i}(\omega_1)|$. Then we see that

$$e_k = E(\epsilon_{k,I_k} h_{k,I_k} | \mathcal{L} \otimes \{\emptyset, \Omega_2\}).$$

Hence, since conditional expectation is a contraction on L_p

$$\left\| \sum_{k=1}^n e_k \right\|_p \leq \left\| \sum_{k=1}^n \epsilon_{k,I_k} h_{k,I_k} \right\|_p.$$

Now we see that (ϵ_{k,I_k}) is a predictable sequence bounded by 1. Hence by Burkholder's inequality, we see that

$$\left\| \sum_{k=1}^n \epsilon_{k,I_k} h_{k,I_k} \right\|_p \leq c_p \left\| \sum_{k=1}^n h_{k,I_k} \right\|_p.$$

Next, observing (5), since P_{k,i_k} are permutation matrices, for each $k \geq 1$, $i_k = 1, 2, \dots, S_k$, h_{k,i_k} is just an x_k -rearrangement of d_k . that is

$$h_{k,i_k}(x_1, \dots, x_{k-1}, j) = d_k(x_1, \dots, x_{k-1}, \pi_{k,i_k}(j))$$

for some permutation π_{k,i_k} . Thus for any sequence (i_k) we have that (h_{k,i_k}) and (d_k) are tangent sequences. But then we see that (h_{k,I_k}) and (d_k) are tangent sequences. Hence there exists a positive constant c_p such that

$$\left\| \sum_{k=1}^n h_{k,I_k} \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p.$$

The result follows. \square

Theorem 7. *Theorem 3 holds in the case that (d_k) and (e_k) are adapted to the filtration (\mathcal{F}_k) described above.*

This will follow immediately from the following well-known result [10, p. 124].

Theorem 8. *$f = (f_1, f_2, \dots, f_N)$, $g = (g_1, g_2, \dots, g_N)$ are N -dimensional real-valued vectors. $f^\# = (f_1^\#, f_2^\#, \dots, f_N^\#)$ is the decreasing rearrangement of $|f| = (|f_1|, |f_2|, \dots, |f_N|)$. Then*

$$\sum_{k=1}^n g_k^\# \leq \sum_{k=1}^n f_k^\#$$

for all $n = 1, 2, \dots, N$ if and only if there exists a matrix $T = [a_{ij}]_{N \times N}$ such that $Tf = g$, $\sum_{i=1}^N |a_{ij}| \leq 1$ and $\sum_{j=1}^N |a_{ij}| \leq 1$.

3. THE GENERAL CASE

The following theorem was proved by Crowe, Zweibel and Rosenbloom [3].

Theorem 9. *Suppose f, g are random variables on $[0, 1]$, then for $1 \leq p \leq \infty$,*

$$\|f^\# - g^\#\|_p \leq \|f - g\|_p$$

Proof of Theorem 3. We will prove this theorem by using the discrete case. For each $k \geq 1$, we approximate d_k and e_k by functions $d'_k(x_k)$ and $e'_k(x_k)$ such that (d'_k) and (e'_k) are adapted to $(\mathcal{L}_{k-1} \times \Sigma)$, keep the martingale property, and

$$\int_0^t (e'_k(x_1, \dots, x_{k-1}, \cdot))^{\#}(s) ds \leq \int_0^t (d'_k(x_1, \dots, x_{k-1}, \cdot))^{\#}(s) ds$$

for all $t \in [0, 1]$. Then we apply Theorem 7.

In what follows, we will fix x_1, \dots, x_{k-1} , and regard the functions below as functions of only one variable x_k on $[0, 1]$. Since we only consider L_p -norm approximation, without loss of generality, we can assume that d_k and e_k are simple functions. Suppose

$$d_k = \sum_{i=1}^S \alpha_{k,i} \chi_{A_{k,i}}$$

$$e_k = \sum_{i=1}^S \beta_{k,i} \chi_{B_{k,i}}$$

For $1 \leq p < \infty$, each $A_{k,i}$, $B_{k,i}$ can be approximated by $A'_{k,i}$, $B'_{k,i}$ which are the finite unions of disjoint intervals $[a, b)$ with rational endpoints such that for any $\gamma > 0$,

$$\mu(\cup_{i=1}^S (A_{k,i} \triangle A'_{k,i})) < \gamma^p \|d_k\|_1^p$$

$$\mu(\cup_{i=1}^S (B_{k,i} \triangle B'_{k,i})) < \gamma^p \|e_k\|_1^p$$

Let N be the least common denominator of these rational endpoints. Hence $[\frac{i-1}{N}, \frac{i}{N})$ is either contained in some $A'_{k,j}$ or disjoint to all $A'_{k,j}$. Let $\alpha'_{k,i} = \alpha_{k,j}$ if $[\frac{i-1}{N}, \frac{i}{N}) \subset A'_{k,j}$ for some j , $\alpha'_{k,i} = 0$ otherwise. $\beta'_{k,i}$ are likewise.

$$\left(\int_{\cup_{i=1}^S (A_{k,i} \cap A'_{k,i})} + \int_{\cup_{i=1}^S (A_{k,i} \triangle A'_{k,i})} \right) \left| \sum_{i=1}^N \alpha'_{k,i} \chi_i - d_k \right|^p \leq \gamma^p \|d_k\|_{\infty}^p \|d_k\|_1^p$$

where $\chi_i = \chi_{[\frac{i-1}{N}, \frac{i}{N})}$.

$$(6) \quad \left\| \sum_{i=1}^N \alpha'_{k,i} \chi_i - d_k \right\|_p \leq \delta \|d_k\|_1$$

where $\delta = \gamma \|d_k\|_{\infty}$. Similarly, for $\beta'_{k,i}$, we have

$$\left\| \sum_{i=1}^N \beta'_{k,i} \chi_i - e_k \right\|_p \leq \epsilon \|e_k\|_1$$

where $\epsilon = \gamma \|e_k\|_\infty$. Since

$$(7) \quad \left(\sum_{i=1}^N \alpha'_{k,i} \chi_i \right)^\# = \sum_{i=1}^N \varepsilon_{\sigma(i)} \alpha'_{k,\sigma(i)} \chi_i$$

for some permutation σ , where $\varepsilon_j = \text{sgn}(\alpha'_{k,j})$. By Theorem 9,

$$(8) \quad \left\| \sum_{i=1}^N \varepsilon_{\sigma(i)} \alpha'_{k,\sigma(i)} \chi_i - d_k^\# \right\|_p < \delta \|d_k\|_1$$

Define

$$(9) \quad \alpha''_{k,i} = N \cdot \int_{\frac{i-1}{N}}^{\frac{i}{N}} d_k^\#$$

Note that

$$(10) \quad \int_0^t \sum_{i=1}^N \alpha''_{k,i} \chi_i = \int_0^t d_k^\#$$

if $t = \frac{i}{N}$ for some $i = 0, 1, 2, \dots, N$.

Since $d_k^\#$ is monotone decreasing, thus Remain integrable, if N is large enough (if not, pick the multiple of N), we can get

$$(11) \quad \left\| \sum_{i=1}^N \alpha''_{k,i} \chi_i - d_k^\# \right\|_p < \delta \|d_k\|_1$$

From (8) and (11),

$$(12) \quad \left\| \sum_{i=1}^N \alpha''_{k,i} \chi_i - \sum_{i=1}^N \varepsilon_{\sigma(i)} \alpha'_{k,\sigma(i)} \chi_i \right\|_p < 2\delta \|d_k\|_1$$

By doing the reverse process of taking decreasing rearrangement of $|\sum_{i=1}^N \alpha'_{k,i} \chi_i|$,

$$(13) \quad \left\| \sum_{i=1}^N \varepsilon_{\sigma^{-1}(i)} \alpha''_{k,\sigma^{-1}(i)} \chi_i - \sum_{i=1}^N \alpha'_{k,i} \chi_i \right\|_p < 2\delta \|d_k\|_1$$

From (6) and (13),

$$(14) \quad \left\| \sum_{i=1}^N \varepsilon_{\sigma^{-1}(i)} \alpha''_{k,\sigma^{-1}(i)} \chi_i - d_k \right\|_p < 3\delta \|d_k\|_1$$

Repeat the same discussion from (7) to (14) for $\beta'_{k,i}$, we can obtain a permutation τ and a sequence $(\beta''_{k,\tau^{-1}(i)})$ such that

$$\left\| \sum_{i=1}^N \varepsilon_{\tau^{-1}(i)} \beta''_{k,\tau^{-1}(i)} \chi_i - e_k \right\|_p < 3\epsilon \|e_k\|_1$$

Here we realize that the integer N should be chosen large enough for approximating both $d_k^\#$ and $e_k^\#$. For shorter notations, we set

$$\left(\varepsilon_{\sigma^{-1}(i)} \alpha''_{k,\sigma^{-1}(i)} \right)_{\varepsilon_{\sigma^{-1}(i)}=1}^N = (\hat{\alpha}_{k,j})_{j=1}^N$$

$$\left(\varepsilon_{\tau^{-1}(i)} \beta''_{k,\tau^{-1}(i)} \right)_{\varepsilon_{\tau^{-1}(i)}=1}^N = (\hat{\beta}_{k,j})_{j=1}^N$$

$$E \left[\sum_{j=1}^N \hat{\alpha}_{k,j} \chi_j \right] = \zeta$$

$$E \left[\sum_{j=1}^N \hat{\beta}_{k,j} \chi_j \right] = \eta$$

For $t = \frac{j}{N}$, $j = 0, 1, 2, \dots, N$, it is clear that

$$\int_0^t \left(\sum_{j=1}^N \hat{\alpha}_{k,j} \chi_j \right)^\# = \int_0^t \sum_{j=1}^N \alpha''_{k,j} \chi_j = \int_0^t d_k^\#,$$

$$\int_0^t \left(\sum_{j=1}^N \hat{\beta}_{k,j} \chi_j \right)^\# = \int_0^t \sum_{j=1}^N \beta''_{k,j} \chi_j = \int_0^t e_k^\#,$$

and $|\zeta| \leq 3\delta\|d_k\|_1$, $|\eta| \leq 3\epsilon\|e_k\|_1$.

$$\begin{aligned}
 (15) \quad & \int_0^t \left(\sum_{j=1}^N (\hat{\beta}_{k,j} - \eta) \chi_j \right)^\# \\
 & \leq \int_0^t \left(\sum_{j=1}^N (|\hat{\beta}_{k,j}| + |\eta|) \chi_j \right)^\# \\
 & \leq \int_0^t \left(\sum_{j=1}^N \hat{\beta}_{k,j} \chi_j \right)^\# + 3\epsilon\|e_k\|_1 \cdot t \\
 & \leq \int_0^t d_k^\# + 3\epsilon\|d_k\|_1 \cdot t \\
 & \leq (1 + 3\epsilon) \int_0^t d_k^\#
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 (16) \quad & \int_0^t \left(\sum_{j=1}^N (\hat{\alpha}_{k,j} - \zeta) \chi_j \right)^\# \geq \int_0^t d_k^\# - 3\delta\|d_k\|_1 \cdot t \\
 & \geq (1 - 3\delta) \int_0^t d_k^\#
 \end{aligned}$$

Finally, we are ready to define d'_k and e'_k . Let

$$\begin{aligned}
 d'_k &= (1 + 3\epsilon) \sum_{j=1}^N (\hat{\alpha}_{k,i} - \zeta) \chi_i \\
 e'_k &= (1 - 3\delta) \sum_{j=1}^N (\hat{\beta}_{k,i} - \eta) \chi_i
 \end{aligned}$$

where $E[d'_k] = E[e'_k] = 0$ is obvious. Combining (15) and (16), we have

$$\int_0^t (e'_k)^\# \leq \int_0^t (d'_k)^\#.$$

By Theorem 7, there exist a positive constant c_p such that

$$\left\| \sum_{k=1}^n e'_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d'_k \right\|_p$$

This is true for almost all x_1, \dots, x_{k-1} , thus it is not hard to check that

$$\left\| \sum_{k=1}^n d_k - \sum_{k=1}^n d'_k \right\|_p \leq \sum_{k=1}^n \|d_k - d'_k\|_p \leq 6\gamma \left(\sum_{k=1}^n \|d_k\|_\infty \|d_k\|_1 \right)$$

$$\left\| \sum_{k=1}^n e_k - \sum_{k=1}^n e'_k \right\|_p \leq \sum_{k=1}^n \|e_k - e'_k\|_p \leq 6\gamma \left(\sum_{k=1}^n \|e_k\|_\infty \|e_k\|_1 \right)$$

Therefore

$$\left\| \sum_{k=1}^n e_k \right\|_p \leq c_p \left\| \sum_{k=1}^n d_k \right\|_p + 12\gamma \left(\sum_{k=1}^n \|d_k\|_\infty \|d_k\|_1 + \sum_{k=1}^n \|e_k\|_\infty \|e_k\|_1 \right)$$

for arbitrary γ . \square

Proof of Theorem 2. If f is a random variable on (Ω, \mathcal{F}, P) , $1 \leq p < \infty$, $0 \leq t \leq 1$, we define the K -functional by

$$K(t, f; L_p, L_\infty) = \inf_{f_0 + f_1 = f} \{ \|f_0\|_p + t \|f_1\|_\infty \}.$$

J. Peetre [13] has shown that

$$K(t, f; L_1, L_\infty) = \int_0^t f^\#(s) ds.$$

Hence it follows that if T is an operator on both $L_1([0, 1])$ and $L_\infty([0, 1])$ with norm bounded by 1, then for $t \geq 0$

$$\int_0^t (Tf)^\#(s) ds \leq \int_0^t f^\#(s) ds.$$

Thus the result follows from Theorem 3. \square

Lemma 3. *Let f and g are real-valued random variables on (Ω, \mathcal{F}, P) . Then*

$$(17) \quad E[M \vee |g|] \leq E[M \vee |f|]$$

for all nonnegative number M if and only if

$$\int_0^t g^\#(s) ds \leq \int_0^t f^\#(s) ds$$

for all $t \in [0, 1]$.

Proof. Equation (17) is equivalent to $E[M \vee g^\#] \leq E[M \vee f^\#]$. For the "if" part, let

$$\alpha = \sup \{ t : f^\#(t) \geq M \}$$

$$\beta = \sup \{ t : g^\#(t) \geq M \}$$

Note that

$$\begin{aligned} E [M \vee f^\#] &= \int_0^\alpha f^\# + (1 - \alpha)M \\ E [M \vee g^\#] &= \int_0^\beta g^\# + (1 - \beta)M \end{aligned}$$

If $\alpha \geq \beta$,

$$\begin{aligned} \int_0^\alpha f^\# + (1 - \alpha)M &= \int_0^\beta f^\# + \int_\beta^\alpha f^\# + (1 - \alpha)M \\ &\geq \int_0^\beta g^\# + \int_\beta^\alpha M + (1 - \alpha)M \\ &= \int_0^\beta g^\# + (1 - \beta)M \end{aligned}$$

If $\alpha < \beta$,

$$\begin{aligned} &\left[\int_0^\alpha f^\# + (1 - \alpha)M \right] - \left[\int_0^\beta g^\# + (1 - \beta)M \right] \\ &= \int_0^\alpha [f^\# - g^\#] - \int_\alpha^\beta g^\# + (\beta - \alpha)M \\ &= \left\{ \int_0^\beta [f^\# - g^\#] - \int_\alpha^\beta [f^\# - g^\#] \right\} - \int_\alpha^\beta g^\# + (\beta - \alpha)M \\ &= \int_0^\beta [f^\# - g^\#] + \left[(\beta - \alpha)M - \int_\alpha^\beta f^\# \right] \geq 0 \end{aligned}$$

Conversely, for any $\alpha \in [0, 1]$, let

$$\begin{aligned} M &= f^\#(\alpha) \\ \beta &= \inf \{ t : g^\#(t) \geq M \} \\ \int_0^\alpha f^\# + (1 - \alpha)M &= E [M \vee f^\#] \\ &\geq E [M \vee g^\#] \\ &= \int_0^\beta g^\# + (1 - \beta)M \\ &\geq \int_0^\alpha g^\# + (1 - \alpha)M \end{aligned}$$

Thus

$$\int_0^\alpha g^\#(s)ds \leq \int_0^\alpha f^\#(s)ds$$

□

Given a random variable f and a sigma field \mathcal{G} , we will say that f is nowhere constant with respect to \mathcal{G} if $P(f = g) = 0$ for every \mathcal{G} measurable function g . The following theorem [12] shows a concrete representation of a sequence of random variables.

Theorem 10. *Let (f_n) be a sequence of random variables taking values in a separable sigma field (S, \mathcal{S}) . Then there exists a sequence of measurable functions $(g_n : [0, 1]^n \rightarrow S)$ that has the same law as (f_n) . If further we have that f_{n+1} is nowhere constant with respect to $\sigma(f_1, \dots, f_n)$ for all $n \geq 0$, then we may suppose that $\sigma(g_1, \dots, g_n) = \mathcal{L}_n$ for all $n \geq 0$.*

Proof of Theorem 1. First we claim that (2) is equivalent to

$$(18) \quad E[(s_k \vee |e_k|)|\mathcal{F}_{k-1}] \leq E[(s_k \vee |d_k|)|\mathcal{F}_{k-1}]$$

For any $A_k \in \mathcal{F}_{k-1}$, $\lambda \geq 0$, $(\lambda \chi_{A_k} \vee s_k)$ is predictable.

$$E[(\lambda \chi_{A_k} \vee s_k) \vee |e_k|] \leq E[(\lambda \chi_{A_k} \vee s_k) \vee |d_k|]$$

When λ intends to infinity,

$$E[(s_k \vee |e_k|)\chi_{A_k^c}] \leq E[(s_k \vee |d_k|)\chi_{A_k^c}]$$

which is equivalent to (18).

Let's consider the map $D_k = (d_k, e_k, f_k) : \Omega \times [0, 1]^N \rightarrow \mathbf{R}^3$ by $(\omega, (x_k)) \mapsto (d_k(\omega), e_k(\omega), x_k)$. It is clear that D_k is nowhere constant with respect to $\sigma(D_1, D_2, \dots, D_{k-1})$. Apply the previous theorem to get $\tilde{D}_k = (\tilde{d}_k, \tilde{e}_k, \tilde{f}_k) : [0, 1]^k \rightarrow \mathbf{R}^3$ such that (\tilde{D}_k) has the same law as (D_k) and $\sigma(\tilde{D}_1, \tilde{D}_2, \dots, \tilde{D}_k) = \mathcal{L}_k$. Let $\phi : [0, 1]^{k-1} \rightarrow [0, \infty)$ be any bounded nonnegative measurable function. There exists a bounded Borel measurable function $\theta : \mathbf{R}^{3(k-1)} \rightarrow \mathbf{R}$ such that $\phi = \theta(\tilde{D}_1, \tilde{D}_2, \dots, \tilde{D}_{k-1})$

almost everywhere in $[0, 1]^{k-1}$. For particular $v_k = M$,

$$\begin{aligned}
\int_{[0,1]^k} (M \vee |\tilde{e}_k|) \cdot \phi &= \int_{[0,1]^k} (M \vee |\tilde{e}_k|) \cdot \theta(\tilde{D}_1, \dots, \tilde{D}_{k-1}) \\
&= \int_{\Omega} (M \vee |e_k|) \cdot \theta(D_1, \dots, D_{k-1}) \\
&= \int_{\Omega} E[(M \vee |e_k|) | \mathcal{F}_{k-1}] \cdot \theta(D_1, \dots, D_{k-1}) \\
&\leq \int_{\Omega} E[(M \vee |d_k|) | \mathcal{F}_{k-1}] \cdot \theta(D_1, \dots, D_{k-1}) \\
&= \int_{\Omega} (M \vee |d_k|) \cdot \theta(D_1, \dots, D_{k-1}) \\
&= \int_{[0,1]^k} (M \vee |\tilde{d}_k|) \cdot \theta(\tilde{D}_1, \dots, \tilde{D}_{k-1}) \\
&= \int_{[0,1]^k} (M \vee |\tilde{d}_k|) \cdot \phi
\end{aligned}$$

Since this is true for all such $\phi \geq 0$, thus $E[(M \vee |\tilde{e}_k|) | \mathcal{L}_{k-1}] \leq E[(M \vee |\tilde{d}_k|) | \mathcal{L}_{k-1}]$. Also if $\phi : [0, 1]^{k-1} \rightarrow \mathbf{R}$ is any bounded measurable function, there exists a bounded Borel measurable function $\theta : \mathbf{R}^{3(k-1)} \rightarrow \mathbf{R}$ such that $\phi = \theta(\tilde{D}_1, \tilde{D}_2, \dots, \tilde{D}_{k-1})$ almost everywhere in $[0, 1]^{k-1}$. Similarly,

$$\int_{[0,1]^k} \tilde{d}_k \cdot \phi = \int_{\Omega} E[d_k | \mathcal{F}_{k-1}] \cdot \theta(D_1, \dots, D_{k-1}) = 0.$$

Thus $E[\tilde{d}_k | \mathcal{L}_{k-1}] = 0$. i.e. (\tilde{d}_k) , (\tilde{e}_k) are martingale difference sequences with respect to (\mathcal{L}_k) and the result follows from Lemma 3 and Theorem 3. \square

Acknowledgments. We would like to mention the help of Jim Reeds and David Boyd in obtaining the argument about doubly stochastic matrices and other useful remarks.

REFERENCES

1. D. L. Burkholder, *Distribution function inequalities for martingales*. Ann. Probability **1** (1973), 19–42.
2. D. L. Burkholder, *A geometrical characterization of banach spaces in which martingale difference sequences are unconditional*. Ann. Probability **9** (1981), 997–1011.
3. L. A. Crowe, J. A. Zweibel and P. C. Rosenbloom, *Rearrangements of functions*, Journal of functional analysis **66** (1986), 43291–438.
4. P. Hitzenko, *Comparison of moments for tangent sequences of random variables*. Probab. Theory Related Fields **78** (1988), 223–230.

5. P. Hitczenko, *On a domination of sums of random variables by sums of conditionally independent ones*. Ann. Probability **22** (1994), 453–468.
6. P. Hitczenko and S. J. Montgomery-Smith, *Tangent sequences in Orlicz and rearrangement invariant spaces*. Math. Proc. Camb. Phil. Soc. **119** (1996), 91–101.
7. P. Kree, *Interpolation d'espaces qui ne sont ni normes, ni complets*. Applications, Seminaire Lions-Schwartz, semestre 1964-1965, Secretariat Mathematique, 11 rue Pierre Curie, Paris 5e.
8. S. Kwapien and W. A. Woyczynski, *Semimartingale integrals via decoupling inequalities and tangent processes*. Case Western Reserve University, preprint, (1986).
9. S. Kwapien and W. A. Woyczynski, *Random series and stochastic integrals. Single and multiple*. Birkhauser, Boston. (1996).
10. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II*. Springer-Verlag. (1979).
11. H. Minc, *Nonnegative matrices*. Wiley Interscience. (1988).
12. S. J. Montgomery-Smith, *Concrete representation of martingales*. Electronic J. Probab. **3**, (1998), paper 15
13. J. Peetre, *Espaces d'interpolation, generalisations, applications*. Rend. Sem. Mat. Fis. Milano **34** (1964), 83-92.
14. J. Zinn, *Comparison of martingale difference sequences*. Probability in Banach spaces, V (Medford, Mass., 1984), 453–457, Lecture Notes in Math., 1153, Springer, Berlin, 1985.

DEPARTMENT OF MATHEMATICS,, UNIVERSITY OF MISSOURI, COLUMBIA, MISSOURI 65211, USA., stephen@math.missouri.edu

DEPARTMENT OF MATHEMATICS,, UNIVERSITY OF MISSOURI, COLUMBIA, MISSOURI 65211, USA., mathgr75@math.missouri.edu