



A. Plasma GFAP
log plasma GFAP (pg/mL)
6.5
6.0
5.5
5.0
4.5
4.0
0
2
4
6
Years from baseline
Non−carrier
APOE .4 carrier
B. Plasma p−tau217 (UPENN−Fujire
log plasma p−tau217 (pg/mL)
1
0
−1
−2
−3
0
2
4
6
8
Years from baseline
Non−carrier
APOE .4 carrier


A. GFAP slope vs ADA
Pearson r = 0.28
ADAS−Cog 13 rate (pts/yr)
GFAP slope (log−units/yr)
B. GFAP slope vs MM
Pearson r = −0.35
MMSE rate (pts/yr)
GFAP slope (log−units/yr)
C. GFAP slope vs HV
Pearson r = −0.07
Hippocampal volume rate (mm³/yr)
GFAP slope (log−units/yr)
D. p−tau217 slope vs A
Pearson r = 0.44
ADAS−Cog 13 rate (pts/yr)
p−tau217 slope (log−units/yr)
E. p−tau217 slope vs I
Pearson r = −0.36
MMSE rate (pts/yr)
p−tau217 slope (log−units/yr)
F. p−tau217 slope vs
Pearson r = −0.32
Hippocampal volume rate (mm³/yr)
p−tau217 slope (log−units/yr)


A. UPENN−Fujirebio vs C2N
Pearson r = 0.82 (n = 489)
C2N PrecivityAD2 (pg/mL)
UPENN−Fujirebio (pg/mL)
0
5
10
15
0.0
0.5
1.0
1.5
2.0
B. UPENN−Fujirebio vs Lilly MSD
Pearson r = 0.94 (n = 33)
Lilly MSD600 (pg/mL)
UPENN−Fujirebio (pg/mL)
0.25
0.50
0.75
1.00
1.25
0.5
1.0
1.5