

Clinical Assessments - Summary

Timepoint:	Baseline	Assigned Sex at Birth:	Female	Date:	2023-01-01	Instrument Version:	Youth (11-17)
		Gender:	Gender fluid Girl/Woman Genderqueer Non-binary	Age:	14	Report Version:	V1; 1-Jan-2022

This dashboard is a summary of research results from participation in the XXX study. These assessments were conducted for research purposes only and are not meant to be interpreted as clinical assessments. If you have any questions, please contact the study team.

The participant has provided consent for us to share these results with you. This report has not been shared with the participant or their family.

Functioning

Sub-Threshold

Threshold

Columbia Impairment Scale (CIS)

Total Score

Diagnostic

Kiddie Schedule for Affective Disorders and Schizophrenia (K-SADS-PL)

Diagnoses determined by diagnostic consensus to be “Not Present” or “No Information” are omitted from the below tables. Age of Onset is in brackets.

Most Severe Past (MSP)

Attention deficit and hyperactivity disorder	Threshold [4]
Social anxiety disorder	Threshold [4]
Generalized anxiety disorder	Threshold [13]
Major depressive episode	Threshold [13]
Obsessive-compulsive disorder	Sub-Threshold [10]

Current (CE)

Attention deficit and hyperactivity disorder	Threshold [4]
Social anxiety disorder	Threshold [4]
Generalized anxiety disorder	Threshold [13]
Obsessive-compulsive disorder	Sub-Threshold [10]

Psychosis

Sub-Threshold

Threshold

Prodromal Questionnaire - Brief (PQ-B)

Total Score

Distress Score

Structured Interview for Prodromal Symptoms (SIPS)

Negative Symptoms Score

Disorganized Symptoms Score

Suicidal Ideation and Behaviours

Sub-Threshold

Borderline

Threshold

Suicidal Ideation Questionnaire (SIQ-Jr)

Total Score

Columbia Suicide Severity Rating Scale (C-SSRS)

	Past 12 Months:	Lifetime:
Passive Suicidal Thoughts/Ideation:	Yes	Yes
Active Suicidal Thoughts/Ideation:	Yes	Yes
Suicidal Behaviour:	Yes	Yes

Non-Suicidal Self Injury (NSSI)

	Past 12 Months:	Lifetime:
Cut or Carved Your Skin:	Yes	Yes
Hit Yourself on Purpose:	No	No
Burnt Your Skin:	No	No
Overdosed on Medication:	No	No

Psychopathology

Borderline

Clinical

ASEBA

Youth Report

Caregiver Report

Anxious-Depressed		
Withdrawn-Depressed		
Somatic Complaints		
Social Problems		
Thought Problems		
Attention Problems		
Rule-Breaking Behavior		
Aggressive Behavior		
Affective Problems		
Anxiety Problems		
Somatic Problems		
ADHD Problems		
Oppositional Defiant		
Conduct Problems		

T-Score

100 50 0 0 50 100

Substance Use

Adolescent Alcohol and Drug Involvement Scale (AADIS)

Alcohol: Endorsed Drug: Endorsed

Total Score

Endorsed Items:

1. Smoking tobacco (cigarettes, cigars), chewing tobacco

2. Alcohol (beer, wine, liquor)

3. Marijuana, hashish, poppers (marijuana with tobacco), K2, spice

4. Ketamine, LSD/acid, mushrooms, peyote, MDA, PCP, salvia, other hallucinogens

6. Powder cocaine (coke, blow)

7. Rock cocaine (crack, rock, freebase)

9. Heroin, other opiates (opium, morphine, oxys, T3s)

## Clinical Assessments - Supplementary Tables

Timepoint:	Baseline	Assigned Sex at Birth:	Female	Date:	2023-01-01	Instrument Version:	Youth (11-17)
		Gender:	Gender fluid Girl/Woman    Genderqueer Non-binary	Age:	14	Report Version:	V1; 1-Jan-2022

This dashboard displays a high level summary of research results. The full assessments are also available upon request. Please contact the study team for more information.

## Suicidal Ideation and Behaviours

Sub-Threshold	
Borderline	
Threshold	

### Suicidal Ideation Questionnaire (SIQ-Jr)




Items that the participant scored responses 5 (couple of times a week) or 6 (almost every day)

1. I thought it would be better if I was not alive
3. I thought about how I would kill myself
4. I thought about when I would kill myself
5. I thought about people dying
6. I thought about death
7. I thought about what to write in a suicide note
8. I thought about writing a will
9. I thought about telling people I plan to kill myself
10. I thought about how people would feel if I killed myself
11. I wished I were dead
12. I thought that killing myself would solve my problems
13. I thought that others would be happier if I was dead

### Non-Suicidal Self Injury (NSSI)

Engaged in NSSI to:




Get rid of bad feelings? (eg. anxiety or guilt)

Feel something, because you were feeling numb or empty?

Communicate with someone else or to get attention?

Get out of doing something or to get away from others?



### Columbia Suicide Severity Rating Scale (C-SSRS)

**Suicidal Thoughts/ Ideation (Past 12 Months):**

1. Wish to be dead
2. Non-specific active suicidal thoughts
3. Active suicidal ideation with any methods (not plan) without intent to act
4. Active suicidal ideation with some intent to act, without specific plan

**Suicidal Thoughts/ Ideation (Lifetime):**

1. Wish to be dead
2. Non-specific active suicidal thoughts
3. Active suicidal ideation with any methods (not plan) without intent to act
4. Active suicidal ideation with some intent to act, without specific plan

**Suicidal Behaviour (Past 12 Months):**

1. Actual attempt

**Suicidal Behaviour (Lifetime):**

- ### 1. Actual attempt

## Functioning

## World Health Organization Disability Assessment Schedule 2.0 (WHODAS)



## Psychopathology

**ASEBA**

Risky Items	Responses
Deliberately harms self or attempt suicide: -----	0 = Not True -----
Sets fires: -----	0 = Not True -----
Talk about killing self/Thinks about killing self:	2 = Very True or Often True