

PDE NOTES

# Notes on Inequalities and Embeddings

*Kiyuob Jung*

(in progress)



Department of Mathematics,  
Kyungpook National University

December 9, 2022

# Contents

<b>1</b>	<b>Definitions and notations</b>	<b>3</b>
1.1	Definitions . . . . .	3
<b>2</b>	<b>Basic properties</b>	<b>4</b>
2.1	Change of variable . . . . .	4
2.2	Coordinates . . . . .	4
2.2.1	Polar coordinates . . . . .	4
<b>3</b>	<b>Inequalities</b>	<b>5</b>
3.1	Power inequalities . . . . .	5
<b>4</b>	<b>Embeddings</b>	<b>6</b>
4.1	Sobolev Embedding . . . . .	6
4.1.1	The case $1 \leq p < n$ . . . . .	6
4.1.2	The case $p = n$ . . . . .	7
4.1.3	The case $n < p \leq \infty$ . . . . .	7
4.1.4	The case $p = \infty$ . . . . .	7
4.1.5	Summary . . . . .	7

## Preface

Todo.

## Abbreviation

Since these notes is not for formal research, we almost always employ the following abbreviations:

◇ s.t. : s.t.

◇ TFAE : The following are equivalent

## Notation

Let a set  $X$ . We employ the following notations:

◇  $\mathbb{R}^n$  :  $n$ -dimensional real Euclidean space,  $\mathbb{R} = \mathbb{R}^1$

◇  $\mathbb{C}^n$  :  $n$ -dimensional complex space,  $\mathbb{C} = \mathbb{C}^1$

◇  $\mathbb{K}$  : either  $\mathbb{R}$  or  $\mathbb{C}$ .

◇  $\partial X$  : boundary of  $X$ .

◇  $\forall$  : for all.

# Chapter 1

## Definitions and notations

In this note, we always denote  $x = (x_1, x_2, \dots, x_n)$  to be a point in  $\mathbb{R}^n$ .

### 1.1 Definitions

Here, we collect definitions for averages of a function.

**Definition 1.1.1.** Let  $f \in L^1(U)$  with an open set  $U \subset \mathbb{R}^n$ .

(a) An average of  $f$  over set  $E$  is

$$\oint_E f dx := \frac{1}{\text{meas}(E)} \int_E f(x) dx.$$

(b) An average of  $f$  over the ball  $B_r(x_0)$  is

$$\oint_{B_r(x_0)} f dx := \frac{1}{\alpha(n)r^n} \int_{B_r(x_0)} f(x) dx.$$

(c) An average of  $f$  over the sphere  $\partial B_r(x_0)$  is

$$\oint_{\partial B_r(x_0)} f dS := \frac{1}{n\alpha(n)r^{n-1}} \int_{\partial B_r(x_0)} f(x) dS.$$

As for integrability, TODO.

**Definition 1.1.2.** Let  $p > 1$  and define  $q \in \mathbb{R}$  by

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Then  $p$  and  $q$  are called *conjugate exponents*.

**Remark 1.1.3.** A simple calculation shows the following:

- (i)  $pq = p + q$ ,
- (ii)  $1 = \frac{p+q}{pq}$ ,
- (iii)  $(p-1)(q-1) = 1$ ,
- (iv)  $q = \frac{p}{p-1}$ .

## Chapter 2

# Basic properties

### 2.1 Change of variable

**Proposition 2.1.1.** *Let  $f : U \rightarrow \mathbb{R}$  with an open set  $U \subset \mathbb{R}^n$ .*

$$(a) \int_{B_r(x_0)} f(ax+b)dx = \frac{1}{r^n} \int_{B_{ar}(x_0+b)} f(x)dx.$$

$$(b) \int_{B_r(x_0)} f(ax+b)dx = \int_{B_{ar}(x_0+b)} f(x)dx.$$

*Proof.* TODO

To show (b), let  $\tilde{x} := ax + b$ . Then  $d\tilde{x} = r^n dx$ . Also, since  $|\tilde{x} - b| = |ax| < |a|r$ , we get  $\tilde{x} \in B_{|a|r}(x_0 + b)$ . Hence,

$$\begin{aligned} \int_{B_r(x_0)} f(ax+b)dx &= \frac{1}{\alpha(n)r^n} \int_{B_r(x_0)} f(x)dx \\ &= \int_{B_{ar}(x_0+b)} f(\tilde{x})d\tilde{x} \\ &= \int_{B_{ar}(x_0+b)} f(\tilde{x})d\tilde{x}. \end{aligned}$$

□

**Remark 2.1.2.** Convolution. TODO

### 2.2 Coordinates

#### 2.2.1 Polar coordinates

**Proposition 2.2.1.** *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be continuous and summable. Then*

$$(a) \int_{\mathbb{R}^n} f(x)dx = \int_0^\infty \left( \int_{\partial B_r(x_0)} f(x)d\mathcal{S} \right) dr \quad \forall x_0 \in \mathbb{R}^n.$$

$$(b) \frac{d}{dr} \left( \int_{B_r(x_0)} f(x)dx \right) = \int_{\partial B_r(x_0)} f(x)d\mathcal{S} \quad \forall r > 0.$$

**Proposition 2.2.2.** *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be continuous and summable. Then*

$$(a) \int_{B_\varepsilon(0)} f(x)dx = \int_0^\varepsilon \left( \int_{\partial B_r(0)} f(x)d\mathcal{S}(x) \right) dr \quad \forall \varepsilon > 0.$$

# Chapter 3

## Inequalities

### 3.1 Power inequalities

**Theorem 3.1.1.** *The following statements holds.*

- (a)  $1 + x \leq e^x \quad \forall x \in \mathbb{R}.$
- (b) (Cauchy's inequality)  
$$xy \leq \frac{x^2}{2} + \frac{y^2}{2} \quad \forall x, y \in \mathbb{R}.$$
- (c)  $e^{(x+y)/2} < \frac{e^y - e^x}{y - x} \quad \forall x, y \in \mathbb{R} \text{ with } x \neq y.$

**Theorem 3.1.2.** *The following statements holds.*

- (a)  $\left(\frac{1}{e}\right)^{\frac{1}{e}} \leq x^x \quad \forall x > 0.$
- (b)  $x \leq x^{x^x} \quad \forall x > 0.$
- (c)  $1 < x^y + y^x \quad \forall x, y > 0.$
- (d)  $x^y + y^x \leq x^x + y^y \quad \forall x, y > 0.$
- (e)  $x^{ey} + y^{ex} \leq x^{ex} + y^{ey} \quad \forall x, y > 0.$
- (f)  $\frac{1 - \frac{1}{x^y}}{y} \leq \ln(x) \leq \frac{x^y - 1}{y} \quad \forall x, y > 0.$  The upper and lower bounds converge to  $\ln(x)$  as  $y \rightarrow 0.$
- (g)  $2 < (x + y)^z + (x + z)^y + (y + z)^x \quad \forall x, y, z > 0.$
- (h)  $x^{2y} + y^{2z} + z^{2x} \leq x^{2x} + y^{2y} + z^{2z} \quad \forall x, y, z > 0.$
- (i)  $(xyz)^{(x+y+z)/3} \leq x^x y^y z^z \quad \forall x, y, z > 0.$

**Theorem 3.1.3.** *The following statements holds.*

- (a) (Cauchy's inequality with  $\varepsilon$ )  
$$xy \leq \varepsilon x^2 + \frac{y^2}{4\varepsilon} \quad \forall x, y > 0, \forall \varepsilon > 0.$$

**Theorem 3.1.4.** *The following statements holds.*

- (a)  $(x + y)^p < x^p + y^p \quad \forall x, y > 0, \forall p \in (0, 1).$
- (b)  $(x + y)^p \leq 2^{p-1} (x^p + y^p) \quad \forall x, y \geq 0, \forall p \in [1, \infty).$

# Chapter 4

## Embeddings

### 4.1 Sobolev Embedding

In this section, we deal with embeddings of diverse Sobolev spaces into others. Given a Sobolev space, it automatically belongs to certain other space, depending on the relationship between the integrability  $p$  and the dimension  $n$ .<sup>1</sup> There are three cases:

$$\begin{aligned} p &\in [1, n), \\ p &= n, \\ p &\in (n, \infty]. \end{aligned}$$

In particular, the second case  $p = n$  is called the *borderline case*. What are we trying to obtain from the Sobolev embedding theory? Broadly speaking, given a Sobolev space  $W^{k,p}$ , imbeddings of  $W^{k,p}$  target two types of Banach spaces: either another Sobolev space  $W^{j,q}$  or Hölder spaces  $C^{j,\alpha}$  for some constants  $j \leq k$ ,  $q \geq p$ , and  $0 \leq \alpha \leq 1$ .

#### 4.1.1 The case $1 \leq p < n$

**Definition 4.1.1.** If  $1 \leq p < n$ , the *Sobolev conjugate*  $p^*$  of  $p$  is defined by

$$p^* := \frac{np}{n-p}.$$

**Remark 4.1.2.** A simple calculation shows the following:

- (i)  $p^* > p$ ,
- (ii)  $\frac{1}{p^*} = \frac{1}{p} - \frac{1}{n}$ ,
- (iii)  $p^* \rightarrow \infty$  as  $p \rightarrow n$ .

**Theorem 4.1.3.** (Gagliardo-Nirenberg-Sobolev inequality) *If  $1 \leq p < n$ , then there exists a constant  $C$ , depending only on  $p$  and  $n$ , such that*

$$\|u\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du\|_{L^p(\mathbb{R}^n)},$$

for all  $u \in C_c^1(\mathbb{R}^n)$ .

**Theorem 4.1.4.** (Estimates for  $W^{1,p}$ ,  $1 \leq p < n$ ) *Let  $U$  be a bounded, open subset of  $\mathbb{R}^n$ , and suppose  $\partial U$  is  $C^1$ . Assume  $1 \leq p < n$  and  $u \in W^{1,p}(U)$ . Then  $u \in L^{p^*}(U)$ , with the estimate*

$$\|u\|_{L^{p^*}(U)} \leq C \|u\|_{W^{1,p}(U)},$$

the constant  $C$  depending only on  $p, n$ , and  $U$ .

---

<sup>1</sup>The regularity of domains also affects the inclusion.

#### 4.1.2 The case $p = n$

#### 4.1.3 The case $n < p \leq \infty$

For convenience as in the notation for the Sobolev conjugate, we write

$$\gamma := 1 - \frac{n}{p},$$

whenever  $n < p \leq \infty$ .

**Theorem 4.1.5.** (Morrey's inequality) *If  $n < p \leq \infty$ , then there exists a constant  $C$ , depending only on  $p$  and  $n$ , such that*

$$\|u\|_{C^{0,\gamma}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}$$

for all  $u \in C^1(\mathbb{R}^n)$ .

**Theorem 4.1.6.** (Estimates for  $W^{1,p}$ ,  $n < p \leq \infty$ , [1]) *Let  $U$  be a bounded, open subset of  $\mathbb{R}^n$ , and suppose  $\partial U$  is  $C^1$ . Assume  $n < p \leq \infty$  and  $u \in W^{1,p}(U)$ . Then  $u$  has a version  $u^* \in C^{0,\gamma}(\bar{U})$  with the estimate*

$$\|u^*\|_{C^{0,\gamma}(\bar{U})} \leq C \|u\|_{W^{1,p}(U)},$$

the constant  $C$  depending only on  $p, n$  and  $U$ .

#### 4.1.4 The case $p = \infty$

**Theorem 4.1.7.** (Characterization of  $W^{1,\infty}$ , [1]) *Let  $U$  be open and bounded, with  $\partial U$  of class  $C^1$ . Then  $u : U \rightarrow \mathbb{R}$  is Lipschitz continuous if and only if  $u \in W^{1,\infty}(U)$*

#### 4.1.5 Summary

**Theorem 4.1.8.** (Sobolev, [2, Theorem 7.29]) *Let  $u \in W_0^{1,p}(B_R(0))$ , where  $B_R(0) \subset \mathbb{R}^n$ . Then there are universal constants  $c, c_1, c_2$  and  $c_3$ , depending on  $n$ , such that*

(a) if  $1 < p < n$  then

$$\|u\|_{L^{p^*}} \leq c \|Du\|_{L^p},$$

(b) if  $p = n$  then

$$\int_{B_R(0)} \exp\left(c_1 \frac{|u|}{\|Du\|_{L^n}}\right)^{\frac{n}{n-1}} dx \leq c_2,$$

(c) if  $p > n$  then

$$\|u\|_{L^\infty} \leq c_3 R^{1-\frac{n}{p}} \|Du\|_{L^p}.$$



# Bibliography

- [1] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [2] M. Giaquinta and L. Martinazzi. *An introduction to the regularity theory for elliptic systems, harmonic maps and minimal graphs*, volume 11 of *Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)]*. Edizioni della Normale, Pisa, second edition, 2012.