EntroDiff: Taming Hyperbolic Shocks via Double-Burgers Coupling

Anonymous Author(s)
Affiliation
Address
email

Abstract

[Placeholder] Abstract to be drafted in W10. The four-sentence skeleton lives in this comment.

1 Introduction

Generative diffusion models in function space have rapidly become a leading paradigm for data-driven PDE solving, yet their successes concentrate on regimes in which the underlying solutions remain smooth. The state of the art—DiffusionPDE [5] for forward and inverse problems under partial observation, and the FunDPS framework [15] that lifts Tweedie's posterior to infinite-dimensional Banach spaces—reports striking accuracy on Darcy, Helmholtz, and viscous Navier–Stokes benchmarks at 3% observation density. Hyperbolic conservation laws sit outside this comfort zone in two reinforcing senses. *Empirically*, none of the five PDE benchmarks of **?**] is hyperbolic, and the Burgers experiment of Huang et al. [5] fixes $\nu_{\mathrm{phys}} = 10^{-2}$ to sidestep the inviscid limit at which entropy admissibility becomes the central question. *Analytically*, prevailing convergence guarantees for diffusion-based PDE solvers measure proximity in L^2, W_2, or TV under classical smoothness assumptions; none accommodates weak solutions, L^1-contraction, or shock geometry. *Structurally*, the PDE-residual guidance term that powers both DiffusionPDE and FunDPS is evaluated by central differences—a discretisation that is ill-defined across discontinuity interfaces, making the guidance velocity itself singular at the very feature that distinguishes hyperbolic solutions from their elliptic counterparts. The shock regime is therefore an empirical, analytical, *and* algorithmic blind spot of generative PDE solving.

A recent identity reframes the landscape: the score field of a VE-SDE diffusion model itself satisfies a viscous Burgers equation along the diffusion-time axis. Concretely, Sarkar [12] establish that, with $s(u,\tau) = \nabla_u \log p_\tau(u)$, the score field obeys $\partial_\tau s + 2\, s \cdot \nabla_u s = \Delta_u s$ (cf. (10))—equivalently, $u \mapsto -2\, s$ is itself a Burgers velocity field in (u,τ). When the target distribution $\rho^\star$ is supported on solutions of a hyperbolic conservation law of the form (2), this score-level Burgers structure runs parallel to a solution-level Burgers structure on (x,t)—and, in fact, the two are *coupled* (Theorem 1): physical shocks define the singular support of the data law and thereby pin the location of the score-level interfacial layer through the correspondence $\Sigma_{\mathrm{phys}}(t) \leftrightarrow \Sigma_{\mathrm{score}}(\tau)$. This coupling is not a mere curiosity. It is the structural lever that allows shock geometry to be promoted from an obstacle of the smooth-data toolbox to an architectural prior on the score network, and—through that prior—to a sharp Wasserstein rate against the Kruzhkov entropy solution.

We turn the double-Burgers correspondence into both an algorithm and a theoretical guarantee; quantitative validation appears in §5.

Contributions.

Submitted to 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Do not distribute.

- **(C1) Double-Burgers coupling (Theorem 1).** We formalise the geometric correspondence between $\Sigma_{\mathrm{phys}}(t)$ and $\Sigma_{\mathrm{score}}(\tau)$ as a coupled Burgers system on (τ, t), providing the structural foundation for the rest of the paper.
- **(C2) BV-aware score parameterization (§3).** We embed the exact $\tanh$ interfacial profile of the score field [12, Proposition 5.4] directly into the network, decomposing s_θ into a smooth potential plus a shock layer modulated by a Rankine–Hugoniot–admissible jump amplitude. This excises the $\exp(\Lambda T_d)$ amplification source identified by Sarkar [12, Theorem 6.3].
- **(C3) Improved Wasserstein rate $W_1 \leq \mathcal{O}(\varepsilon^{1/2})$ (Theorem 3).** Under (C2) and a BV regulariser on the denoiser, we prove that $\mathrm{Law}(u^\theta)$ converges to the Kruzhkov entropy solution $\mathbf{u}^\star$ at rate $\varepsilon^{1/2}$ in the score-matching error, with constants *independent* of the Score-Shocks amplification exponent Λ.

2 Related Work and Background

Function-space diffusion solvers. Diffusion-based generative models have rapidly become the dominant paradigm for data-driven PDE solving in function space. Lim et al. [11] introduced score-based diffusion on Banach space; Huang et al. [5] demonstrated PDE-residual–guided sampling for forward and inverse problems under partial observation; building on these threads, the *FunDPS* framework of Yao et al. [15] extends Tweedie's posterior sampling to infinite-dimensional Banach spaces and reports state-of-the-art accuracy on five PDE benchmarks at 3% observation density. None of these benchmarks is hyperbolic; the Burgers experiment in Huang et al. [5] fixes $\nu_{\mathrm{phys}} = 10^{-2}$, sidestepping the inviscid limit at which entropy admissibility becomes the central question. Their PDE-residual term, evaluated by central differences, is silently undefined at shock interfaces—a structural failure mode that motivates this work.

Flow matching for PDE and function spaces. Stochastic interpolants [1] and their function-space lifts unify flows and diffusions and have fuelled a wave of PDE-oriented variants; recent contributions include the residual-augmented flow-matching operators of Bhola and Duraisamy [3], which map low-fidelity surrogates onto the high-fidelity solution manifold. Our contribution is structurally orthogonal to this line: rather than varying the generative dynamics, we retain the VE-SDE machinery and exploit the *intrinsic* Burgers structure of its score field.

The PDE perspective on diffusion. A complementary line treats diffusion models themselves as PDE flows. Sarkar [12] establishes that the score field of a VE-SDE diffusion model satisfies a viscous Burgers equation in diffusion time, develops a detailed theory of score shocks, Rankine–Hugoniot jumps, and a Lax-type entropy condition for diffusion modes, and identifies an $\exp(\Lambda T_d)$ error-amplification phenomenon (their Theorem 6.3) that is sharp under standard parameterizations. Armegioiu [2] establishes Wasserstein-stability bounds for measure-transport samplers of chaotic PDEs, controlling growth via average-strain exponents rather than worst-case Lipschitz constants. We turn the score–Burgers correspondence from a structural observation into both an architectural prior and a sharp convergence rate (Theorem 3).

Hyperbolic conservation laws as ground truth. Classical numerics—WENO5 [13], Runge–Kutta discontinuous Galerkin schemes [4], and Godunov-type solvers—serve in this paper exclusively as ground-truth generators and as comparators in the entropy-solution sense; we do not propose them as generative baselines. The entropy theory of Kruzhkov [8] and the viscous-approximation rate of Kuznetsov [9] provide the analytical scaffolding for our Theorem 3.

2.1 Background

VE-SDE diffusion preliminaries. We adopt the EDM formulation [7]. The forward process injects Gaussian noise of variance $\sigma^2(\tau)$ at diffusion time $\tau \in [0, T_d]$, yielding the noised density $p_\tau = \rho^\star * G_\tau$ with associated score field $s(u, \tau) = \nabla_u \log p_\tau(u)$. The forward density evolution satisfies the heat equation $\partial_\tau p_\tau = \Delta_u p_\tau$, from which the score-level viscous Burgers structure (10) follows via the Cole–Hopf transformation (see Theorem 1). The reverse-time SDE used at inference reads

$$du = -2\,\dot{\sigma}^2(\tau)\, s(u, \tau)\, d\tau + \sqrt{2\,\dot{\sigma}^2(\tau)}\, d\bar{W}_\tau, \tag{1}$$

and is trained at the network level by minimising the denoising score-matching loss $\mathcal{L}_{\mathrm{DSM}}$ of s_θ. We write u^θ for the sample produced at $\tau = 0$ under the learned reverse process and $\mathrm{Law}(u^\theta)$ for its distribution.

Hyperbolic target and entropy admissibility. The target PDE is the scalar conservation law

$$\partial_t \mathbf{u} + \partial_x f(\mathbf{u}) = 0, \qquad \mathbf{u}(\cdot, 0) = \mathbf{u}_0, \qquad x \in \Omega \subset \mathbb{R}, \quad t \in [0, T_{\mathrm{phys}}], \tag{2}$$

with C^1 flux f (canonical example: Burgers, $f(u) = u^2/2$). Classical solutions may break down in finite time and weak solutions are non-unique; the unique admissible entropy solution $\mathbf{u}^\star$ is selected by the Kruzhkov condition [8], whose semigroup is L^1-contractive—the structural pivot for our convergence proofs in §4.

Wasserstein and BV metrics. We measure the proximity between generated and target distributions through the 1-Wasserstein distance W_1 in Kantorovich–Rubinstein form, and control the relevant function class through the bounded-variation seminorm $\mathrm{TV}(\cdot)$. The compact embedding $\mathrm{BV}(\Omega) \hookrightarrow L^1(\Omega)$ together with Helly's selection theorem provides the relative compactness exploited in §4.

3 Method: *EntroDiff*

Theorem 1 establishes that the score-level and solution-level Burgers structures are geometrically coupled—the physical shock set $\Sigma_{\mathrm{phys}}(t)$ determines the interfacial layer $\Sigma_{\mathrm{score}}(\tau)$ of the score field. In this section we translate this structural coupling into four independent, trainable algorithmic components: (i) a viscosity-matched noise schedule that aligns the viscous profiles of the two Burgers systems (§3.1), (ii) a BV-aware score parameterization that hard-codes the exact $\tanh$ interfacial profile into the network architecture (§3.2), (iii) a composite loss family in which each term addresses a specific theoretical role (§3.3), and (iv) a reverse-time sampler equipped with Godunov-form PDE guidance that remains well-defined across shocks (§3.4). The four components are not additive conveniences; their joint presence is necessary for Theorem 3 to excise the $\exp(\Lambda T_d)$ factor from the convergence bound.

3.1 Viscosity-matched noise schedule

The score-level Burgers equation (10) carries an effective viscosity $g(\tau)^2/2$, where $g(\tau) = \sqrt{\dot{\sigma}^2(\tau)}$ for a VE-SDE. The solution-level conservation law (2), in its vanishing-viscosity (Kruzhkov) limit, is the $\nu_{\mathrm{phys}} \to 0^+$ limit of the viscous regularisation $\partial_t \mathbf{u} + \partial_x f(\mathbf{u}) = \nu_{\mathrm{phys}}\, \partial_{xx} \mathbf{u}$. Theorem 1 achieves its tightest geometric correspondence when the two viscous scales coincide. We therefore replace the default EDM schedule (cosine or polynomial) with a viscosity-matched law:

$$\sigma^2(\tau) = 2\, \nu_{\mathrm{phys}}\, \tau, \qquad \tau \in [0, T_d]. \tag{3}$$

Under (3), the forward SDE becomes pure Brownian rescaling, $du = \sqrt{2\, \nu_{\mathrm{phys}}}\, dw$, and the Fokker–Planck equation degenerates to $\partial_\tau p_\tau = \nu_{\mathrm{phys}}\, \Delta_u p_\tau$. The Cole–Hopf correspondence then yields the score-Burgers equation with viscosity precisely ν_{phys} (cf. the proof of Theorem 1 in Appendix A). This alignment has two consequences: (i) the interfacial layer $\Sigma_{\mathrm{score}}(\tau)$ of the score field develops with the same width scaling $\sim \sqrt{2\, \nu_{\mathrm{phys}}\, \tau}$ as the viscous shock profile of the target PDE, and (ii) the speciation time at which score shocks nucleate coincides with the physical shock-formation time (Theorem 25), enabling Rankine–Hugoniot admissibility to be inherited directly from the score-level dynamics.

3.2 BV-aware score parameterization

The central observation guiding our architecture is that the exact score field in the interfacial layer admits a closed-form $\tanh$ profile. Sarkar [12, Proposition 5.4] show that, for a bimodal target distribution separated by a hyperplane, the score field in the transition layer reduces to

$$s(u, \tau) \;\approx\; \frac{\Delta s_{\mathrm{jump}}}{2}\, \tanh\!\left(\frac{d_\Sigma(u)}{2\, \sigma^2(\tau)}\right) \nabla_u d_\Sigma(u), \tag{4}$$

where d_Σ is the signed distance to the separating manifold and Δs_{jump} is the local jump vector. A standard EDM denoiser must learn this functional form from data; the approximation error in the interfacial region is the source of the $\exp(\Lambda T_d)$ blow-up factor identified by Sarkar [12, Theorem 6.3].

Table 1: Loss family of *EntroDiff*. Each regularizer targets a specific hypothesis in the convergence proof of Theorem 3. **Key take-away**: removing any single term breaks the proof at a distinct step—$\mathcal{L}_{\mathrm{ent}}$ at entropy admissibility, $\mathcal{L}_{\mathrm{BV}}$ at the Helly compactness argument, $\mathcal{L}_{\mathrm{Burg}}$ at the Gronwall estimate.

Loss	Form	Theoretical role
$\mathcal{L}_{\mathrm{DSM}}$	Denoising score matching	Baseline regression target; provides the ε^2 sc
$\lambda_{\mathrm{ent}}\,\mathrm{R}_{\mathrm{ent}}$	Kruzhkov entropy indicator	Selects the physically admissible weak solut
$\lambda_{\mathrm{BV}}\,\mathrm{TV}(D_\theta)$	Total-variation penalty on the denoiser	Guarantees $\hat{u} \in \mathrm{BV}(\Omega)$, enabling the L^1-co
$\lambda_{\mathrm{Burg}}\,\lVert\partial_\tau s_\theta + 2 s_\theta \nabla_u s_\theta - \Delta_u s_\theta\rVert^2$	Score–Burgers consistency residual	Confines s_θ to the score-Burgers manifold (1

We eliminate this source by *embedding the* $\tanh$ *profile directly into the network architecture*. The score network is decomposed into three sub-networks,

$$s_\theta(u,\tau) \;=\; \nabla\phi_\theta^{\mathrm{sm}}(u,\tau) \;+\; \frac{\kappa_\theta(u,\tau)}{2}\;\tanh\!\left(\frac{\phi_\theta^{\mathrm{sh}}(u,\tau)}{2\,\sigma(\tau)^2}\right)\;\nabla\phi_\theta^{\mathrm{sh}}(u,\tau), \tag{5}$$

with the following roles:

- $\phi_\theta^{\mathrm{sm}}(u,\tau)$: a smooth background potential, implemented as a small MLP or U-Net backbone (structurally equivalent to the baseline EDM denoiser); its gradient $\nabla\phi_\theta^{\mathrm{sm}}$ captures the score field in regions far from any shock.
- $\phi_\theta^{\mathrm{sh}}(u,\tau)$: a signed-distance-to-shock function, zero on the interfacial manifold $\Sigma_{\mathrm{score}}(\tau)$ and satisfying $|\nabla\phi_\theta^{\mathrm{sh}}| = 1$ on the manifold; this encodes the shock geometry.
- $\kappa_\theta(u,\tau)$: the local jump amplitude, constrained to satisfy $\kappa_\theta \geq \kappa_0 > 0$ via a Softplus gate; the Rankine–Hugoniot condition of Theorem 25 provides the physical value $\kappa_\theta = f(u_L) - f(u_R)$.

In the limit $\tau \to 0$, the $\tanh$ factor degenerates to a step function, and (5) exactly reproduces the limiting score discontinuity across the shock—without requiring the network to approximate this singular behaviour. This is the architectural lever that degrades the Gronwall constant in Theorem 3 from Λ to $\mathcal{O}(1)$.

3.3 Loss family

The training objective is a composite loss in which each term targets a distinct theoretical requirement of the convergence proof:

$$\mathcal{L} \;=\; \mathcal{L}_{\mathrm{DSM}} \;+\; \lambda_{\mathrm{ent}}\,\mathrm{R}_{\mathrm{ent}}(D_\theta) \;+\; \lambda_{\mathrm{BV}}\,\mathrm{TV}(D_\theta) \;+\; \lambda_{\mathrm{Burg}}\,\big\|\partial_\tau s_\theta + 2\,s_\theta\cdot\nabla_u s_\theta - \Delta_u s_\theta\big\|^2. \tag{6}$$

The four terms, together with their theoretical and empirical roles, are summarised in Table 1.

The Kruzhkov entropy regulariser $\mathrm{R}_{\mathrm{ent}}$ is defined as the expected violation of the entropy inequality over all constant test levels $k \in \mathbb{R}$:

$$\mathrm{R}_{\mathrm{ent}}(D_\theta) \;=\; \mathbb{E}_k\left[\max\!\big\{0,\; \partial_t|D_\theta - k| + \partial_x\big(\mathrm{sgn}(D_\theta - k)\,(f(D_\theta) - f(k))\big)\big\}\right]. \tag{7}$$

The TV term is the standard total-variation seminorm evaluated on the denoiser output; together with the BV-aware parameterization (5), it places the generated samples in a uniformly BV-bounded set $\mathcal{K}_M = \{\mathbf{u} : \mathrm{TV}(\mathbf{u}) \leq M\}$ (Assumption A0 of Theorem 3). The Burgers-consistency term penalises deviations of the learned score from the parabolic manifold (10); when combined with the viscosity-matched schedule (3), this regulariser aligns the score-level dynamics with the viscous approximation of the target PDE.

3.4 Reverse-time sampler with Godunov-form guidance

At inference, we evolve the sample from the noise prior $u_{T_d} \sim \mathcal{N}(0, \sigma^2(T_d)\,I)$ to the data end $\tau = 0$ via a probability-flow ODE with guidance. The key structural difference from DiffusionPDE [5] and FunDPS [15] is the treatment of the PDE residual term: where prior work evaluates $\partial_t u + \partial_x f(u)$ via central differences—a discretisation that is ill-defined at discontinuity interfaces—we employ a

Algorithm 1 *EntroDiff* reverse-time sampling. **Key take-away**: the Godunov flux (9) replaces central differences everywhere a PDE residual is evaluated, making the guidance term well-defined across shock interfaces.

Input: Noise schedule $\sigma(\tau)$, score network s_θ, PDE flux f, guidance weights $\zeta_{\text{obs}}, \zeta_{\text{PDE}}$, steps N_τ
Output: Generated sample u^θ

$\tau \leftarrow T_d, \quad u \leftarrow u_{T_d} \sim \mathcal{N}(0, \sigma^2(T_d)\, I)$
$\Delta\tau \leftarrow T_d / N_\tau$
for $n = 1$ **to** N_τ **do**
  $s \leftarrow s_\theta(u, \tau)$ ▷ BV-aware score evaluation (5)
  $g_{\text{obs}} \leftarrow \zeta_{\text{obs}}\, \nabla_u \mathcal{L}_{\text{obs}}(u)$ ▷ Observation guidance (if applicable)
  $g_{\text{PDE}} \leftarrow \zeta_{\text{PDE}}\, \nabla_u \mathcal{L}_{\text{PDE}}^{\text{Godunov}}(u)$ ▷ Godunov-form PDE guidance (9)
  $v \leftarrow -\,\sigma(\tau)\,\dot\sigma(\tau)\, s \;-\; g_{\text{obs}} \;-\; g_{\text{PDE}}$
  $\tilde u \leftarrow u + \Delta\tau \cdot v$ ▷ Heun predictor
  $\tilde v \leftarrow -\,\sigma(\tau - \Delta\tau)\,\dot\sigma(\tau - \Delta\tau)\, s_\theta(\tilde u, \tau - \Delta\tau) \;-\; \ldots$
  $u \leftarrow u + \frac{\Delta\tau}{2}\,(v + \tilde v)$ ▷ Heun corrector
  $\tau \leftarrow \tau - \Delta\tau$
end for
return $u^\theta \leftarrow u$

Godunov-form flux that is *entropy-consistent* and respects the Rankine–Hugoniot condition across shocks:

$$\frac{du}{d\tau} \;=\; -\sigma(\tau)\,\dot\sigma(\tau)\, s_\theta(u,\tau) \;-\; \zeta_{\text{obs}}\, \nabla_u\, \mathcal{L}_{\text{obs}}(u) \;-\; \zeta_{\text{PDE}}\, \nabla_u\, \mathcal{L}_{\text{PDE}}^{\text{Godunov}}(u). \tag{8}$$

The Godunov residual $\mathcal{L}_{\text{PDE}}^{\text{Godunov}}$ is constructed from the weak form of the conservation law (2): on each spatial cell $[x_{i-1/2},\, x_{i+1/2}]$,

$$\mathcal{L}_i^{\text{Godunov}}(u) \;=\; \frac{\Delta t}{\Delta x}\,\big[f^{\text{God}}(u_i, u_{i+1}) - f^{\text{God}}(u_{i-1}, u_i)\big], \tag{9}$$

where $f^{\text{God}}(u_L, u_R)$ is the Godunov numerical flux—the exact Riemann-solution flux evaluated at the cell interface. This flux is monotone, entropy-satisfying, and well-defined at arbitrary jumps; it seamlessly handles the transition between smooth and shocked regions without oscillation. The guidance coefficients ζ_{obs} and ζ_{PDE} control the strength of the observation and PDE constraints, respectively.

The numerical integration of (8) uses the Heun second-order method with N_τ time steps (default $N_\tau = 50$; full hyperparameters in Appendix C). Algorithm 1 summarises the complete sampling procedure.

Theorem 25 proves that, in the $\tau \to 0$ limit, samples produced by Algorithm 1 satisfy the Rankine–Hugoniot condition and the Lax entropy condition at every shock location. Theorem 29 recasts the entire reverse-time procedure as a JKO gradient flow, establishing a rigorous variational interpretation of the sampling process.

4 Theory

The theorems of this section sequentially erect the convergence theory of *EntroDiff*. §4.1 establishes the structural foundation by coupling the score-level and solution-level Burgers structures (Theorem 1); §4.2 gives a baseline stability bound that exposes the $\exp(\Lambda T_d)$ amplification factor of standard parameterizations; §4.3 states the main contribution—an improved $W_1 \le \mathcal{O}(\varepsilon^{1/2})$ rate that excises this exponent (Theorem 3); §4.4 certifies shock-location admissibility via Rankine–Hugoniot and Lax conditions; and §4.5 closes the loop by recasting the reverse process as a JKO gradient flow.

4.1 Double-Burgers coupling

The point of departure is a structural identity established in Sarkar [12]: along the diffusion-time axis τ, the score field of any VE-SDE diffusion model satisfies a viscous Burgers equation,

$$\partial_\tau s(u,\tau) + 2\, s(u,\tau) \cdot \nabla_u s(u,\tau) = \Delta_u s(u,\tau). \tag{10}$$

Equivalently, $u \mapsto -2\,s$ is a Burgers velocity field in (u, τ). When the target distribution $\rho^\star$ is itself supported on solutions of a hyperbolic conservation law of the form (2), a second Burgers structure emerges in the orthogonal direction: the data law at physical time t, $\rho_t := (\Phi_t^f)_{\#}\rho_0$ with Φ_t^f the entropy semigroup of f, develops shocks at locations $\Sigma_{\mathrm{phys}}(t) \subset \Omega$ in finite time. The two structures are not merely parallel; they are *coupled*, in that physical shocks define the singular support of ρ_t and thereby fix the location of the interfacial layer $\Sigma_{\mathrm{score}}(\tau)$ at which (10) develops a $\tanh$-shaped jump.

Theorem 1 (Double-Burgers coupling)**.** *Let ρ_t denote the law of the Kruzhkov entropy solution of* (2) *at physical time $t \in [0, T_{\mathrm{phys}}]$, and write $p_{\tau,t} := \rho_t * G_\tau$ for the corresponding noised density at diffusion time τ. Setting $s_{\tau,t} := \nabla_u \log p_{\tau,t}$, the joint density $p_{\tau,t}$ satisfies the coupled system*

$$\partial_\tau s_{\tau,t} = \Delta_u s_{\tau,t} + 2\, s_{\tau,t} \cdot \nabla_u s_{\tau,t}, \tag{11}$$

$$\partial_t \rho_t + \partial_x\big(f(\mathbf{u})\,\rho_t\big) = 0 \qquad \textit{(in the entropy-distribution sense).}$$

Moreover, for τ sufficiently small, the interfacial layer $\Sigma_{\mathrm{score}}(\tau)$ of $s_{\tau,t}$ is supported in an $\mathcal{O}(\sigma(\tau))$ neighbourhood of the lift of the physical shock set $\Sigma_{\mathrm{phys}}(t)$ to state space.

Proof sketch. The τ-direction equation follows from the Cole–Hopf transformation applied to the heat equation $\partial_\tau p_{\tau,t} = \Delta_u p_{\tau,t}$ (a restatement of Score Shocks [12, Theorem 4.3] applied pointwise in t). The t-direction equation is the Liouville continuity equation along the characteristic flow of f, lifted to the entropy-solution sense via L^1-contractivity [8]. The geometric correspondence $\Sigma_{\mathrm{phys}}(t) \leftrightarrow \Sigma_{\mathrm{score}}(\tau)$ is obtained from the singular support of ρ_t at physical shocks combined with Laplace's method on the convolution $\rho_t * G_\tau$, yielding the $\mathcal{O}(\sigma(\tau))$ neighbourhood bound. Full proof in Appendix A.1.

4.2 Entropy-solution stability (baseline rate)

Theorem 2 (Entropy-solution stability)**.** *Let the score training error satisfy $\mathbb{E}_\tau \|s_\theta - s_\tau\|^2 \le \varepsilon^2$. Let $\widehat{u}(0)$ be the sample produced by the reverse sampler under the standard (non-BV-aware) parameterization, and $\mathbf{u}^\star$ the Kruzhkov entropy solution of* (2) *at $t = T_{\mathrm{phys}}$. Then*

$$W_1\big(\mathrm{Law}(\widehat{u}(0)), \mathrm{Law}(\mathbf{u}^\star)\big) \le C_1\, \varepsilon\, \exp(\Lambda T_d), \tag{12}$$

where $\Lambda = \sup_\tau \mathrm{SNR}(\tau)/2$ is the Score Shocks amplification exponent [12, Theorem 6.3], and C_1 depends on the schedule bound C_σ and a distributional stability constant C_{stab}.

Proof sketch. The proof proceeds by coupling the true reverse trajectory $u^\natural$ and the learned trajectory $\widehat{u}$ through the ODE flow, then applying Gronwall's inequality in reverse time. The Gronwall constant is $\Lambda = \sup_\tau |\sigma(\tau)\,\dot\sigma(\tau)|\, L_s(\tau)$, where $L_s(\tau)$ is the Lipschitz constant of the true score Jacobian. Under standard parameterizations, $L_s(\tau)$ grows as $1/\sigma^2(\tau)$ near the data end ($\tau \to 0$), leading to the $\exp(\Lambda T_d)$ blow-up. A measure-discrepancy correction via the Kantorovich–Rubinstein duality relates the empirical expectation over $\widehat{u}$ to the true expectation over $u^\natural$. The final W_1 bound then follows from L^1-contractivity of the Kruzhkov entropy semigroup [8], which ensures that numerical discretisation errors in the PDE solution do not propagate. Full proof in Appendix A.2.

4.3 Improved rate via BV-aware parameterization

Theorem 3 (Improved rate via BV-aware parameterization)**.** *Under the BV-aware parameterization* (5) *and the composite loss* (6) *with $\lambda_{\mathrm{BV}} > 0$, and under the regularity assumptions A0–A6 (stated in full in Appendix A.3), we have*

$$W_1\big(\mathrm{Law}(u^\theta), \mathrm{Law}(\mathbf{u}^\star)\big) \le C_2\, \varepsilon^{1/2}, \tag{13}$$

removing the $\exp(\Lambda T_d)$ blow-up factor of Theorem 2. The constant C_2 depends on the BV ball radius M, the viscosity ν_{phys}, and the schedule length T_d, but is independent of Λ and ε.

Proof sketch. The proof proceeds in six stages. **Stage 1** establishes refined trajectory bounds: because the BV-aware parameterization (5) inherits the exact $\tanh$ interfacial profile from Sarkar [12, Proposition 5.4], the score Jacobian's Lipschitz constant degenerates from $L_s(\tau) \sim 1/\tau^2$ to $\mathcal{O}(\tau^{-1/2})$, reducing the Gronwall exponent from Λ to $\mathcal{O}(1)$. **Stage 2** closes the Gronwall loop with this reduced constant, yielding an $\mathcal{O}(\varepsilon)$ trajectory-error bound independent of Λ. **Stage 3** maps the

state-space error to the physical domain through the Kruzhkov L^1-contraction semigroup. **Stage 4** bridges the L^1 and W_1 metrics: the BV penalty $\lambda_{\mathrm{BV}}\mathrm{TV}(D_\theta)$ and the hard-coded $\tanh$ structure jointly guarantee that u^θ belongs to the uniformly bounded-TV ball $\mathcal{K}_M$, on which L^1 and W_1 are equivalent via the Kantorovich–Rubinstein dual. **Stage 5** controls the viscosity error between $\mathbf{u}^\nu$ and $\mathbf{u}^\star$ using the Kuznetsov rate $\mathcal{O}(\nu_{\mathrm{phys}}^{1/2})$ [9], and **Stage 6** assembles all components via the triangle inequality to deliver the $\varepsilon^{1/2}$ rate. Full proof in Appendix A.3.

Remark 4. The rate $\varepsilon^{1/2}$ is consistent with the optimal rate for entropy-solution approximation by vanishing-viscosity methods [9]. Whether the full ε^1 rate is achievable under the present structural assumptions remains an open question; the bottleneck is the $\varepsilon^{1/2}$ conversion from L^2 score error to W_1 distributional distance via Kantorovich–Rubinstein duality, which is tight in the absence of further log-Sobolev or Talagrand-type inequalities on the target distribution.

4.4 Shock-location admissibility

Theorem 5 (Shock-location admissibility)**.** *In the limit $\sigma(\tau) \to 0$, the shock $x_s(t)$ of u^θ satisfies the Rankine–Hugoniot condition*

$$\dot{x}_s(t) = \frac{f(u_L) - f(u_R)}{u_L - u_R}, \tag{14}$$

and the Lax entropy condition $f'(u_L) \geq \dot{x}_s \geq f'(u_R)$ holds almost everywhere.

Proof sketch. The viscosity-matched schedule (3) aligns the effective viscosity of the score-level Burgers equation with the physical viscosity ν_{phys}, establishing an exact interfacial width scaling $\sqrt{2\nu_{\mathrm{phys}}\tau}$ (Lemma 1, Appendix A.9). Under the BV-aware parameterization (5), the $\tanh$ profile degenerates to a Heaviside step in the $\tau \to 0$ limit; the Godunov-form guidance (9) then enforces the weak-form jump condition across every cell interface via a Green's theorem argument on the sample trajectory (Lemma 3). The Rankine–Hugoniot relation—an algebraic consequence of the weak formulation of (2)—is thereby inherited directly from the score-level jump correspondence [12, Theorem 5.5]. The Lax entropy condition follows from Kruzhkov's inequality applied to the Godunov flux, which is monotone and hence entropy-satisfying. Full proof in Appendix A.9.

4.5 JKO correspondence

Theorem 6 (JKO correspondence)**.** *Under the viscosity-matched schedule* (3) *and the Burgers-consistency loss $\mathcal{L}_{\mathrm{Burg}}$, the reverse-time ODE* (8) *in the time-discretization limit $\Delta\tau \to 0$ corresponds to the JKO step*

$$\rho^{n+1} = \arg\min_{\rho \in \mathcal{P}_2} \left\{ \mathcal{F}(\rho) + \frac{1}{2\Delta\tau} W_2^2(\rho, \rho^n) \right\}, \tag{15}$$

where $\mathcal{F}(\rho) = \mathrm{Ent}(\rho \mid \rho^\star) + \lambda\,\mathcal{R}(\rho)$ is the free-energy functional composed of the relative entropy and a PDE-constraint functional $\mathcal{R}$.

Proof sketch. The reverse ODE induces a continuity equation $\partial_\tau p_\tau + \nabla \cdot (p_\tau v) = 0$ on the particle density, with the total velocity field v decomposed into a Fokker–Planck component (from the standard score term) and a guidance component (from $\mathcal{L}_{\mathrm{ent}}$ and $\mathcal{L}_{\mathrm{Burg}}$). The first variation of the free energy $\mathcal{F}$ identifies the gradient-flow velocity as $-\nabla(\delta\mathcal{F}/\delta\rho)$, and the Burgers-consistency penalty $\mathcal{L}_{\mathrm{Burg}}$ ensures that the learned score satisfies the entropy-dissipation relation required for the Wasserstein gradient-flow structure [6]. Under the time reparameterization $\theta(\tau) = \int_0^\tau a(r)\,dr$ with $a(\tau) = \dot{\sigma}^2(\tau)/2$, the density evolution becomes the W_2-gradient flow of $\mathcal{F}$, and the implicit Euler discretisation recovers the JKO variational formulation (15). Full proof in Appendix A.10.

5 Experiments

5.1 Setup

TODO: State that all experiments are toy 1D problems intended to validate the theoretical predictions of Theorems 3 and 25. Ground truth is generated with WENO5 [13]; baselines are EDM [7] and DiffusionPDE [5]; metrics are W_1, L^1, and shock-location error.

Figure 1: **TODO:** Figure 2: Burgers shock comparison. Left: ground-truth WENO5; middle: standard EDM (oscillation visible); right: EntroDiff (clean shock). Key take-away: BV-aware parameterization captures the shock without TV blow-up, and the residual error scales as $\varepsilon^{1/2}$ in line with Theorem 3.

Table 2: **TODO:** Table 2: main result. W_1 error on E1–E3, comparing EDM baseline, DiffusionPDE, FNO, and EntroDiff (ours). Key take-away: EntroDiff achieves the lowest W_1 on every shock-containing benchmark, with the largest gain on Sod (E3).

Method	E1 (Burgers)	E2 (B–L)	E3 (ρ)	E3 (u)
FNO [10]	—	—	—	—
EDM [7]	—	—	—	—
DiffusionPDE [5]	—	—	—	—
EntroDiff (ours)	—	—	—	—

5.2 E1 · Inviscid Burgers

TODO: Show: (a) standard EDM produces oscillations near shock, (b) BV-aware parameterization (5) sharpens the shock without overshoot, (c) the empirical W_1 scaling matches the $\varepsilon^{1/2}$ prediction of Theorem 3 as ε varies (training horizon ablation).

5.3 E2 · Buckley–Leverett

TODO: Non-convex flux $f(u) = u^2/(u^2 + (1-u)^2)$ produces a shock + rarefaction wave. EntroDiff resolves both correctly; we verify Rankine–Hugoniot speed (Theorem 25) numerically.

5.4 E3 · 1D Euler (Sod shock tube)

TODO: Three-component system. Shock + contact discontinuity + rarefaction simultaneously. EntroDiff applied component-wise. Compare W_1 per component.

5.5 Ablation

TODO: Ablate: (a) viscosity-matched schedule vs. EDM default; (b) BV-aware vs. plain parameterization; (c) each loss term in $\mathcal{L}_{\mathrm{ent}}/\mathcal{L}_{\mathrm{BV}}/\mathcal{L}_{\mathrm{Burg}}$. Each ablation should isolate one design choice and link back to the corresponding theorem.

6 Conclusion and Limitations

TODO: Conclusion paragraph: We turned the Score Shocks observation — that a diffusion model's score itself satisfies a Burgers equation — from a purely theoretical curiosity into an algorithmic and theoretical lever for solving hyperbolic PDEs. The double-Burgers structure (Theorem 1) leads naturally to a BV-aware score parameterization, which (Theorem 3) achieves $W_1 \leq \mathcal{O}(\varepsilon^{1/2})$ convergence to the Kruzhkov entropy solution, free of the $\exp(\Lambda)$ blow-up of the standard analysis.

Limitations. **TODO:** Three explicit limitations:

- **Scalar 1D only (theory).** Theorems 3–25 are stated for 1D scalar conservation laws. The system case (Euler) is supported empirically (E3) but a formal extension is left as future work.
- **VE-SDE only.** We work in the variance-exploding setting where Score Shocks gives a clean Burgers identity. The VP / EDM-Karras case is similar in spirit but the algebra is messier; we leave it to a follow-up.
- ϕ^{sh} **in 2D.** Constructing a signed-distance-to-shock function in two dimensions is non-trivial; we use a mesh-based level-set surrogate in the appendix. A learned, mesh-free ϕ^{sh} is an interesting follow-up.

Future work. **TODO:** Three future directions (one sentence each): (i) extension to systems and multi-D; (ii) plasma kinetic equations (Vlasov–Poisson) leveraging the entropy-aware machinery; (iii) connection to flow-matching variants and Hilbert-space diffusion (FunDPS [15]).

References

[1] Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *Journal of Machine Learning Research (JMLR)*, 26, 2025.

[2] Victor Armegioiu. Statistical error bounds for generative solvers of chaotic PDEs: Wasserstein stability, generalization, and turbulence. *arXiv preprint arXiv:2602.18794*, 2026.

[3] Sahil Bhola and Karthik Duraisamy. Flow matching operators for residual-augmented probabilistic learning of partial differential equations. *arXiv preprint arXiv:2512.12749*, 2025.

[4] Bernardo Cockburn and Chi-Wang Shu. Runge–kutta discontinuous Galerkin methods for convection-dominated problems. *Journal of Scientific Computing*, 16(3):173–261, 2001.

[5] Jiahe Huang, Guandao Yang, Zichen Wang, and Jeong Joon Park. DiffusionPDE: Generative PDE-solving under partial observation. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2024.

[6] Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the Fokker–Planck equation. *SIAM Journal on Mathematical Analysis*, 29(1):1–17, 1998.

[7] Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022.

[8] S. N. Kruzhkov. First-order quasilinear equations in several independent variables. *Mathematics of the USSR-Sbornik*, 10(2):217–243, 1970.

[9] N. N. Kuznetsov. Accuracy of some approximate methods for computing the weak solutions of a first-order quasi-linear equation. *USSR Computational Mathematics and Mathematical Physics*, 16(6):105–119, 1976.

[10] Zongyi Li, Nikola B. Kovachki, Kamyar Azizzadenesheli, Burigede Liu, Kaushik Bhattacharya, Andrew M. Stuart, and Anima Anandkumar. Fourier neural operator for parametric partial differential equations. In *International Conference on Learning Representations (ICLR)*, 2021.

[11] Jae Hyun Lim et al. Score-based diffusion models in function space. In *International Conference on Machine Learning (ICML)*, 2023. TODO: confirm full author list at W11.

[12] Krisanu Sarkar. Score shocks: The Burgers equation structure of diffusion generative models. *arXiv preprint arXiv:2604.07404*, 2026.

[13] Chi-Wang Shu. High order weighted essentially non-oscillatory schemes for convection dominated problems. *SIAM Review*, 51(1):82–126, 2009. The canonical WENO reference.

[14] TBA. Deep kinetic JKO schemes for Vlasov–Fokker–Planck equations. 2026. arXiv:2603.23901; TODO: confirm authors at W11.

[15] Jiachen Yao, Abbas Mammadov, Julius Berner, Gavin Kerrigan, Jong Chul Ye, Kamyar Azizzadenesheli, and Anima Anandkumar. Guided diffusion sampling on function spaces with applications to PDEs. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2025.

A Proofs of Main Theorems

A.1 Proof of Theorem 1 (Double-Burgers coupling)

Theorem 7 (Double-Burgers coupling — restated)**.** *Let $\Omega \subset \mathbb{R}$ be the physical domain and consider the 1D scalar conservation law*

$$\partial_t u + \partial_x f(u) = 0, \qquad f \in C^2(\mathbb{R}), \quad u(0,\cdot) = \mathbf{u}_0. \tag{16}$$

*Let S_t denote the Kruzhkov entropy semigroup [**?**]. Let the initial data be random: $\mathbf{u}_0 \sim \rho_0$ with $\rho_0 \in \mathcal{P}(L^\infty(\Omega) \cap \mathrm{BV}(\Omega))$, and define the pushforward distribution at physical time t:*

$$\rho_t \coloneqq (S_t)_\# \rho_0. \tag{17}$$

At each fixed t, apply Gaussian smoothing with diffusion time $\tau > 0$:

$$p_{\tau,t}(u) \coloneqq (\rho_t * G_\tau)(u), \qquad G_\tau(u) = (4\pi\tau)^{-d/2} \exp\!\left(-\frac{\|u\|^2}{4\tau}\right), \tag{18}$$

where $u \in \mathbb{R}^d$ is the state variable (for scalar conservation laws, $d = 1$; the Score Burgers part holds for all $d \geq 1$). Define the score and its Cole–Hopf conjugate:

$$s_{\tau,t}(u) \coloneqq \nabla_u \log p_{\tau,t}(u), \qquad w_{\tau,t}(u) \coloneqq -2\, s_{\tau,t}(u). \tag{19}$$

Then the following hold:

(A) ***Score Burgers (τ-direction).*** *For each fixed t, for all $\tau > 0$ where $p_{\tau,t} > 0$,*

$$\boxed{\partial_\tau w_{\tau,t} + (w_{\tau,t} \cdot \nabla_u) w_{\tau,t} = \Delta_u w_{\tau,t}} \tag{20}$$

with $\nabla \times w_{\tau,t} = 0$ identically (since $w = -2\nabla \log p$).

(B) ***Physical transport (t-direction).*** *The curve $t \mapsto \rho_t$ in $\mathcal{P}_2(\mathbb{R}^d)$ satisfies the continuity equation*

$$\boxed{\partial_t \rho_t + \nabla_u \cdot (\rho_t V_t) = 0, \qquad V_t(u)(x) = -\partial_x f(u(x))} \tag{21}$$

in the sense of distributions, obtained as the vanishing-viscosity limit $\varepsilon \to 0^+$ of the regularized evolution.

(C) ***Shock set co-location.*** *Let*

$$\Sigma_{\mathrm{phys}}(t) \coloneqq \{\, x \in \Omega : u(\cdot,t) \text{ has a jump at } x \,\} \tag{22}$$

be the physical shock set of the entropy solution. For each $\tau > 0$, define the score interfacial layer set

$$\Sigma_{\mathrm{score}}(\tau,t) \coloneqq \{\, u \in \mathbb{R}^d : \|\nabla_u w_{\tau,t}(u)\| \gtrsim 1/\tau \,\}, \tag{23}$$

*i.e., the region where the score gradient is large (the $\tanh$-layer identified by Score Shocks [**?**]). Let $\pi_\Omega : \mathbb{R}^d \to \Omega$ be the projection from state space to physical space (identifying the spatial coordinate of the shock value). Then*

$$\boxed{\pi_\Omega\big(\Sigma_{\mathrm{score}}(\tau,t)\big) \xrightarrow{\tau \to 0^+} \Sigma_{\mathrm{phys}}(t) \qquad \text{in Hausdorff distance.}} \tag{24}$$

Assumptions.

(A1) **Gaussian smoothing.** For each fixed t, ρ_t is a finite Borel measure, and the convolution $p_{\tau,t} = \rho_t * G_\tau$ is strictly positive and C^∞ for all $\tau > 0$.

(A2) **Entropy semigroup.** S_t is the Kruzhkov entropy semigroup, so $\rho_t = (S_t)_\# \rho_0$ is well-defined and $t \mapsto \rho_t$ is continuous in the narrow topology.

(A3) **BV regularity.** ρ_0 is supported on $\mathrm{BV}(\Omega) \cap L^\infty(\Omega)$. Consequently, for each t, $u(\cdot, t) \in \mathrm{BV}(\Omega)$ with $\mathrm{TV}(u(\cdot, t)) \leq \mathrm{TV}(\mathbf{u}_0)$, and the jump set $\Sigma_{\mathrm{phys}}(t)$ is a finite (or countably infinite) set of rectifiable curves in (t, x)-space.

(A4) **Convex flux, for concreteness.** f is uniformly convex ($f'' \geq c > 0$); the non-convex case follows via Oleinik's generalization [**?**] and is discussed in the appendix.

Proof. We establish the three structural identities (A), (B), (C) in order.

Part A — Score Burgers (τ-direction). This part follows directly from the Cole–Hopf correspondence and is essentially a restatement of Score Shocks Theorem 4.3 [**?**].

By definition, $p_{\tau,t} = \rho_t * G_\tau$. Since G_τ is the heat kernel, $\partial_\tau G_\tau = \Delta_u G_\tau$, and differentiation under the integral (justified by A1) yields

$$\partial_\tau p_{\tau,t} = \Delta_u p_{\tau,t}. \tag{25}$$

Set $\varphi \coloneqq \log p_{\tau,t}$ ($p_{\tau,t} > 0$ by A1). Then

$$\partial_\tau \varphi = \frac{\partial_\tau p}{p} = \frac{\Delta p}{p} = \Delta\varphi + |\nabla\varphi|^2, \tag{26}$$

where we used $\nabla \cdot (p\nabla\varphi) = \nabla p \cdot \nabla\varphi + p\Delta\varphi = p|\nabla\varphi|^2 + p\Delta\varphi$ and $\nabla \cdot (p\nabla\varphi) = \nabla \cdot (\nabla p) = \Delta p$, so $\Delta p/p = \Delta\varphi + |\nabla\varphi|^2$.

Now $s_{\tau,t} = \nabla_u \varphi$. Taking the gradient of (26),

$$\partial_\tau s_{\tau,t} = \nabla(\Delta\varphi) + \nabla(|\nabla\varphi|^2) = \Delta s_{\tau,t} + \nabla(|s_{\tau,t}|^2). \tag{27}$$

Since $s_{\tau,t} = \nabla\varphi$ is a gradient, $\nabla s_{\tau,t} = \nabla^2\varphi$ is symmetric: $\partial_{u_j} s_i = \partial_{u_i} s_j$. Hence

$$[\nabla(|s|^2)]_i = \partial_{u_i} \sum_j s_j^2 = 2\sum_j s_j\, \partial_{u_i} s_j = 2\sum_j s_j\, \partial_{u_j} s_i = 2[(s \cdot \nabla)s]_i. \tag{28}$$

Substituting (28) into (27) gives the score PDE:

$$\partial_\tau s_{\tau,t} = \Delta_u s_{\tau,t} + 2(s_{\tau,t} \cdot \nabla_u) s_{\tau,t}. \tag{29}$$

Finally, define $w_{\tau,t} \coloneqq -2\, s_{\tau,t}$. Then

$$\partial_\tau w = -2\, \partial_\tau s = -2\big(\Delta s + 2(s \cdot \nabla)s\big) = \Delta w - 2(w \cdot \nabla)s. \tag{30}$$

Since $s = -w/2$, we have $(w \cdot \nabla)s = -\frac{1}{2}(w \cdot \nabla)w$, hence

$$\partial_\tau w = \Delta w + (w \cdot \nabla)w. \tag{31}$$

Rearranging yields the viscous Burgers equation (20). The curl-free condition $\nabla \times w = -\nabla \times (2\nabla \log p) = 0$ is automatic.

Part B — Physical Transport (t-direction). We prove (21) via the vanishing viscosity method, following the structure sketched in **?**] and adapted to the distributional setting.

B.1 Viscous regularization. Introduce the viscous perturbation of the conservation law:

$$\partial_t u^\varepsilon + \partial_x f(u^\varepsilon) = \varepsilon\, \partial_x^2 u^\varepsilon, \qquad u^\varepsilon(0, \cdot) = \mathbf{u}_0. \tag{32}$$

Let S_t^ε be the solution operator of (32). For BV initial data, the solution $u^\varepsilon(\cdot, t)$ is smooth for all $t > 0$ (parabolic regularization). Define the regularized pushforward distribution

$$\rho_t^\varepsilon \coloneqq (S_t^\varepsilon)_\# \rho_0. \tag{33}$$

By Kruzhkov's theory [**?**], for any $T < \infty$,

$$u^\varepsilon(\cdot, t) \to u(\cdot, t) \quad \text{in } L^1_{\mathrm{loc}}(\Omega), \quad \text{uniformly in } t \in [0, T], \tag{34}$$

as $\varepsilon \to 0^+$, where u is the unique entropy solution. In particular, $\rho_t^\varepsilon \rightharpoonup \rho_t$ in the narrow topology.

B.2 Liouville equation for the viscous flow. For the viscous PDE (32), the solution at each fixed x evolves according to

$$\partial_t u^\varepsilon(t,x) = -\partial_x f(u^\varepsilon(t,x)) + \varepsilon\, \partial_x^2 u^\varepsilon(t,x). \tag{35}$$

Treating this as an ODE in the state variable u for each fixed x, the velocity field is

$$V_t^\varepsilon(u)(x) \coloneqq -\partial_x f(u) + \varepsilon\, \partial_x^2 u. \tag{36}$$

Let $\psi \in C_c^\infty(\mathbb{R}^d)$ be a test function in state space. By the chain rule and the definition $\rho_t^\varepsilon = (S_t^\varepsilon)_\# \rho_0$,

$$\begin{aligned} \frac{d}{dt}\int_{\mathbb{R}^d} \psi(u)\, d\rho_t^\varepsilon(u) &= \mathbb{E}_{\mathbf{u}_0\sim\rho_0}\big[\nabla\psi(u^\varepsilon(t,\cdot))\cdot \partial_t u^\varepsilon(t,\cdot)\big] \\ &= \int_{\mathbb{R}^d} \nabla\psi(u)\cdot V_t^\varepsilon(u)\, d\rho_t^\varepsilon(u) + \varepsilon \int_{\mathbb{R}^d} \Delta\psi(u)\, d\rho_t^\varepsilon(u), \end{aligned} \tag{37}$$

where we used $\mathbb{E}[\nabla\psi\cdot\varepsilon\partial_x^2 u] = -\varepsilon\mathbb{E}[\nabla^2\psi : \partial_x u\otimes\partial_x u + \Delta\psi]$ after an integration by parts in x; the second-order term simplifies to $\varepsilon\Delta_u\psi$ in the state-space measure.

Integration by parts in u yields the weak form

$$\int \psi\, \partial_t \rho_t^\varepsilon\, du = -\int \nabla\psi\cdot V_t^\varepsilon\ \rho_t^\varepsilon\, du + \varepsilon\int \Delta\psi\ \rho_t^\varepsilon\, du, \tag{38}$$

which is the distributional form of

$$\partial_t\rho_t^\varepsilon + \nabla_u\cdot(\rho_t^\varepsilon V_t^\varepsilon) = \varepsilon\Delta_u\rho_t^\varepsilon. \tag{39}$$

B.3 Vanishing viscosity limit. Take $\varepsilon\to 0^+$ in (39). The second-order term $\varepsilon\Delta_u\rho_t^\varepsilon \to 0$ in the sense of distributions. By the L^1-contraction property of the entropy semigroup and BV compactness [**? ?**]:

- $\rho_t^\varepsilon \rightharpoonup \rho_t$ narrowly;
- $V_t^\varepsilon(u)(x) = -\partial_x f(u) + \varepsilon\partial_x^2 u \to -\partial_x f(u) =: V_t(u)(x)$ almost everywhere in (t,x).

Passing to the limit in the weak formulation of (39) gives

$$\partial_t\rho_t + \nabla_u\cdot(\rho_t\, V_t) = 0, \qquad V_t(u)(x) = -\partial_x f(u(x)), \tag{40}$$

in the distributional sense. This is precisely equation (21).

Part C — Shock Set Co-location. This is the core geometric observation of Theorem 1. We sketch the argument; a fully rigorous measure-theoretic proof (including the Hausdorff convergence rate) appears in the supplementary material.

C.1 BV structure of ρ_t. By Assumption A3, $u(\cdot,t)\in \mathrm{BV}(\Omega)$. The Lebesgue decomposition of $u(\cdot,t)$ splits it into an absolutely continuous part and a jump part:

$$u(\cdot,t) = u^{\mathrm{ac}}(\cdot,t) + \sum_{x_s\in\Sigma_{\mathrm{phys}}(t)} [\![u]\!]_{x_s}\, \mathbf{1}_{[x_s,\infty)}, \tag{41}$$

where $[\![u]\!]_{x_s} \coloneqq u(x_s^+,t) - u(x_s^-,t)$ is the jump magnitude at shock x_s.

Correspondingly, the data distribution decomposes as

$$\rho_t = \rho_t^{\mathrm{ac}} + \rho_t^{\mathrm{sing}}, \tag{42}$$

where ρ_t^{sing} is a sum of weighted Dirac masses (or narrow concentrations) at the shock values $\{u(x_s^-,t), u(x_s^+,t)\}_{x_s\in\Sigma_{\mathrm{phys}}(t)}$.

C.2 Mode boundary formation under Gaussian smoothing. Consider a single shock at x_s with left/right limits u_L, u_R. Locally (in state space near u_L, u_R), ρ_t behaves approximately as

$$\rho_t \approx w_L \delta_{u_L} + w_R \delta_{u_R} + \text{smooth background}, \tag{43}$$

where $w_L, w_R > 0$ are the spatial weights on each side of the shock.

Gaussian convolution with G_τ yields

$$p_{\tau,t}(u) \approx w_L G_\tau(u - u_L) + w_R G_\tau(u - u_R) + \text{smooth correction}. \tag{44}$$

This is a two-component Gaussian mixture. By Score Shocks Proposition 3.1 (speciation threshold) [**?**], the score field $s_{\tau,t} = \nabla_u \log p_{\tau,t}$ develops an interfacial layer — a narrow region where $|\nabla_u s|$ is large — located near the "mode boundary," i.e., the set of u where the two Gaussian components have equal weight. As $\tau \to 0^+$, this mode boundary converges to the continuum of values between u_L and u_R.

C.3 Interfacial profile and shock set convergence. By Score Shocks Proposition 5.4 [**?**] (the interfacial profile result), near each mode boundary the score field admits the exact decomposition

$$w_{\tau,t}(u) = w^{\mathrm{sm}}(u) + \frac{\kappa(u)}{2} \tanh\left(\frac{\phi(u)}{2}\right) \nabla\phi(u) + o(1), \tag{45}$$

where $\phi(u)$ is the signed log-ratio of the local mode densities and κ is the interfacial strength (related to the jump magnitude $|u_L - u_R|$). The interfacial layer set is precisely

$$\Sigma_{\mathrm{score}}(\tau, t) = \{\, u : \|\nabla_u w_{\tau,t}(u)\| \gtrsim 1/\tau \,\}, \tag{46}$$

which, to leading order, coincides with the set where $|\nabla\phi|^{-1} \lesssim \sqrt{\tau}$.

Now, the key observation: the mode boundaries of ρ_t — and therefore the interfacial layers of $w_{\tau,t}$ — are in one-to-one correspondence with the shock jumps of $u(\cdot, t)$. This is because:

1. **Each physical shock at x_s contributes a sharp bimodal feature to ρ_t** (the Dirac-like concentrations at u_L, u_R).
2. **Gaussian smoothing of a bimodal feature always produces an interfacial layer** in the score field — this is a deterministic consequence of the heat-kernel/Cole–Hopf dynamics (Score Shocks Theorem 5.5).
3. **Conversely**, smooth (non-shock) regions of $u(\cdot, t)$ contribute only unimodal or slowly varying features to ρ_t, which produce no interfacial layer.

Therefore, projecting $\Sigma_{\mathrm{score}}(\tau, t)$ to physical space via π_Ω (which maps each state value u to the spatial location where that value occurs) yields a set that converges to $\Sigma_{\mathrm{phys}}(t)$ as $\tau \to 0^+$.

C.4 Hausdorff convergence. Formally, for each shock $x_s \in \Sigma_{\mathrm{phys}}(t)$ with jump values u_L, u_R, the interfacial layer in $w_{\tau,t}$ at diffusion time τ has width $\ell_\tau \sim \sqrt{\tau}/|u_L - u_R|$ (Score Shocks Proposition 5.4). Its spatial projection satisfies

$$\mathrm{dist}_H\big(\pi_\Omega(\Sigma_{\mathrm{score}}(\tau, t)),\, \Sigma_{\mathrm{phys}}(t)\big) \le C\,\sqrt{\tau}, \tag{47}$$

where the constant C depends on the shock strength and the local regularity of $u(\cdot, t)$. The proof of this bound relies on the BV structure (to control the number of shocks and their separation) and the $\tanh$-profile asymptotics of Score Shocks. The detailed computation is deferred to the supplementary material. □

Assembling Parts A, B, and C, we have established:

- In the **diffusion time** τ, the score field evolves as a vector viscous Burgers equation (A) — this is the Cole–Hopf structure that underlies all diffusion models.
- In the **physical time** t, the data distribution is transported by the entropy-solution dynamics, satisfying a Liouville equation (B) — this links the PDE to the distributional level.

- The **shock sets** of the physical solution and the score field coincide in the $\tau \to 0^+$ limit (C) — this is the geometric coupling that motivates the EntroDiff architecture: the network's $\tanh$-interfacial-layer structure can be aligned with the known shock locations, enabling the exponential-error-amplification removal of Theorem 3.

Theorem 1 is thus the structural foundation on which the entire EntroDiff methodology is built.

A.2 Proof of Theorem 2 (Entropy-solution stability)

Theorem 8 (Entropy-solution stability — restated)**.** *Let $\mathbf{u}^\star$ be the Kruzhkov entropy solution of the scalar conservation law $\partial_t \mathbf{u} + \partial_x f(\mathbf{u}) = 0$ at a fixed physical time T_{phys}. View the solution as a state vector $u \in \mathbb{R}^d$ ($d = N_x$ is the number of spatial grid points) and define the target distribution*

$$\rho^\star \coloneqq \mathrm{Law}(\mathbf{u}^\star). \tag{48}$$

The forward diffusion process (VE-SDE, schedule $\sigma(\tau)$):

$$p_\tau \coloneqq \rho^\star * G_\tau, \qquad G_\tau(u) = (4\pi\tau)^{-d/2} \exp\left(-\frac{\|u\|^2}{4\tau}\right), \quad \tau \in [0, T_d]. \tag{49}$$

The true score field:

$$s(u,\tau) \coloneqq \nabla_u \log p_\tau(u). \tag{50}$$

The reverse ODE (standard EDM form, τ integrated from T_d to 0):

True trajectory:

$$\frac{du^\natural}{d\tau} = b(u^\natural, \tau), \qquad b(u,\tau) \coloneqq -\sigma(\tau)\,\dot\sigma(\tau)\,s(u,\tau), \quad u^\natural(T_d) \sim p_{T_d}, \tag{51}$$

Network-approximated trajectory:

$$\frac{d\widehat{u}}{d\tau} = \hat{b}(\widehat{u}, \tau), \qquad \hat{b}(u,\tau) \coloneqq -\sigma(\tau)\,\dot\sigma(\tau)\,s_\theta(u,\tau), \quad \widehat{u}(T_d) \sim p_{T_d}. \tag{52}$$

Notation: $u^\natural(\tau)$ is the true-score-driven reverse trajectory; $\widehat{u}(\tau)$ is the learned-score-driven reverse trajectory. $\|\cdot\|$ denotes the Euclidean norm on $\mathbb{R}^d$. $W_1(\mu,\nu)$ is the 1-Wasserstein distance (Kantorovich–Rubinstein dual). The expectation $\mathbb{E}$ is taken over randomness (initial noise + learned score approximation).

Main theorem. *Under Assumptions (A1)–(A5), the generated sample distribution $\mathrm{Law}(\widehat{u}(0))$ and the target distribution $\rho^\star$ satisfy*

$$\boxed{W_1\big(\mathrm{Law}(\widehat{u}(0)),\ \rho^\star\big) \ \le\ C\,\varepsilon\,\exp(\Lambda T_d)}, \tag{53}$$

where $\Lambda \coloneqq \sup_{\tau\in[0,T_d]} |\sigma(\tau)\,\dot\sigma(\tau)|\,L_s(\tau)$, $L_s(\tau)$ is the Lipschitz constant of $\nabla_u s(\cdot,\tau)$, and ε is the score-matching training error bound. The constant $C = C_\sigma\sqrt{T_d}\,(1 + C_{\text{stab}})$ depends only on the schedule upper bound C_σ and the distributional stability constant C_{stab}, and is independent of ε and Λ.

Connection to the PDE solution. *Equation* (53) *provides a quantitative guarantee that the diffusion model's sampled solution distribution is close to the PDE solution distribution. If the training data originates from a numerical solver, the Kruzhkov L^1-contractivity ensures that the numerical error does not amplify during physical time propagation; the bound* (53) *transfers stably to the true PDE solution (see § PDE Connection below).*

Assumptions.

(A1) **Score gradient Lipschitz regularity.** For each $\tau > 0$, the true score field $s(\cdot,\tau)$ is C^1 and its Jacobian's operator norm is uniformly bounded: there exists $L_s(\tau) < \infty$ such that

$$\|\nabla_u s(u,\tau)\|_{\text{op}} \le L_s(\tau), \quad \forall\, u \in \mathbb{R}^d,\ \tau \in (0, T_d]. \tag{54}$$

Define $\Lambda(\tau) := |\sigma(\tau)\,\dot\sigma(\tau)|\,L_s(\tau)$ and set $\Lambda := \sup_{\tau\in[0,T_d]}\Lambda(\tau) < \infty$.

Verifiability: Since $p_\tau = \rho^\star * G_\tau$ and G_τ is the heat kernel, for any $\tau > 0$, $p_\tau \in C^\infty$ and $\nabla_u s = \nabla_u^2 \log p_\tau$ is uniformly bounded. $\rho^\star$ is a pushforward of a BV measure (containing Dirac point masses at shocks); smoothness holds after convolution. $L_s(\tau) \sim 1/\tau$ may diverge as $\tau \to 0$, but we only require $\sup_{\tau\in[\tau_{\rm end},T_d]} L_s(\tau) < \infty$ (early-stopping $\tau_{\rm end} > 0$), consistent with Sarkar [12].

(A2) **Noise schedule boundedness.** There exists $C_\sigma > 0$ such that

$$|\sigma(\tau)\,\dot\sigma(\tau)| \le C_\sigma, \quad \forall\,\tau \in [0, T_d]. \tag{55}$$

Verifiability: For VE-type schedules (e.g., $\sigma^2(\tau) = 2\nu_{\rm phys}\tau$), $\sigma\dot\sigma = \nu_{\rm phys}$ is constant, which directly satisfies this.

(A3) **Score training error (weighted L^2 bound).** The learned score network s_θ satisfies, under the true data distribution p_τ,

$$\int_0^{T_d} \mathbb{E}_{u\sim p_\tau}\big[\,\|s_\theta(u,\tau) - s(u,\tau)\|^2\,\big]\,d\tau \;\le\; \varepsilon^2. \tag{56}$$

Verifiability: This is the training objective of denoising score matching [**?**] (with weight $\sigma(\tau)^2$); ε is determined by the training loss.

(A4) **Distributional stability (Kantorovich–Rubinstein control).** There exists a constant $C_{\rm stab} > 0$ such that for any $\tau \in [0, T_d]$ and any Lipschitz function $f : \mathbb{R}^d \to \mathbb{R}$,

$$|\,\mathbb{E}_{\widehat{u}(\tau)}[f(\widehat{u})] - \mathbb{E}_{u^\natural(\tau)}[f(u^\natural)]\,| \;\le\; C_{\rm stab}\cdot W_1\big(\mathrm{Law}(\widehat{u}(\tau)),\,\mathrm{Law}(u^\natural(\tau))\big)\cdot \mathrm{Lip}(f), \tag{57}$$

where $\mathrm{Lip}(f) := \sup_{u\neq v}|f(u) - f(v)|/\|u - v\|$.

Verifiability: This is a direct generalization of the Kantorovich–Rubinstein dual $|\int f d\mu - \int f d\nu| \le \mathrm{Lip}(f)\cdot W_1(\mu,\nu)$, for which $C_{\rm stab} = 1$. Introducing $C_{\rm stab}$ accommodates additional smoothness offsets introduced by the score network; under compactly supported measures and smooth score assumptions, $C_{\rm stab} = \mathcal{O}(1)$.

(A5) **Network Lipschitz bound.** The learned score network $s_\theta(\cdot,\tau)$ is uniformly Lipschitz across all $\tau \in [0, T_d]$: there exists $L_{s_\theta} < \infty$ independent of τ such that for all $u, v \in \mathbb{R}^d$,

$$\|s_\theta(u,\tau) - s_\theta(v,\tau)\| \le L_{s_\theta}\|u - v\|. \tag{58}$$

Verifiability: Standard MLPs/UNets are Lipschitz on compact input domains; boundedness of weights is ensured by spectral normalization or training regularization.

Proof. The proof proceeds in nine steps. Steps 1–3 set up the synchronous coupling framework; Steps 4–5 address the core technical difficulty (measure mismatch); Steps 6–8 synthesize the Gronwall estimate; Step 9 yields the final W_1 bound.

Step 1 — Synchronous coupling. Define the error process

$$e(\tau) := \widehat{u}(\tau) - u^\natural(\tau), \qquad \tau \in [0, T_d]. \tag{59}$$

Since both trajectories share the same initial distribution p_{T_d}, we have $e(T_d) = 0$ almost surely. Differentiating along the reverse ODE with respect to τ (with τ decreasing from T_d to 0):

$$\frac{d}{d\tau}e(\tau) = \hat b(\widehat{u}(\tau),\tau) - b(u^\natural(\tau),\tau). \tag{60}$$

Add-and-subtract decomposition:

$$\begin{aligned}\frac{d}{d\tau}e(\tau) &= \big[b(\widehat{u}(\tau),\tau) - b(u^\natural(\tau),\tau)\big] \;+\; \big[\hat b(\widehat{u}(\tau),\tau) - b(\widehat{u}(\tau),\tau)\big] \\ &=: \mathrm{(I)} + \mathrm{(II)}.\end{aligned} \tag{61}$$

The two terms differ in nature:

- (I): propagation error due to trajectory deviation $\widehat{u} \neq u^\natural$ under the *same* drift field b;
- (II): model error due to the score approximation $s_\theta \neq s$ in the drift field itself.

Step 2 — Controlling (I) **(propagation error).** From $b(u,\tau) = -\sigma(\tau)\,\dot\sigma(\tau)\,s(u,\tau)$ and Assumption (A1):

$$\begin{aligned}\|b(\widehat{u},\tau) - b(u^\natural,\tau)\| &\le |\sigma(\tau)\,\dot\sigma(\tau)| \cdot \|s(\widehat{u},\tau) - s(u^\natural,\tau)\| \\ &\le |\sigma(\tau)\,\dot\sigma(\tau)| \cdot L_s(\tau) \cdot \|\widehat{u} - u^\natural\| \\ &= \Lambda(\tau)\,\|e(\tau)\| \le \Lambda\,\|e(\tau)\|. \end{aligned} \tag{62}$$

Here $\Lambda = \sup_{\tau\in[0,T_d]} \Lambda(\tau)$ is precisely the Score Shocks amplification exponent defined in Sarkar [12]. This is the **origin** of the $\exp(\Lambda T_d)$ factor — the Lipschitz constant of the drift field b equals Λ, so errors propagate at exponential rate Λ.

Step 3 — Controlling (II) **(score model error).** From the definitions of $b, \hat{b}$ and Assumption (A2):

$$\begin{aligned}\|\hat{b}(\widehat{u},\tau) - b(\widehat{u},\tau)\| &= |\sigma(\tau)\,\dot\sigma(\tau)| \cdot \|s_\theta(\widehat{u},\tau) - s(\widehat{u},\tau)\| \\ &\le C_\sigma\,\|s_\theta(\widehat{u},\tau) - s(\widehat{u},\tau)\|. \end{aligned} \tag{63}$$

Step 4 — Expectation inequality. Take the Euclidean norm and expectation of (61), and define

$$\delta(\tau) \coloneqq \mathbb{E}\,\|e(\tau)\|. \tag{64}$$

By Jensen's inequality $\mathbb{E}\|\frac{d}{d\tau}e\| \ge |\frac{d}{d\tau}\mathbb{E}\|e\||$ (strictly valid under absolute continuity; handled via the chain rule; see Remark 1), combining (62) and (63):

$$\delta'(\tau) \;\le\; \Lambda\,\delta(\tau) \;+\; C_\sigma\,\mathbb{E}_{\widehat{u}(\tau)}\big[\|s_\theta(\widehat{u},\tau) - s(\widehat{u},\tau)\|\big], \tag{65}$$

where $\delta'(\tau) = d\delta/d\tau$ along the reverse time direction ($\tau \searrow 0$).

Remark 9. **(Direction of Jensen's inequality).** Strictly, $\frac{d}{d\tau}\mathbb{E}\|e\| = \mathbb{E}\big[\frac{e}{\|e\|}\cdot\frac{de}{d\tau}\big] \le \mathbb{E}\|\frac{de}{d\tau}\|$, so the inequality direction is correct. On the null set where $\|e(\tau)\| = 0$, the inequality degenerates to $0 \le \mathbb{E}\|\frac{de}{d\tau}\|$, which does not affect the integral bound.

Step 5 — Repairing the measure mismatch (core technical step).

Problem: The second expectation in (65) is taken under the distribution of $\widehat{u}(\tau)$ (the learned trajectory), whereas the training error Assumption (A3) is controlled under p_τ (the distribution of the true score trajectory $u^\natural(\tau)$). The two **measures differ**.

Repair via triangle decomposition. Define

$$g(u,\tau) \coloneqq \|s_\theta(u,\tau) - s(u,\tau)\|. \tag{66}$$

Then

$$\begin{aligned}\mathbb{E}_{\widehat{u}(\tau)}[g(\widehat{u},\tau)] &= \mathbb{E}_{u^\natural(\tau)}[g(u^\natural,\tau)] \;+\; \Big(\mathbb{E}_{\widehat{u}(\tau)}[g(\widehat{u},\tau)] - \mathbb{E}_{u^\natural(\tau)}[g(u^\natural,\tau)]\Big) \\ &\le \mathbb{E}_{u^\natural(\tau)}[g(u^\natural,\tau)] \;+\; |\,\mathbb{E}_{\widehat{u}(\tau)}[g(\widehat{u},\tau)] - \mathbb{E}_{u^\natural(\tau)}[g(u^\natural,\tau)]\,|. \end{aligned} \tag{67}$$

By Assumption (A4) (distributional stability), applied to the function $g(\cdot,\tau)$:

$$|\,\mathbb{E}_{\widehat{u}(\tau)}[g] - \mathbb{E}_{u^\natural(\tau)}[g]\,| \;\le\; C_{\mathrm{stab}} \cdot W_1\big(\mathrm{Law}(\widehat{u}(\tau)),\,\mathrm{Law}(u^\natural(\tau))\big) \cdot \mathrm{Lip}(g). \tag{68}$$

By construction of the synchronous coupling,

$$W_1\big(\mathrm{Law}(\widehat{u}(\tau)), \mathrm{Law}(u^\natural(\tau))\big) \le \mathbb{E}\|\widehat{u}(\tau) - u^\natural(\tau)\| = \delta(\tau). \tag{69}$$

By Assumptions (A5) and (A1), the Lipschitz constant of $g(\cdot,\tau) = \|s_\theta(\cdot,\tau) - s(\cdot,\tau)\|$ satisfies

$$\mathrm{Lip}(g(\cdot,\tau)) \;\le\; L_{s_\theta} + L_s(\tau) \;\le\; L_{s_\theta} + \sup_\tau L_s(\tau) \;=:\; L_g < \infty, \tag{70}$$

where finiteness is guaranteed by early-stopping $\tau_{\text{end}} > 0$ (see the remark following Assumption A1). Combining these:

$$\mathbb{E}_{\widehat{u}(\tau)}[g(\widehat{u},\tau)] \;\le\; \mathbb{E}_{u^\natural(\tau)}[g(u^\natural,\tau)] \;+\; C_{\text{stab}}\, L_g\, \delta(\tau). \tag{71}$$

Define $C' \coloneqq C_{\text{stab}}\, L_g$.

Step 6 — Assembling the differential inequality. Substituting (71) into (65):

$$\boxed{\delta'(\tau) \;\le\; (\Lambda + C')\,\delta(\tau) \;+\; C_\sigma\, \mathbb{E}_{u^\natural(\tau)}\big[\|s_\theta(u^\natural,\tau) - s(u^\natural,\tau)\|\big]}. \tag{72}$$

This is a standard first-order linear differential inequality (Gronwall type), where the distribution of $u^\natural(\tau)$ is precisely p_τ, matching the measure in the training error Assumption (A3) after Step 5's repair.

Step 7 — Score error integral bound (Cauchy–Schwarz). Define

$$\beta(\tau) \coloneqq \mathbb{E}_{u^\natural(\tau)}\big[\|s_\theta(u^\natural,\tau) - s(u^\natural,\tau)\|\big]. \tag{73}$$

By Cauchy–Schwarz:

$$\begin{aligned}\int_0^{T_d} \beta(\tau)\, d\tau &\le \sqrt{T_d}\, \left(\int_0^{T_d} \mathbb{E}_{u^\natural(\tau)}\big[\|s_\theta - s\|^2\big]\, d\tau\right)^{1/2} \\ &\le \sqrt{T_d}\, \varepsilon,\end{aligned} \tag{74}$$

where the last step uses Assumption (A3). The key point: the expectation in (A3) is taken under the **true trajectory distribution** $p_\tau = \mathrm{Law}(u^\natural(\tau))$, which matches the measure in (72) after the Step 5 repair.

Step 8 — Gronwall reverse-time integration. Apply the Gronwall argument to the differential inequality (72) over reverse time $[T_d, 0]$. Set $\bar\Lambda \coloneqq \Lambda + C'$. Rewrite (72) as

$$\frac{d}{d\tau}\big[\delta(\tau)\, e^{-\bar\Lambda\tau}\big] \;\le\; C_\sigma\, \beta(\tau)\, e^{-\bar\Lambda\tau}, \tag{75}$$

where the derivative is along the reverse direction (τ decreasing). Integrating forward over $[0, T_d]$ (equivalently: writing the reverse Gronwall in forward form):

$$\delta(0)\, e^{-\bar\Lambda\cdot 0} - \delta(T_d)\, e^{-\bar\Lambda T_d} \;\le\; C_\sigma \int_0^{T_d} \beta(\tau)\, e^{-\bar\Lambda\tau}\, d\tau. \tag{76}$$

Since the initial conditions coincide at the noise endpoint ($\widehat{u}(T_d) \sim p_{T_d}$, $u^\natural(T_d) \sim p_{T_d}$), we have $\delta(T_d) = 0$. Hence

$$\begin{aligned}\delta(0) &\le C_\sigma\, e^{\bar\Lambda T_d} \int_0^{T_d} \beta(\tau)\, e^{-\bar\Lambda\tau}\, d\tau \\ &\le C_\sigma\, e^{\bar\Lambda T_d} \int_0^{T_d} \beta(\tau)\, d\tau.\end{aligned} \tag{77}$$

Substituting (74):

$$\delta(0) \;\le\; C_\sigma \sqrt{T_d}\, \varepsilon\, e^{(\Lambda + C')T_d}. \tag{78}$$

Step 9 — W_1 upper bound and absorption of constants. By the synchronous coupling construction, the W_1 distance is bounded above by the L^1 expectation under the coupling:

$$W_1\big(\mathrm{Law}(\widehat{u}(0)),\, \rho^\star\big) \;\le\; \mathbb{E}\|\widehat{u}(0) - u^\natural(0)\| \;=\; \delta(0). \tag{79}$$

Note: strictly, W_1 above couples $\mathrm{Law}(\widehat{u}(0))$ vs. $\mathrm{Law}(u^\natural(0))$, and $\mathrm{Law}(u^\natural(0)) = \rho^\star$ (since the true reverse ODE exactly recovers the target distribution). Therefore $W_1(\mathrm{Law}(\widehat{u}(0)), \rho^\star) \le \delta(0)$ holds.

Substituting (78) and setting $C := C_\sigma \sqrt{T_d}\, e^{C' T_d}$:

$$W_1\big(\mathrm{Law}(\widehat{u}(0)),\, \rho^\star\big) \;\le\; C\, \varepsilon\, e^{\Lambda T_d}. \tag{80}$$

Observe that $C' = C_{\mathrm{stab}}\, L_g$, where L_g depends on the schedule and network structure but is independent of ε. In the asymptotic regime $\varepsilon \ll 1$, $e^{C' T_d} = \mathcal{O}(1)$ and can be absorbed into the constant C. The final form is as stated in (53). □

PDE Connection — Stable transfer from numerical solution to true PDE solution. The above proof is carried out entirely within the diffusion model domain (state space $\mathbb{R}^d$) and yields a W_1 bound for $\mathrm{Law}(\widehat{u}(0))$ approximating $\rho^\star = \mathrm{Law}(\mathbf{u}^\star)$. The following ensures that this bound **stably transfers** to the true PDE solution when the training data originates from a numerical solver, without amplification during physical time propagation.

Proposition 10 (Kruzhkov L^1-contractivity)**.** *Let $\mathbf{u}, \mathbf{v}$ be two Kruzhkov entropy solutions of the same scalar conservation law $\partial_t \mathbf{u} + \partial_x f(\mathbf{u}) = 0$ with initial data $\mathbf{u}_0, \mathbf{v}_0$ respectively. Then at any physical time $t > 0$,*

$$\|\mathbf{u}(\cdot, t) - \mathbf{v}(\cdot, t)\|_{L^1(\Omega)} \;\le\; \|\mathbf{u}_0 - \mathbf{v}_0\|_{L^1(\Omega)}. \tag{81}$$

*(See **?**], Theorem 1, or **?**], § 6.3.)*

Corollary 11 (Numerical error does not amplify)**.** *If the training data is generated by a numerical solution $\mathbf{u}_h$ satisfying $\|\mathbf{u}_h - \mathbf{u}^\star\|_{L^1} \le \delta_{\mathrm{num}}$, then under the discrete correspondence $\|u\| \approx \sqrt{\Delta x}\, \|\mathbf{u}\|_{L^2}$,*

$$\|\widehat{u}(0) - u_h\| \;\le\; \underbrace{\|\widehat{u}(0) - u^\star\|}_{\text{Theorem 2 bound } \mathcal{O}(\varepsilon e^{\Lambda T_d})} + \underbrace{\|u^\star - u_h\|}_{\text{numerical error } \mathcal{O}(\delta_{\mathrm{num}})}. \tag{82}$$

By (81), δ_{num} does not grow with physical time T_{phys} — this is a core property distinguishing conservation law entropy solutions from general PDEs (e.g., Navier–Stokes L^2 errors may propagate in time, whereas conservation law L^1 errors are bounded by the initial data).

Practical implication: In experiments, high-resolution Godunov/WENO solvers are used to generate training reference solutions; δ_{num} is much smaller than the score-matching error ε (typical grid resolution $N_x \ge 2048$ yields $\delta_{\mathrm{num}} \sim 10^{-4}$), so the Theorem 2 bound is the dominant error term.

Remarks on the form and limitations of the bound.

Remark 12 (Origin and tightness of $\exp(\Lambda)$). The exponential factor $\exp(\Lambda T_d)$ in (53) originates from the Lipschitz constant $\Lambda = \sup_\tau |\sigma\dot{\sigma}|\, L_s(\tau)$ of the drift field b. When the target distribution contains shocks (i.e., $\rho^\star$ has Dirac point masses), $\nabla_u s$ blows up as $\mathcal{O}(1/\tau)$ in the interfacial layer, leading to $L_s(\tau) \sim 1/\tau$ and hence Λ diverges (or becomes large, depending on the early-stopping τ_{end}). This exponential amplification is the **inherent deficiency** of standard score-based diffusion when handling discontinuous solutions.

Theorem 3 reduces this amplification factor to $\mathcal{O}(1)$ via BV-aware parameterization, constituting the core contribution of this paper.

Remark 13 (Relation to Score Shocks [12]). The bound structure of this theorem is consistent with Score Shocks Theorem 6.3 ($W_1 \lesssim \varepsilon \exp(\Lambda T)$). Our independent derivation supplements the following technical points not detailed in Score Shocks:

- Formal treatment of the measure mismatch (Step 5, Assumption A4);
- Explicit tracing of the Lipschitz constant (Assumption A1);
- Bridging to Kruzhkov L^1-contractivity, enabling the bound to transfer from the diffusion model domain to the PDE domain.

Remark 14 (The constant C' and the network Lipschitz bound). The auxiliary constant $C' = C_{\mathrm{stab}}\, L_g$ that appears in Steps 5–6 accumulates as $e^{C' T_d}$ in the exponent. At practical scales ($T_d \sim 1$, L_g controlled by spectral normalization or weight decay), $e^{C' T_d} = \mathcal{O}(1)$, so this constant does not affect the asymptotic rate but only the absolute constant C. Formally, one may require the network to satisfy $\|s_\theta\|_{\mathrm{Lip}} \le L_{\mathrm{net}}$ as part of training regularization.

A.3 Proof of Theorem 3 (Improved rate)

Theorem 15. *Under Assumptions A.4–A.4, consider the BV-aware parameterization* (5) *and the loss $\mathcal{L}_3$ (including the TV regularizer). Let the score matching error satisfy A3, with $\varepsilon \in (0,1)$. Then the generated distribution $\mu_\theta = \mathrm{Law}(u^\theta)$ and the Dirac measure $\delta_{\mathbf{u}^\star}$ at the Kruzhkov entropy solution $\mathbf{u}^\star$ satisfy*

$$W_1(\mu_\theta, \delta_{\mathbf{u}^\star}) \le C_{\mathrm{final}}\, \varepsilon^{1/2}, \tag{83}$$

where C_{final} depends only on $M, T_d, T_{\mathrm{phys}}, \kappa_0, L_{\mathrm{sm}}, \nu_{\mathrm{phys}}/\varepsilon$ (all independent of ε). In particular, the bound does not *contain the exponential amplification factor* $\exp(\Lambda T)$ *(cf. Theorem 2).*

A.4 Assumption System (A0–A6)

The following assumptions hold throughout the proof. The physical meaning and verifiability of each are stated alongside.

Assumption A0 (BV boundedness) The TV regularizer $\lambda_{\mathrm{BV}}\,\mathrm{TV}(D_\theta)$ in the loss $\mathcal{L}_3$ together with the hard-coded $\tanh$ structure of parameterization (5) jointly guarantee that the generated sample u^θ almost surely belongs to the bounded TV ball

$$\mathcal{K}_M \coloneqq \big\{\mathbf{u} \in L^1(\Omega) \cap L^\infty(\Omega) : \mathrm{TV}(\mathbf{u}) \le M\big\}, \tag{84}$$

and the true entropy solution satisfies $\mathrm{TV}(\mathbf{u}^\star) \le M$, with $M > 0$ a constant independent of ε. *Verifiability:* M can be estimated from the TV bound of the initial data and the TV-non-increasing property of the BV semigroup; λ_{BV} is a training hyperparameter.

Assumption A1 (Locally bimodal score structure) In a neighborhood of the shock manifold $\Sigma_{\mathrm{score}}(\tau)$, the true score field admits the asymptotic form

$$s(u,\tau) = s^{\mathrm{sm}}(u,\tau) + \frac{\Delta s_{\mathrm{jump}}(u)}{2}\tanh\Big(\frac{A(u,\tau) - B(u,\tau)}{2\tau}\Big) + \mathcal{O}(\tau), \tag{85}$$

where A, B are the Cole–Hopf potential functions of the two dominant modes, $\Delta s_{\mathrm{jump}} = \nabla_u B - \nabla_u A$ is the jump vector, and s^{sm} is uniformly Lipschitz on the whole space. This structure follows as a local result of the Laplace asymptotic method, assuming sufficiently small diffusion time and well-separated Gaussian clusters; superposition of finitely many such local structures is allowed. *Verifiability:* Verifiable for discrete distribution mixture models; treated as a reasonable engineering approximation for general continuous distributions (see engineering approximation EA1).

Assumption A2 (Network parameterization constraint) In the parameterization (5):

- $\kappa_\theta(u,\tau) \ge \kappa_0 > 0$ holds strictly for all (u,τ) (enforceable via Softplus activation, i.e., $\kappa_\theta = \kappa_0 + \mathrm{Softplus}(\tilde{\kappa}_\theta)$);
- The maps $\nabla_u^2 \phi_\theta^{\mathrm{sm}}$, $\nabla_u^2 D_\theta$, $\nabla_u \kappa_\theta$ are uniformly bounded on $(u,\tau) \in \mathbb{R}^d \times [\tau_{\mathrm{end}}, T_d]$, with bounds independent of τ.

Verifiability: Enforceable at training time via network architecture design and weight norm constraints.

Assumption A3 (Score matching error) There exists $\varepsilon > 0$ such that in the grid-independent mean-squared error sense:

$$\mathbb{E}_{u,\tau}\Big[\frac{|\Omega|}{d}\sum_{i=1}^{d} e_i(u,\tau)^2\Big] \le \varepsilon^2, \tag{86}$$

where $\Delta x = |\Omega|/d$; equivalently $\mathbb{E}_{u,\tau}\big[\Delta x \|e(u,\tau)\|^2\big] \le \varepsilon^2$, with expectation taken over the training distribution $p(u,\tau)$ ($u \sim p(\cdot,\tau)$, $\tau \sim \mathcal{U}[\tau_{\mathrm{end}}, T_d]$). This setting makes ε^2 a physical approximation of the continuous $\int_\Omega e(x,\tau)^2\,dx$.

Assumption A4 (Prior matching) The squared distance bound of the prior approximation at the initial time $\tau = T_d$ of the reverse ODE is controlled by the same scaling:

$$\mathbb{E}\big[\Delta x \|\widehat{u}(T_d) - u^\natural(T_d)\|^2\big] \le \tilde{C}^2 \varepsilon^2, \tag{87}$$

where $\tilde{C} > 0$ is a constant independent of ε and d. *Typically satisfied when:* p_{T_d} is standard Gaussian and T_d is large enough that the signal-to-noise ratio is very low.

Assumption A5 (Early stopping and viscosity matching balance, engineering approximation EA2) The reverse ODE is early-stopped at $\tau_{\text{end}} > 0$, the effective terminal viscosity is defined by $\sigma(\tau)^2(\tau_{\text{end}}) = 2\nu_{\text{phys}}\tau_{\text{end}}$, and the following holds:

$$\nu_{\text{phys}} = \Theta(\varepsilon), \quad \text{i.e.,} \quad c\varepsilon \leq \nu_{\text{phys}} \leq C\varepsilon, \tag{88}$$

with A3 still valid over the full interval $[\tau_{\text{end}}, T_d]$. *Engineering approximation note:* The achievability of $\nu_{\text{phys}} = \Theta(\varepsilon)$ depends on fine design of the noise schedule and prior estimation of the training error ε; this is an engineering approximation (EA2), whose scope of influence is discussed in Appendix C.

Assumption A6 (Kuznetsov-type error bound) Under A0 and A5, the L^1 error between the viscous PDE solution $\mathbf{u}^\nu$ and the Kruzhkov entropy solution $\mathbf{u}^\star$ satisfies

$$\|\mathbf{u}^\nu - \mathbf{u}^\star\|_{L^1(\Omega)} \leq C_K\sqrt{\nu_{\text{phys}}}, \tag{89}$$

where C_K depends on the initial TV bound M and physical time T_{phys}, but is independent of ε. *Explanation:* This bound follows directly from the classical theorem of Kuznetsov (1976) under A0, with $C_K = C\sqrt{T_{\text{phys}}} \cdot \text{TV}(\mathbf{u}_0) \leq C\sqrt{T_{\text{phys}}} \cdot M$.

Proof. The proof proceeds in four stages:

Stage 1. Using A1 and A2, construct a function-space decomposition and analyze the well-posedness of the BV-aware parameterization.

Stage 2. Based on one-sided Lipschitz analysis and a modified Gronwall inequality, using A2, A3, A4 to establish the trajectory error bound.

Stage 3. Introduce the physical PDE framework and Kruzhkov theory, stating the BV compactness required for Stage 4.

Stage 4. Using A0, A5, A6 to assemble the Wasserstein error, obtaining $W_1 \leq C_{\text{final}}\varepsilon^{1/2}$.

Notation (unified for the entire proof)

State space (diffusion model domain)

$u \in \mathbb{R}^d$ State vector ($d = N_x$ is the number of spatial grid points)

$\tau \in [0, T_d]$ Diffusion time (reverse: $T_d \to 0$, i.e., noise $\to$ data)

$p(u, \tau)$ Smoothed state density

$s(u, \tau) = \nabla_u \log p(u, \tau)$ True score field

$s_\theta(u, \tau)$ Network-approximated score field

$e(u, \tau) = s_\theta(u, \tau) - s(u, \tau)$ Score residual field

$\Sigma_{\text{score}}(\tau) \subset \mathbb{R}^d$ Shock manifold

$d_\Sigma(u, \tau)$ Signed distance function to $\Sigma_{\text{score}}(\tau)$

Physical space (PDE domain)

$x \in \Omega \subset \mathbb{R}$ Physical space coordinate

$t \in [0, T]$ Physical time (evolved by the PDE from initial data, orthogonal to τ)

$\mathbf{u}^\star(x)$ Kruzhkov entropy solution (at fixed $t = T_{\text{phys}}$)

$\mathbf{u}^\nu(x)$ Viscous PDE approximation with viscosity ν_{phys} (at $t = T_{\text{phys}}$)

∂_x Physical space partial derivative

$\|\cdot\|_{L^1(\Omega)}, \|\cdot\|_{L^2(\Omega)}$ L^p norms on physical space

Trajectories and generation

$\widehat{u}(\tau)$ ODE trajectory driven by the learned score ($\mathbb{R}^d$-valued)

$u^\natural(\tau)$ ODE trajectory driven by the true score ($\mathbb{R}^d$-valued)

$u^\theta := \widehat{u}(0)$ Final generated sample (at $\tau = 0$)

$\mu_\theta := \mathrm{Law}(u^\theta)$ Distribution of the generated samples

$\delta_{\mathbf{u}^\star}$ Dirac measure at the target Kruzhkov entropy solution

Norms and distances

$\|v\|$ Euclidean norm on state space $\mathbb{R}^d$

$\langle \cdot, \cdot \rangle$ $\mathbb{R}^d$ inner product

$\mathrm{TV}(f) = \int_\Omega |\partial_x f| \, dx$ Total variation seminorm

$W_1(\mu, \nu)$ 1-Wasserstein distance (Kantorovich–Rubinstein dual)

Time mapping The physical meaning of diffusion time τ decreasing from T_d to 0:

- $\tau = T_d$: pure noise (prior, e.g., standard Gaussian)
- $\tau = 0$: data sample (corresponding to the PDE solution at physical time $t = T_{\mathrm{phys}}$)
- **Viscosity matching**: $\sigma(\tau)^2(\tau) = 2\nu_{\mathrm{phys}}\tau$, so the evolution of τ is equivalent to a physical process with viscosity ν_{phys}; the drift coefficient of the reverse ODE is ν_{phys}
- In Stage 3, the target $\mathbf{u}^\star$ is the PDE entropy solution at a *fixed physical time* $t = T_{\mathrm{phys}}$ (T_{phys} is regarded as a constant and is not written explicitly in the sequel)
- Diffusion time τ and physical time t are *orthogonal*: τ describes the noising–denoising intensity, t describes the PDE evolution

A.5 Stage 1: Function Space and Error Localization

A.5.1 1.1 Setup

Let the dimensionless diffusion time of the target distribution be $\tau \in (\tau_{\mathrm{end}}, T_d]$. Denote the true smoothed data density by

$$p(u, \tau) = (\rho * G_\tau)(u), \qquad G_\tau(u) = (4\pi\tau)^{-d/2} \exp\bigl(-\|u\|^2/4\tau\bigr), \tag{90}$$

where $\rho(u)$ is the target data distribution. The true score field:

$$s(u, \tau) = \nabla_u \log p(u, \tau). \tag{91}$$

Define the shock manifold $\Sigma_{\mathrm{score}}(\tau) \subset \mathbb{R}^d$ as the equipotential surface in state space that connects different modes. For any $u \in \mathbb{R}^d$, define its signed distance to $\Sigma_{\mathrm{score}}(\tau)$:

$$d_\Sigma(u, \tau) = \inf_{v \in \Sigma_{\mathrm{score}}(\tau)} \|u - v\| \cdot \mathrm{sgn}(\text{mode label}). \tag{92}$$

The shock interfacial layer neighborhood and the outer smooth region:

$$\Omega_{\mathrm{sh}}(\tau) = \bigl\{u \in \mathbb{R}^d \;\bigm|\; |d_\Sigma(u, \tau)| \le \mathcal{O}(\tau)\bigr\}, \qquad \Omega_{\mathrm{out}}(\tau) = \mathbb{R}^d \setminus \Omega_{\mathrm{sh}}(\tau). \tag{93}$$

A.5.2 1.2 Lemma 1.1: Asymptotic Expansion and Singularity Decomposition of the True Score Field

Lemma 16. *Under Assumption A1, the true score field $s(u,\tau)$ can be decomposed on the whole space $\mathbb{R}^d$ as*

$$s(u,\tau) = s^{\mathrm{sm}}(u,\tau) + s^{\mathrm{sing}}(u,\tau) + \mathcal{O}(\tau), \tag{94}$$

where:

1. *$s^{\mathrm{sm}}(u,\tau)$ is uniformly Lipschitz continuous on the whole space: $\|\nabla_u s^{\mathrm{sm}}\|_{\mathrm{op}} \le L_{\mathrm{sm}} = \mathcal{O}(1)$.*
2. *$s^{\mathrm{sing}}(u,\tau)$ is non-negligible only inside $\Omega_{\mathrm{sh}}(\tau)$, and has the exact form*

$$s^{\mathrm{sing}}(u,\tau) = \frac{\Delta s_{\mathrm{jump}}(u)}{2}\tanh\Big(\frac{A(u,\tau)-B(u,\tau)}{2\tau}\Big), \tag{95}$$

where A, B are the potential functions of the dominant modes and $\Delta s_{\mathrm{jump}} = \nabla_u B - \nabla_u A$ is the jump intensity vector.

Proof sketch (Laplace method): By A1, inside $\Omega_{\mathrm{sh}}(\tau)$, the integral representation of $p(u,\tau)$ is dominated by the two nearest modes:

$$p(u,\tau) \sim C_A(u)\exp\Big(-\frac{A(u,\tau)}{\tau}\Big) + C_B(u)\exp\Big(-\frac{B(u,\tau)}{\tau}\Big). \tag{96}$$

Taking the log-gradient and using the identity $\nabla_u \log(e^{-A/\tau} + e^{-B/\tau})$ extracts the $\tanh$ part; the slowly varying coefficients C_A, C_B and their gradients are absorbed into s^{sm}. In $\Omega_{\mathrm{out}}(\tau)$, one mode is overwhelmingly dominant, $\tanh(\cdot) \to \pm 1$, $\nabla_u s^{\mathrm{sing}}$ tends to 0, and the score field degenerates to the smooth $\nabla_u A$ or $\nabla_u B$. The error term $\mathcal{O}(\tau)$ originates from the next-order remainder of the Laplace method; the uniform estimate is guaranteed by the mode-separation condition of A1. ■

Remark 17 (Nature of the singularity). Under standard parameterization, if the network directly fits $s(u,\tau)$, the Jacobian $\nabla_u s^{\mathrm{sing}}$ of s^{sing} contains a singular component of $\mathcal{O}(1/\tau)$ inside Ω_{sh}—this is precisely the origin of the global Lipschitz constant $L(\tau) \propto 1/\tau$ in classical theory, which causes the Gronwall integral to diverge and produces the exponential amplification factor $\exp(\Lambda T_d)$. The core idea of parameterization (5) is to *explicitly embed* this singularity into the network structure, so that the residual field $e(u,\tau)$ no longer inherits the $1/\tau$ singularity.

A.5.3 1.3 Lemma 1.2: Well-Posedness of the BV-Aware Parameterization

The parameterization (5) defines the network score field:

$$s_\theta(u,\tau) = \underbrace{\nabla_u \phi_\theta^{\mathrm{sm}}(u,\tau)}_{\text{smooth background term}} + \underbrace{\frac{\kappa_\theta(u,\tau)}{2}\tanh\Big(\frac{D_\theta(u,\tau)}{2\tau}\Big)\nabla_u D_\theta(u,\tau)}_{\text{interfacial singular term}}. \tag{97}$$

Lemma 18. *(Spatial alignment and network well-posedness) Under Assumption A2, the map $\Theta : (u,\tau) \mapsto (\phi_\theta^{\mathrm{sm}}, \kappa_\theta, D_\theta)$ has the following properties:*

1. *Target approximation: $\nabla_u \phi_\theta^{\mathrm{sm}} \approx s^{\mathrm{sm}}$, $\kappa_\theta \nabla_u D_\theta \approx \Delta s_{\mathrm{jump}}$, $D_\theta \approx A - B$.*
2. *Bounded Jacobian: The Jacobian norms $\|\nabla_u^2 \phi_\theta^{\mathrm{sm}}\|_{\mathrm{op}}$, $\|\nabla_u \kappa_\theta\|$, $\|\nabla_u^2 D_\theta\|_{\mathrm{op}}$ are all independent of τ, of order $\mathcal{O}(1)$.*
3. *The network learns only the smooth signed distance field D_θ. The singular factor $1/\tau$ is written as an explicit physical prior* outside *the network structure, appearing only inside* $\tanh$ *and in the singular Jacobian matrix J_{sh} after differentiation; the network involves no $1/\tau$ amplification from automatic differentiation, completely avoiding gradient explosion.*

Proof: By A2, $\nabla_u^2 \phi_\theta^{\mathrm{sm}}$, $\nabla_u^2 D_\theta$, $\nabla_u \kappa_\theta$ are all uniformly bounded. Note that the derivative of $\tanh$ with respect to its argument is $\mathrm{sech}^2(\cdot) \le 1$, so when computing ∇_u of s_θ, the dependence of τ in the denominator does not propagate divergently outward; it passes into the Jacobian and forms a strictly negative semi-definite singular part (proved in detail below). Since A and B are both $\mathcal{O}(1)$ potential functions, the network-learned $D_\theta(u,\tau)$ has gradient $\nabla_u D_\theta = \mathcal{O}(1)$, introducing no additional singular scale. ■

A.5.4 1.4 Boundedness of the Residual Field (Heuristic Discussion)

Remark 19 (Heuristic significance of Corollary 1.4; not a rigorous claim). The following is a heuristic argument for intuition only and is *not relied upon in subsequent steps of the proof.*

Define the score residual field $e(u,\tau) = s_\theta(u,\tau) - s(u,\tau)$, decomposed into $e = e_{\mathrm{sm}} + e_{\mathrm{sh}}$, where e_{sh} is the shock residual arising from interface location/intensity deviation.

In the standard architecture, e_{standard} cannot fit the steep gradient of $\tanh(\cdot/\tau)$ and itself has a gradient component of $-\mathcal{O}(1/\tau)$ inside Ω_{sh}, causing exponential error amplification. The BV-aware architecture embeds the $\tanh$ structure, making the error of e_{sh} manifest as a small shift of the shock center position; this shift is of the same order as the shift magnitude in the W_1 sense ($\Delta W_1 \propto \Delta x$), without involving multiplicative amplification of gradients.

More rigorously, if one can independently prove gradient bounds for e_{sm} and for the shock shift of e_{sh}, one can establish $\int_0^{T_d} \|\nabla_u e(u,\tau)\|_{\mathrm{eff}}\, d\tau \le C_0 = \mathcal{O}(1)$. Rigorization of this direction is left to future work (see technical point C1.4-G in Appendix C). In the present proof, the Gronwall argument of Stage 2 *does not rely on this bound* and uses only the integral estimate of $\|e\|$ itself (see Theorem 2.2).

A.6 Stage 2: Modified Gronwall Inequality and Trajectory Error Bound

A.6.1 2.1 Failure of the Classical Analysis and the Gronwall Trap

In the reverse generative process, consider two ODE trajectories—the reference trajectory $u^\natural(\tau)$ driven by the true score s and the approximate trajectory $\widehat{u}(\tau)$ driven by the network score s_θ. Both obey the probability-flow ODE under the viscosity matching strategy $\sigma(\tau)^2(\tau) = 2\nu_{\mathrm{phys}}\tau$ (drift coefficient $\sigma\dot\sigma = \nu_{\mathrm{phys}}$):

$$\frac{du^\natural}{d\tau} = -\nu_{\mathrm{phys}}\, s(u^\natural,\tau) =: V(u^\natural,\tau), \qquad \frac{d\widehat{u}}{d\tau} = -\nu_{\mathrm{phys}}\, s_\theta(\widehat{u},\tau) =: V_\theta(\widehat{u},\tau). \tag{98}$$

Initial condition (A4). By A4, the error between the two trajectories at $\tau = T_d$ satisfies

$$E(T_d) := \|\widehat{u}(T_d) - u^\natural(T_d)\| \le \tilde{C}\varepsilon, \tag{99}$$

(equal to 0 if the prior matches exactly; small deviations are handled uniformly by the Gronwall formula below).

Define the trajectory error:

$$E(\tau) := \|\widehat{u}(\tau) - u^\natural(\tau)\|. \tag{100}$$

Differentiating E^2:

$$\frac{1}{2}\frac{d}{d\tau}E(\tau)^2 = \big\langle \widehat{u} - u^\natural,\ V_\theta(\widehat{u},\tau) - V(u^\natural,\tau)\big\rangle. \tag{101}$$

Decompose the right-hand side into the self-consistency term (of V_θ) and the error-driven term:

$$\begin{aligned}\frac{1}{2}\frac{d}{d\tau}E(\tau)^2 \le\ & \underbrace{\big\langle \widehat{u} - u^\natural,\ V_\theta(\widehat{u},\tau) - V_\theta(u^\natural,\tau)\big\rangle}_{\text{Term I: drift self-consistency}} \\ & + \underbrace{\nu_{\mathrm{phys}}\big\langle \widehat{u} - u^\natural,\ e(u^\natural,\tau)\big\rangle}_{\text{Term II: score error-driven}}.\end{aligned} \tag{102}$$

Applying Cauchy–Schwarz to Term II: $\nu_{\mathrm{phys}}\langle \widehat{u} - u^\natural, e(u^\natural,\tau)\rangle \le \nu_{\mathrm{phys}}\|e(u^\natural,\tau)\| \cdot E(\tau)$. Thus:

$$\frac{1}{2}\frac{d}{d\tau}E(\tau)^2 \le \underbrace{\big\langle \widehat{u} - u^\natural,\ V_\theta(\widehat{u},\tau) - V_\theta(u^\natural,\tau)\big\rangle}_{\text{Term I}} + \nu_{\mathrm{phys}}\|e(u^\natural,\tau)\| \cdot E(\tau). \tag{103}$$

In the classical proof, Term I is bounded using the global Lipschitz constant $L_\theta(\tau) = \mathcal{O}(1/\tau)$, causing the Gronwall integral to diverge as $\exp(\Lambda T_d)$. Below we replace this with a one-sided Lipschitz constant.

A.6.2 2.2 Lemma 2.1: One-Sided Lipschitz Condition and Negative Semi-Definiteness

Definition 20 (One-sided Lipschitz constant)**.** The one-sided Lipschitz constant $\mu(\tau)$ of V_θ is the smallest number such that for all $u, v \in \mathbb{R}^d$:

$$\big\langle u - v,\ V_\theta(u,\tau) - V_\theta(v,\tau)\big\rangle \le \mu(\tau)\|u - v\|^2. \tag{104}$$

When V_θ is differentiable, $\mu(\tau)$ equals the maximum eigenvalue of the symmetric part of its Jacobian $J_\theta(u,\tau) \coloneqq \nabla_u V_\theta(u,\tau)$:

$$\mu(\tau) = \sup_{u\in\mathbb{R}^d} \lambda_{\max}\Big(\frac{J_\theta(u,\tau) + J_\theta(u,\tau)^\mathsf{T}}{2}\Big). \tag{105}$$

Lemma 21. *(Bounded logarithmic norm of the BV-aware architecture) Under Assumption A2,*

$$\mu(\tau) \le L_{\mathrm{sm}} = \mathcal{O}(1), \tag{106}$$

i.e., the one-sided Lipschitz constant is independent of τ and has no singularity.

Proof: Compute the Jacobian of $V_\theta(u,\tau) = -\nu_{\mathrm{phys}} s_\theta(u,\tau)$:

$$J_\theta = \nabla_u V_\theta = J_{\mathrm{sm}} + J_{\mathrm{sh}} + J_{\mathrm{cross}}, \tag{107}$$

with the following components.

(a) Smooth part. $J_{\mathrm{sm}} = -\nu_{\mathrm{phys}} \nabla_u^2 \phi_\theta^{\mathrm{sm}}$. By A2, $\|\nabla_u^2 \phi_\theta^{\mathrm{sm}}\|_{\mathrm{op}} = \mathcal{O}(1)$, so $\lambda_{\max}(J_{\mathrm{sm}}) \le \nu_{\mathrm{phys}} \cdot \mathcal{O}(1) = \mathcal{O}(1)$ (since $\nu_{\mathrm{phys}} = \Theta(\varepsilon)$ is small but $\varepsilon \le 1$, so it does not exceed $\mathcal{O}(1)$).

(b) Cross terms. J_{cross} contains derivative terms of κ_θ and $\nabla_u D_\theta$. By A2 these are all uniformly bounded; moreover, the dominant terms produced by these derivatives are already controlled by explicit separation and internal derivative control, introducing no $1/\tau$ factor into this term, so $\lambda_{\max}(J_{\mathrm{cross}}) = \mathcal{O}(1)$.

(c) Singular part. Differentiating the $\tanh$ term in the parameterization with respect to ∇_u yields

$$J_{\mathrm{sh}}(u,\tau) = -\nu_{\mathrm{phys}} \cdot \frac{\kappa_\theta(u,\tau)}{4\tau}\ \mathrm{sech}^2\Big(\frac{D_\theta(u,\tau)}{2\tau}\Big)\ \big(\nabla_u D_\theta\big)\big(\nabla_u D_\theta\big)^\mathsf{T}. \tag{108}$$

The matrix $(\nabla_u D_\theta)(\nabla_u D_\theta)^\mathsf{T}$ is a rank-1 positive semi-definite matrix (of the form $\mathbf{n}\mathbf{n}^\mathsf{T}$, where $\mathbf{n} = \nabla_u D_\theta$ is the shock interface normal vector). By A2, $\kappa_\theta \ge \kappa_0 > 0$; together with $\nu_{\mathrm{phys}} > 0$, $\tau > 0$, $\mathrm{sech}^2(\cdot) > 0$, the leading negative sign makes J_{sh} *strictly negative semi-definite*.

Hence the maximum eigenvalue of the symmetric part of J_{sh} satisfies $\lambda_{\max}(J_{\mathrm{sh}}) \le 0$, contributing nothing positive to the one-sided Lipschitz constant.

(d) Physical interpretation. In the direction normal to the shock surface (the $\mathbf{n}$ direction), J_{sh} provides a negative eigenvalue of magnitude $-\nu_{\mathrm{phys}}\kappa_0\|\nabla_u D_\theta\|^2/(4\tau)$, corresponding to a *strong attractive force* of the reverse diffusion flow in the shock normal direction. Errors that deviate from the true shock manifold are compressed at this enormous rate rather than amplified—this is the mathematical essence of Burgers characteristic line convergence, and the core guarantee that this architecture avoids exponential divergence.

Combining (a)–(d),

$$\mu(\tau) \le \lambda_{\max}\big(J_{\mathrm{sm}} + J_{\mathrm{cross}}\big) + \underbrace{\lambda_{\max}(J_{\mathrm{sh}})}_{\le 0} \le L_{\mathrm{sm}} = \mathcal{O}(1). \tag{109}$$

■

A.6.3 2.3 Theorem 2.2: Modified Flow Stability with Constant Upper Bound

Lemma 22. *(Trajectory error bound) Under Assumptions A2, A3, A4, let $\varepsilon_{\mathrm{raw}}$ be the raw (unnormalized) score matching mean-squared error bound, i.e., $\mathbb{E}[\|e\|^2] \le \varepsilon_{\mathrm{raw}}^2$. By the grid-normalized definition of A3, $\mathbb{E}[\Delta x\|e\|^2] \le \varepsilon^2$, so $\varepsilon_{\mathrm{raw}}^2 = \varepsilon^2/\Delta x$. Then*

$$\mathbb{E}_z\big[E(0)\big] \le C_1\varepsilon_{\mathrm{raw}} = C_1\frac{\varepsilon}{\sqrt{\Delta x}}, \tag{110}$$

where C_1 depends only on $T_d, \nu_{\mathrm{phys}}, L_{\mathrm{sm}}$. In the norm conversion of Stage 4, $\sqrt{\Delta x}$ will be canceled, yielding a grid-independent final bound.

Proof: **Step 1: Scalar differential inequality.** Applying Lemma 2.1 to Term I in the derivative of $\frac{1}{2}E^2$ gives

$$\frac{1}{2}\frac{d}{d\tau}E(\tau)^2 \le \mu(\tau)\,E(\tau)^2 + \nu_{\text{phys}}\,\|e(u^\natural,\tau)\| \cdot E(\tau). \tag{111}$$

Dividing by $E(\tau)$ (when $E(\tau) > 0$; the inequality holds trivially when $E(\tau) = 0$),

$$\frac{d}{d\tau}E(\tau) \le \mu(\tau)\,E(\tau) + \nu_{\text{phys}}\,\|e(u^\natural,\tau)\|. \tag{$\star$}$$

Step 2: Apply Gronwall's inequality. From ($\star$), integrating from $\tau = T_d$ to $\tau = 0$,

$$E(0) \le E(T_d) \cdot \exp\Big(\int_0^{T_d} \mu(r)\,dr\Big) + \int_0^{T_d} \exp\Big(\int_0^{\tau} \mu(r)\,dr\Big)\,\nu_{\text{phys}}\,\|e(u^\natural,\tau)\|\,d\tau. \tag{112}$$

By Lemma 2.1, $\mu(r) \le L_{\text{sm}} = \mathcal{O}(1)$, so

$$\exp\Big(\int_0^{\tau} \mu(r)\,dr\Big) \le e^{L_{\text{sm}}T_d} =: C_{\text{flow}} = \mathcal{O}(1). \tag{113}$$

Substituting the initial error bound $E(T_d) \le \tilde{C}\varepsilon$ from A4:

$$E(0) \le \tilde{C}C_{\text{flow}}\,\varepsilon + C_{\text{flow}}\,\nu_{\text{phys}} \int_0^{T_d} \|e(u^\natural,\tau)\|\,d\tau. \tag{$\star\star$}$$

Step 3: Integral bound of the score matching error (directly from A3). Apply Cauchy–Schwarz to the integral $\int_0^{T_d} \|e(u^\natural,\tau)\|\,d\tau$:

$$\int_0^{T_d} \|e(u^\natural,\tau)\|\,d\tau \le \sqrt{T_d} \cdot \Big(\int_0^{T_d} \|e(u^\natural,\tau)\|^2\,d\tau\Big)^{1/2}. \tag{114}$$

Take expectation over z (prior noise) of the right-hand side. By A3, $\mathbb{E}[\Delta x\|e\|^2] \le \varepsilon^2$, i.e., the raw (unnormalized) mean-squared error satisfies $\mathbb{E}[\|e\|^2] \le \varepsilon^2/\Delta x = \varepsilon_{\text{raw}}^2$. Hence

$$\mathbb{E}_z\Big[\int_0^{T_d} \|e(u^\natural,\tau)\|^2\,d\tau\Big] \le T_d \cdot \varepsilon_{\text{raw}}^2 = T_d \cdot \frac{\varepsilon^2}{\Delta x}, \tag{115}$$

(this step holds because the marginal distribution of $u^\natural(\tau)$ is $p(\cdot,\tau)$, so the expectation over $(u^\natural(\tau),\tau)$ matches $\mathbb{E}_{u,\tau}$ in A3).

Now take expectation of the Cauchy–Schwarz result (using Jensen's inequality):

$$\mathbb{E}_z\Big[\int_0^{T_d} \|e(u^\natural,\tau)\|\,d\tau\Big] \le \sqrt{T_d} \cdot \sqrt{T_d\,\varepsilon_{\text{raw}}^2} = T_d\,\varepsilon_{\text{raw}} = T_d\frac{\varepsilon}{\sqrt{\Delta x}}. \tag{116}$$

Step 4: Assembly. Substituting Step 3 into ($\star\star$) and taking expectation:

$$\mathbb{E}_z[E(0)] \le \tilde{C}C_{\text{flow}}\,\varepsilon_{\text{raw}} + C_{\text{flow}}\,\nu_{\text{phys}}\,T_d\,\varepsilon_{\text{raw}} =: C_1\varepsilon_{\text{raw}}, \tag{117}$$

where $C_1 = C_{\text{flow}}(\tilde{C} + \nu_{\text{phys}}T_d) = \mathcal{O}(1)$ (by A5, $\nu_{\text{phys}} = \Theta(\varepsilon) \le 1$, so $\nu_{\text{phys}}T_d = \mathcal{O}(1)$). From $\varepsilon_{\text{raw}} = \varepsilon/\sqrt{\Delta x}$, we obtain $\mathbb{E}_z[E(0)] \le C_1\varepsilon/\sqrt{\Delta x}$. ■

Conclusion. Under the BV-aware architecture, the trajectory endpoint error is $\mathbb{E}[E(0)] = \mathcal{O}(\varepsilon_{\text{raw}})$, free of the exponential amplification factor.

A.7 Stage 3: Kruzhkov Entropy Solution L^1 Contraction Framework

Remark 23 (Convention for Stage 3). We now introduce physical space $x \in \Omega \subset \mathbb{R}$. The components of the state vector $u \in \mathbb{R}^d$ ($d = N_x$) are $u_i = \mathbf{u}(x_i)$, where $\{x_i\}$ is a uniform grid in physical space (spacing $\Delta x = |\Omega|/N_x$). The discrete L^1 norm is defined as $\|\mathbf{u}\|_{L^1} = \Delta x \sum_{i=1}^{N_x} |\mathbf{u}(x_i)|$, which approximates the continuous L^1 norm on physical space (see the norm equivalence claim in Stage 4).

A.7.1 3.1 Kruzhkov Entropy Condition and L^1 Contraction

Consider the target hyperbolic conservation law (on physical space $\Omega \times [0, T]$):

$$\partial_t \mathbf{u} + \partial_x f(\mathbf{u}) = 0, \qquad x \in \Omega \subset \mathbb{R}, \quad t \in [0, T], \tag{118}$$

where f is a flux function (e.g., Burgers flux $f(u) = u^2/2$), with initial condition $\mathbf{u}(\cdot, 0) = \mathbf{u}_0$. The Kruzhkov entropy solution $\mathbf{u}^\star$ satisfies: for every constant $k \in \mathbb{R}$, in the sense of distributions,

$$\partial_t |\mathbf{u}^\star - k| + \partial_x \big(\operatorname{sgn}(\mathbf{u}^\star - k)(f(\mathbf{u}^\star) - f(k))\big) \le 0. \tag{119}$$

Corollary of Kruzhkov's theorem (L^1-contractive semigroup). If $\mathbf{a}(x, t)$ and $\mathbf{b}(x, t)$ both satisfy the Kruzhkov entropy condition, then

$$\|\mathbf{a}(\cdot, t) - \mathbf{b}(\cdot, t)\|_{L^1(\Omega)} \le \|\mathbf{a}(\cdot, 0) - \mathbf{b}(\cdot, 0)\|_{L^1(\Omega)}. \tag{120}$$

Set $\mathbf{u}^\star(x) \coloneqq \mathbf{u}^\star(x, T_{\text{phys}})$ (at fixed physical time T_{phys}; omitted in the sequel).

A.7.2 3.2 Loss Design and BV Compactness

The loss function $\mathcal{L}_3$ augments the standard denoising score matching loss with a BV regularizer:

$$\mathcal{L}_3 = \mathcal{L}_{\text{DSM}} + \lambda_{\text{BV}} \operatorname{TV}(D_\theta). \tag{121}$$

By Assumption A0, together with the hard-coded $\tanh$ structure of parameterization (5), this regularizer guarantees that the generated samples almost surely belong to the bounded BV ball $\mathcal{K}_M$. By Helly's selection theorem, $\mathcal{K}_M$ is relatively compact in the $L^1(\Omega)$ topology.

A.7.3 3.3 Viscous Limit (Application of Kuznetsov's Lemma)

The reverse stage of the diffusion model, under viscosity matching $\sigma(\tau)^2(\tau) = 2\nu_{\text{phys}}\tau$, effectively introduces artificial viscosity ν_{phys}. Denote by $\mathbf{u}^\nu$ the solution of the viscous physical PDE at $t = T_{\text{phys}}$:

$$\partial_t \mathbf{u}^\nu + \partial_x f(\mathbf{u}^\nu) = \nu_{\text{phys}}\, \partial_{xx} \mathbf{u}^\nu. \tag{122}$$

By Assumption A6 (based on Kuznetsov 1976):

$$\|\mathbf{u}^\nu - \mathbf{u}^\star\|_{L^1(\Omega)} \le C_K \sqrt{\nu_{\text{phys}}}. \tag{123}$$

Together with A5 ($\nu_{\text{phys}} = \Theta(\varepsilon)$), we obtain

$$\|\mathbf{u}^\nu - \mathbf{u}^\star\|_{L^1(\Omega)} \le C_K \sqrt{C\varepsilon} = C_K \sqrt{C} \cdot \varepsilon^{1/2}. \tag{3.1}$$

A.8 Stage 4: Assembly of the Wasserstein Error and Emergence of $\mathcal{O}(\varepsilon^{1/2})$

A.8.1 4.1 Grid Renormalization, Norm Equivalence, and Neural Generation Error

Derivation of the discrete L^1–normalized Euclidean norm equivalence. The physical domain Ω is discretized into $d = N_x$ uniform grid points with spacing $\Delta x = |\Omega|/d$. For a state vector $u \in \mathbb{R}^d$, the discrete L^1 norm is

$$\|\mathbf{u}\|_{L^1} = \Delta x \sum_{i=1}^{d} |u_i|. \tag{124}$$

From the ℓ^1–ℓ^2 norm inequality ($\sum |u_i| \le \sqrt{d}\sqrt{\sum u_i^2} = \sqrt{d}\|u\|$), substituting Δx, we obtain

$$\|\mathbf{u}\|_{L^1} \le \Delta x \sqrt{d}\, \|u\| = \sqrt{|\Omega|} \cdot \sqrt{\Delta x}\, \|u\|. \tag{4.1}$$

Deriving the L^1 bound for the neural generation error. Note that in the renormalized Assumption A3, the error bound is given in the form $\mathbb{E}[\Delta x \|e\|^2] \le \varepsilon^2$, which means that for the pure Euclidean norm, we have $\mathbb{E}[\|e\|^2] \le \varepsilon^2/\Delta x$.

Recall from Stage 2, the Gronwall integration and trajectory error bound Lemma 2.2 give, for the standard Euclidean distance $E(0)$:

$$\mathbb{E}_z\big[\|\widehat{u}(0) - u^\natural(0)\|\big] \le C_1\, \mathbb{E}_z\big[\|e\|\big] \le C_1 \frac{\varepsilon}{\sqrt{\Delta x}}. \tag{4.2}$$

The corresponding difference of physical functions is $u^\theta - \mathbf{u}^\nu$. Combining with the norm scaling relation (4.1):

$$\mathbb{E}_z\big[\|u^\theta - \mathbf{u}^\nu\|_{L^1(\Omega)}\big] \le \sqrt{|\Omega|}\sqrt{\Delta x} \cdot \mathbb{E}_z\big[\|\widehat{u}(0) - u^\natural(0)\|\big] \le \sqrt{|\Omega|}\sqrt{\Delta x} \cdot C_1 \frac{\varepsilon}{\sqrt{\Delta x}}. \tag{125}$$

Here d and Δx are *perfectly canceled*! Regardless of how fine the grid is, the error bound remains valid, yielding a grid-independent continuum limit bound:

$$\mathbb{E}_z\big[\|u^\theta - \mathbf{u}^\nu\|_{L^1(\Omega)}\big] \le C_1\sqrt{|\Omega|}\,\varepsilon =: C_2\varepsilon. \tag{4.3}$$

Remark 24 (Place of the Gagliardo–Nirenberg interpolation). An earlier draft used a Gagliardo–Nirenberg-type interpolation inequality $\|h\|_{L^1} \le C_{\mathrm{GN}}\|h\|_{L^2}^{2/3}\mathrm{TV}(h)^{1/3}$ to convert L^2 error into L^1 error, obtaining an $\mathcal{O}(\varepsilon^{2/3})$ bound. The present revised proof uses normalized L^2-matching combined with the norm equivalence (4.1), obtaining a sharper, dimension-independent L^1 error $\mathcal{O}(\varepsilon)$ directly from the scaled Euclidean error.

A.8.2 4.2 Triangle Inequality Decomposition of the Wasserstein Distance

The 1-Wasserstein distance (Kantorovich–Rubinstein dual):

$$W_1(\mu,\nu) = \inf_{\pi\in\Pi(\mu,\nu)} \int \|\mathbf{u} - \mathbf{v}\|_{L^1(\Omega)}\, d\pi(\mathbf{u},\mathbf{v}). \tag{126}$$

Let $\delta_{\mathbf{u}^\nu}$ be the Dirac measure at the viscous solution (given the initial data $\mathbf{u}_0$, the viscous PDE solution is unique, hence deterministic). Decompose via the triangle inequality:

$$W_1(\mu_\theta, \delta_{\mathbf{u}^\star}) \le \underbrace{W_1(\mu_\theta, \delta_{\mathbf{u}^\nu})}_{\text{Neural generation error (Term A)}} + \underbrace{W_1(\delta_{\mathbf{u}^\nu}, \delta_{\mathbf{u}^\star})}_{\text{Viscous limit error (Term B)}}. \tag{4.4}$$

Term A (Neural generation error). Since $\widehat{u}(0)$ and $u^\natural(0)$ are deterministically generated from the same prior noise z (the ODE has a unique solution for fixed z), there exists a natural pointwise coupling:

$$\pi_z = \big(u^\theta(z),\ \mathbf{u}^\nu\big)_{\#}\, p_{T_d}, \tag{127}$$

where p_{T_d} is the prior distribution and $\mathbf{u}^\nu$ is regarded as a constant vector (independent of z). By the feasibility of this coupling and (4.3):

$$W_1(\mu_\theta, \delta_{\mathbf{u}^\nu}) \le \int \|u^\theta(z) - \mathbf{u}^\nu\|_{L^1(\Omega)}\, dp_{T_d}(z) = \mathbb{E}_z\big[\|u^\theta - \mathbf{u}^\nu\|_{L^1(\Omega)}\big] \le C_2\varepsilon. \tag{4.5}$$

Term B (Viscous limit error). Both $\delta_{\mathbf{u}^\nu}$ and $\delta_{\mathbf{u}^\star}$ are single-point distributions; W_1 degenerates to the L^1 distance:

$$W_1(\delta_{\mathbf{u}^\nu}, \delta_{\mathbf{u}^\star}) = \|\mathbf{u}^\nu - \mathbf{u}^\star\|_{L^1(\Omega)}. \tag{128}$$

From (3.1) (the joint conclusion of A6 and A5):

$$W_1(\delta_{\mathbf{u}^\nu}, \delta_{\mathbf{u}^\star}) \le C_K\sqrt{C} \cdot \varepsilon^{1/2}. \tag{4.6}$$

A.8.3 4.3 Final Conclusion

Substituting (4.5) and (4.6) into (4.4):

$$W_1(\mu_\theta, \delta_{\mathbf{u}^\star}) \le C_2\varepsilon + C_K\sqrt{C} \cdot \varepsilon^{1/2}. \tag{129}$$

For $\varepsilon \in (0,1)$, we have $\varepsilon \le \varepsilon^{1/2}$, so the higher-order term $\mathcal{O}(\varepsilon)$ is absorbed by the lower-order term $\mathcal{O}(\varepsilon^{1/2})$:

$$W_1(\mu_\theta, \delta_{\mathbf{u}^\star}) \le \left(C_2 + C_K\sqrt{C}\right) \cdot \varepsilon^{1/2} =: C_{\text{final}} \cdot \varepsilon^{1/2}. \tag{130}$$

Here $C_{\text{final}} = C_2 + C_K\sqrt{C}$ depends only on $M, T_d, T_{\text{phys}}, \kappa_0, L_{\text{sm}}, \nu_{\text{phys}}/\varepsilon$, all independent of ε. Thus

$$\boxed{W_1(\mu_\theta, \delta_{\mathbf{u}^\star}) \le C_{\text{final}}\, \varepsilon^{1/2}.} \tag{131}$$

This completes the proof of Theorem 3. □

Appendix A: Symbol Correspondence Table (Old → Revised)

Table 3: Symbol correspondence table

Old symbol	Revised symbol
$u \in \mathbb{R}^d$ (state point) and $u(x,t)$ (PDE solution) confused	State space: $u \in \mathbb{R}^d$; physical space: $\mathbf{u}(x)$
u^θ used for both true trajectory and PDE solution	$u^\natural(\tau)$ (true trajectory); $\mathbf{u}^\star$ (PDE solution)
$\widehat{u}$ confused as trajectory/sample/distribution	$\widehat{u}(\tau)$ (trajectory); u^θ (sample); μ_θ (distribution)
∇_u vs ∂_x confused	∇_u (state-space gradient); ∂_x (physical space partial derivative)
$\sup_u \lambda_{\max}$ with ambiguous scope	$\sup_{u\in\mathbb{R}^d} \lambda_{\max}$
$\|E(\tau)\|$ with ambiguous norm type	$E(\tau) := \|\widehat{u}(\tau) - u^\natural(\tau)\|$ ($\mathbb{R}^d$ Euclidean norm)
Dual role of u in residual field $e(u,\tau)$	The parameter u in $e(u,\tau)$ is a state-space point
Relation between times τ and t unclear	Time mapping paragraph explicitly defined

Appendix B: Summary of Modifications

Table 4: Summary of modifications

No.	Modification	Treatment
M1	Rigorous gradient bound claim in Corollary 1.4	Downgraded to heuristic remark in §A.5; not relied upon s
M2	The $\int \|e\|\, d\tau$ bound in Lemma 2.2	Directly from A3 + Cauchy–Schwarz; no dependence on C
M3	GN interpolation in Stage 4	Removed; replaced with discrete norm equivalence
M4	Viscous error Term B	Directly cites A5 + A6 (Equation (3.1)); logical chain
M5	Theorem statement	Explicitly references A0–A6; no dangling assumpt
M6	Assumption system	Newly added A0–A6, with verifiability annotations and engineering

Appendix C: Technical Points Requiring Further Rigorization and Scope of Engineering Approximations

Table 5: Checklist of technical points

No.	Technical point	Status	Scope of influence
C1.1-P	"Finitely separated Gaussian clusters" assumption of Lemma 1.1	Upgraded to A1	If A1 fails, the score singula
C1.4-G	Residual gradient bound of Corollary 1.4	Left to future work	This proof does not rely on
C2.1-κ	Training-time enforcement of $\kappa_\theta > 0$	Upgraded to A2	If not strictly enforced, conc
EA1	Applicability of A1 to general continuous distributions	Engineering approximation	Affects the uniformity of th
EA2	Feasibility of schedule $\nu_{\text{phys}} = \Theta(\varepsilon)$ in A5	Engineering approximation	If $\nu_{\text{phys}} \gg \varepsilon$, Term B domi
P4.1-GN	GN interpolation argument in the continuum limit	Retained in remark	Does not affect conclusion
K3.3-ν	Generalization of Kuznetsov bound in A6	Rigorously established	Only requires initial BV bo

Table 6: Summary of modifications

No.	Modification	Treatment
M1	Rigorous gradient bound claim in Corollary 1.4	Downgraded to heuristic remark in §A.5; not relied upon s
M2	The $\int \|e\| \, d\tau$ bound in Lemma 2.2	Directly from A3 + Cauchy–Schwarz; no dependence on C
M3	GN interpolation in Stage 4	Removed; replaced with discrete norm equivalence
M4	Viscous error Term B	Directly cites A5 + A6 (Equation (3.1)); logical chain
M5	Theorem statement	Explicitly references A0–A6; no dangling assumpt
M6	Assumption system	Newly added A0–A6, with verifiability annotations and engineering

Table 7: Checklist of technical points

No.	Technical point	Status	Scope of influence
C1.1-P	"Finitely separated Gaussian clusters" assumption of Lemma 1.1	Upgraded to A1	If A1 fails, the score singula
C1.4-G	Residual gradient bound of Corollary 1.4	Left to future work	This proof does not rely on
C2.1-κ	Training-time enforcement of $\kappa_\theta > 0$	Upgraded to A2	If not strictly enforced, conc
EA1	Applicability of A1 to general continuous distributions	Engineering approximation	Affects the uniformity of th
EA2	Feasibility of schedule $\nu_{\text{phys}} = \Theta(\varepsilon)$ in A5	Engineering approximation	If $\nu_{\text{phys}} \gg \varepsilon$, Term B domi
P4.1-GN	GN interpolation argument in the continuum limit	Retained in remark	Does not affect conclusion
K3.3-ν	Generalization of Kuznetsov bound in A6	Rigorously established	Only requires initial BV bo

Appendix B: Summary of Modifications

Appendix C: Technical Points Requiring Further Rigorization and Scope of Engineering Approximations

A.9 Proof of Theorem 25 (Shock-location admissibility)

Theorem 25 (Shock-location admissibility)**.** *In the limit $\sigma(\tau) \to 0$ (i.e., the final step of inference), the solution produced by the proposed sampler satisfies the Rankine–Hugoniot condition at each shock location $x_s(t)$ almost everywhere,*

$$\dot{x}_s(t) = \frac{f(u_L) - f(u_R)}{u_L - u_R}, \tag{132}$$

and the Lax entropy condition $f'(u_L) \geq \dot{x}_s \geq f'(u_R)$.

Proof. **Assumptions.**

1. **(Perfect score matching at the interfacial layer.)** The BV-aware score parameterization (5) is optimized to global optimality in a neighborhood of the discontinuity, i.e. $s_\theta(u, \tau) \equiv \nabla_u \log p_\tau(u)$. We thus analyze the true score field.

2. **(Single-shock target manifold.)** The target distribution $\rho_t(u)$ concentrates on a submanifold $\mathcal{M}_t$ in function space (or its discretization $\mathbb{R}^N$).For the sampling trajectory under consideration, the corresponding target Kruzhkov entropy solution $\mathbf{u}^\star(x, t)$ at the current time t possesses a single isolated shock located at $x_s(t)$, with left and right limits

$$u_L \coloneqq \mathbf{u}^\star(x_s^-, t), \qquad u_R \coloneqq \mathbf{u}^\star(x_s^+, t). \tag{133}$$

3. **(Local planarity in high dimension.)** Since the grid dimension N is large, in the limit $\tau \to 0$ the separating manifold of data modes (i.e., the locus where the score exhibits a shock) can be locally approximated as a hyperplane. Its signed distance function $\phi_\theta^{\text{sh}}(u, \tau)$ satisfies $|\nabla_u \phi_\theta^{\text{sh}}| = 1$.

4. **(Compressive score gradient.)** When the reverse sampling trajectory $u(\tau)$ approaches the target manifold $\mathcal{M}_t$, the local flow induced by the network parameterization s_θ is compressive along the shock normal $\nabla \phi_\theta^{\text{sh}}$:

$$\partial_{\phi_\theta^{\text{sh}}} s_\theta \cdot \nabla \phi_\theta^{\text{sh}} < 0. \tag{134}$$

This guarantees that the score field acts as a centripetal force, pushing probability mass toward the interfacial manifold.

Part I: Geometric correspondence via the viscosity-matched schedule.

Let the forward noising process be a variance-exploding SDE. Under the schedule (3), the noise variance is tied to the physical viscosity ν_{phys}:

$$\sigma(\tau)^2 = 2\nu_{\text{phys}}\,\tau, \qquad \tau \in [0, T_d]. \tag{135}$$

The forward SDE then reads

$$du = \sqrt{2\nu_{\text{phys}}}\,dw. \tag{136}$$

The corresponding Fokker–Planck equation for the density $p(u,\tau)$ in data space $\mathbb{R}^N$ is

$$\partial_\tau p(u,\tau) = \nu_{\text{phys}}\,\Delta_u p(u,\tau). \tag{137}$$

Differentiating the score $s(u,\tau) = \nabla_u \log p = \nabla_u p/p$ with respect to τ and invoking (137) gives the exact evolution of the score:

$$\begin{aligned}\partial_\tau s &= \nu_{\text{phys}}\,\nabla_u\Big(\frac{\Delta_u p}{p}\Big)\\ &= \nu_{\text{phys}}\,\Delta_u s + \nu_{\text{phys}}\,\nabla_u(|s|^2).\end{aligned} \tag{138}$$

Since s is a gradient field, $\nabla_u(|s|^2) = 2(s\cdot\nabla_u)s$. Note that the relation $s = \nabla_u \log p$ is essentially the classical *inverse Cole–Hopf transform*, which explicitly converts the linear heat equation (Fokker–Planck) into a nonlinear PDE. We thus obtain the **high-dimensional viscous Burgers equation at the score level**:

$$\partial_\tau s + (-2\nu_{\text{phys}}\,s\cdot\nabla_u)\,s = \nu_{\text{phys}}\,\Delta_u s. \tag{139}$$

Lemma 26 (Effective viscosity equivalence)**.** *Via the schedule* (3)*, the effective viscosity of the score dynamics is numerically identical to the physical viscosity* ν_{phys} *of the target equation.*

The target PDE with small physical viscosity is

$$\partial_t u(x,t) + \partial_x f(u(x,t)) = \nu_{\text{phys}}\,\partial_{xx} u(x,t). \tag{140}$$

Under discretization, the components of $u \in \mathbb{R}^N$ satisfy $u_i \approx u(x_i,t)$. The target distribution $\rho_t(u)$ (the boundary condition at $\tau = 0$) is the physical weak-solution distribution. In the limit $\tau \to 0$, the reverse sampling trajectory $u(\tau)$ is pulled by the score $s(u,\tau)$ toward the support manifold $\mathcal{M}_t$ of $\rho_t(u)$.

The BV-aware score parameterization (5) reads

$$s_\theta(u,\tau) = \nabla\phi_\theta^{\text{sm}} + \frac{\kappa_\theta}{2}\tanh\Big(\frac{\phi_\theta^{\text{sh}}}{2}\Big)\nabla\phi_\theta^{\text{sh}}. \tag{141}$$

Near the interface, the smooth background $\nabla\phi_\theta^{\text{sm}}$ is negligible, and the true score exhibits the asymptotic inner-scale form

$$s(u,\tau) \sim \frac{\kappa_\theta(u)}{2}\tanh\Big(\frac{\operatorname{dist}(u,\mathcal{M}_t)}{\sqrt{2\nu_{\text{phys}}\,\tau}}\Big)\mathbf{n}(u), \tag{142}$$

where $\mathbf{n}(u) = \nabla_u \operatorname{dist}(u,\mathcal{M}_t)$ is the normal vector to the manifold.

Lemma 27 (Geometric correspondence in the double limit)**.** *When* $\tau \to 0$ *and* $\nu_{\text{phys}} \to 0$ *hold simultaneously, the* $\tanh$ *profile in* (142) *degenerates to a step function. Consequently, the location where the score field exhibits a shock is determined in data space by* $\operatorname{dist}(u,\mathcal{M}_t) = 0$*. Mapped back to physical space, this manifold boundary coincides exactly with the grid coordinates where* $u(x,t)$ *exhibits a jump, i.e., the physical shock position* $x_s(t)$*.*

Part II: Weak-form limit and derivation of the Rankine–Hugoniot condition.

2.1 Tweedie profile near the shock.

For the reverse diffusion, the clean physical state $\widehat{u}_0(x,t)$ is estimated via the Tweedie formula

$$\widehat{u}_0(x,t) = u(\tau) + \sigma(\tau)^2\, s(u,\tau). \tag{143}$$

Under the schedule (3) and the BV-aware parameterization, restricted to a one-dimensional physical section, the signed distance expands as $\phi_\theta^{\mathrm{sh}} \approx \dfrac{x - x_s(t)}{\sqrt{2\nu_{\mathrm{phys}}\,\tau}}$.

Lemma 28 (Viscous shock profile)**.** *In a neighborhood of $x_s(t)$, the Tweedie-reconstructed physical field $\widehat{u}^\tau(x,t)$ possesses a smooth, monotone shock transition layer. To make the physical scales explicit, define the layer thickness $\varepsilon = \sigma(\tau) = \sqrt{2\nu_{\mathrm{phys}}\,\tau}$. It is described by*

$$\widehat{u}^\tau(x,t) \approx \frac{u_L + u_R}{2} - \frac{u_L - u_R}{2}\tanh\Big(\frac{x - x_s(t)}{2\varepsilon}\Big). \tag{144}$$

Thus $\widehat{u}^\tau$ is strictly between u_L and u_R and has controlled total variation.

2.2 Godunov guidance and weak form.

During generation, an entropy-compatible Godunov flux difference is used as PDE guidance. Thus, as $\tau \to 0$, the generated sequence $\widehat{u}^\tau$ approximately satisfies a weak form: for every smooth test function $\varphi \in C_c^\infty(\mathbb{R} \times (0,T))$,

$$I(\varepsilon) = \iint_{\mathbb{R}\times\mathbb{R}^+} \big(\widehat{u}^\tau\, \partial_t\varphi + f(\widehat{u}^\tau)\, \partial_x\varphi\big)\, dx\, dt = \mathcal{O}(\varepsilon). \tag{145}$$

Because the $\tanh$ profile guarantees $\widehat{u}^\tau \in L^\infty \cap \mathrm{BV}$, the dominated convergence theorem allows the limit $\varepsilon \to 0$ to be taken inside the integral. **This step is particularly emphasized: the local monotonicity induced by parameterization** (5) **is the key topological constraint that renders this weak limit legitimate.** Traditional diffusion models (e.g., those forcing a Gaussian fit across shock interfaces) inevitably produce Gibbs oscillations, causing the BV norm to blow up in the limit and precluding application of dominated convergence. The profile (144) converges strongly to the piecewise-constant function

$$\mathbf{u}^\star(x,t) = \begin{cases} u_L, & x < x_s(t), \\ u_R, & x > x_s(t). \end{cases} \tag{146}$$

Partition the spacetime domain along the shock trajectory $\Gamma : x = x_s(t)$ into a left region D^- ($x < x_s$) and a right region D^+ ($x > x_s$). Then

$$0 = \lim_{\varepsilon\to 0} I(\varepsilon) = \iint_{D^-} \big(u_L\, \partial_t\varphi + f(u_L)\, \partial_x\varphi\big)\, dx\, dt + \iint_{D^+} \big(u_R\, \partial_t\varphi + f(u_R)\, \partial_x\varphi\big)\, dx\, dt. \tag{147}$$

Inside each $D^\pm$, the solution u_L, u_R is a strong solution of the conservation law. Applying Green's theorem to transfer derivatives from φ onto u,

$$\begin{aligned} \iint_D \big(u\, \partial_t\varphi + f(u)\, \partial_x\varphi\big)\, dx\, dt = &\int_{\partial D} \varphi\, \big(u\, n_t + f(u)\, n_x\big)\, ds \\ &- \iint_D \varphi\, \big(\partial_t u + \partial_x f(u)\big)\, dx\, dt. \end{aligned} \tag{148}$$

Since the PDE holds pointwise in $D^\pm$, the area term vanishes. Because φ has compact support, only the boundaries along Γ contribute. On Γ, the outward normal (pointing toward D^+) is

$$\mathbf{n} = (n_x, n_t) = \frac{1}{\sqrt{1 + \dot{x}_s^2}}\big(1,\ -\dot{x}_s\big). \tag{149}$$

Accounting for orientation, the jump condition emerges:

$$\int_\Gamma \varphi \cdot \Big[\big(u_L(-\dot{x}_s) + f(u_L)\big) - \big(u_R(-\dot{x}_s) + f(u_R)\big)\Big] \frac{1}{\sqrt{1 + \dot{x}_s^2}}\, ds = 0. \tag{150}$$

Since φ is arbitrary along Γ, the algebraic bracket must vanish identically, yielding

$$\dot{x}_s\,(u_L - u_R) = f(u_L) - f(u_R). \tag{151}$$

For a genuine shock $u_L \neq u_R$, division by the difference gives the Rankine–Hugoniot condition

$$\dot{x}_s(t) = \frac{f(u_L) - f(u_R)}{u_L - u_R}. \qquad \square \tag{152}$$

Part III: Entropy compatibility and the Lax condition.

Background: the non-uniqueness of weak solutions. The Rankine–Hugoniot condition derived above is a *necessary* but not sufficient condition for a weak solution. Physically inadmissible "expansion shocks", which dissipate energy in an entropy-violating manner, also satisfy the R–H condition in nonlinear evolution models. Only by imposing a rigorous entropy condition—the Lax entropy condition, motivated by physical mechanisms such as the Double-Burgers coupling—do we ensure that the generated solution is not merely a "numerical fit" to data, but a genuine approximation of the *physically correct* conservation law.

3.1 Divergence–characteristic relation.

For a convex flux f (e.g., the Burgers flux $f(u) = u^2/2$), a physical shock must satisfy the Lax entropy condition

$$f'(u_L) > \dot{x}_s > f'(u_R). \tag{153}$$

In the reverse-time generative ODE,

$$\frac{du}{d\tau} = -\frac{1}{2}\,\beta(\tau)\big[u + s_\theta(u,\tau)\big], \tag{154}$$

Tweedie reconstruction yields $\partial_x \widehat{u} \approx \partial_x(u + \sigma(\tau)^2\, s_\theta) \to -\infty$ near x_s. This singular negative gradient corresponds to the compression of characteristics toward the shock in physical space.

3.2 Kruzhkov entropy inequality.

To prove that $\widehat{u}$ is an entropy solution, we verify the Kruzhkov inequality for every constant k (in the sense of distributions):

$$\partial_t\,|\widehat{u} - k| + \partial_x\big[\mathrm{sgn}(\widehat{u} - k)\,(f(\widehat{u}) - f(k))\big] \leq 0. \tag{155}$$

By Theorem 1 (Double-Burgers coupling), the score field itself satisfies a viscous Burgers equation. Hence the generative dynamics driven by s_θ constitute a JKO gradient flow that minimizes an energy functional. A jump $u_L \to u_R$ violating the Lax condition (i.e., $u_L < u_R$) would correspond to an expansion (rarefaction) fan whose characteristics diverge outward. Under the BV-aware parameterization (5), the sign of the score gradient is determined by the $\tanh$ profile.

3.3 Proof by contradiction.

Suppose a non-physical shock (an expansion shock) satisfies the R–H condition but has $u_L < u_R$.

- **(i) Score behavior.** The Tweedie formula then requires the score s_θ to supply an outward "push" in order to sustain this unstable discontinuity.
- **(ii) Burgers consistency loss.** The loss term $\mathcal{L}_{\mathrm{Burg}}$ (Burgers consistency) enforces $\partial_\tau s + (s \cdot \nabla)s = \nu_{\mathrm{phys}}\,\Delta s$. An expansion jump would be immediately regularized by the diffusion term Δs, producing a large residual in $\mathcal{L}_{\mathrm{Burg}}$.
- **(iii) Conclusion.** Minimizing the total loss systematically eliminates all non-physical jumps, because only compressive configurations satisfying the Lax condition (153) are compatible with the stable evolution operator of $\mathcal{L}_{\mathrm{Burg}}$.

3.4 Stability in the $\tau \to 0$ limit.

Finally, the Oleinik entropy condition provides the estimate near the shock:

$$\frac{\widehat{u}(x+a,t) - \widehat{u}(x,t)}{a} \leq \frac{C}{t}, \tag{156}$$

which is guaranteed by the controlled width of the $\tanh$ layer in parameterization (5). As τ decreases, the profile steepens but remains strictly monotone whenever $u_L > u_R$, thereby precluding non-physical oscillations.

Combining the three parts above, the generated solution satisfies both the Rankine–Hugoniot condition and the Lax entropy condition almost everywhere. □

A.10 Proof of Theorem 29 (JKO correspondence)

Theorem 29 (JKO correspondence)**.** *Let p_τ denote the density of the reverse sampler at diffusion time $\tau \in [0, T]$, and let $\rho^\star$ be the target distribution induced by the PDE solution in data space. Define the free energy functional*

$$\mathcal{F}(\rho) = \mathrm{Ent}(\rho \mid \rho^\star) + \lambda\, \mathcal{R}(\rho), \qquad \mathrm{Ent}(\rho \mid \rho^\star) = \int_{\mathbb{R}^N} \rho(x)\, \log \frac{\rho(x)}{\rho^\star(x)}\, dx, \tag{157}$$

where $\mathcal{R}$ is a density functional encoding the PDE constraint and $\lambda \geq 0$.

Assume the following hypotheses:

1. ***(Exact score limit.)*** *After sufficient training, $s_\theta(x, \tau) = \nabla_x \log p_\tau(x)$ pointwise.*
2. ***(Viscosity-matched schedule.)*** *$a(\tau) = g(\tau)^2/2 > 0$, aligned with the physical viscosity.*
3. ***(PDE constraint as a differentiable functional.)*** *$\mathcal{R}$ is Fréchet-differentiable on $\mathcal{P}_2(\mathbb{R}^N)$, with first variation $\delta\mathcal{R}/\delta\rho$ well-defined; $\nabla_x(\delta\mathcal{R}/\delta\rho)$ is locally bounded.*
4. ***(Rigid $\mathcal{L}_{\mathrm{Burg}}$ consistency.)*** *The Burgers consistency penalty $\mathcal{L}_{\mathrm{Burg}}$ is sufficiently strong that, in the infinite-capacity noiseless limit, the learned score field satisfies the viscous Burgers structure*
$$\partial_\tau s_\tau + 2\, s_\tau \cdot \nabla\, s_\tau = \Delta s_\tau. \tag{158}$$
Moreover, $p_\tau \in \mathcal{P}_2(\mathbb{R}^N)$ is smooth in (x, τ) and strictly positive ($p_\tau > 0$).
5. ***(PDE guidance as a gradient drift.)*** *In the continuous-time limit, the combined PDE constraint losses $\mathcal{L}_{\mathrm{ent}}$ (Kruzhkov entropy penalty) and $\mathcal{L}_{\mathrm{Burg}}$ (Burgers consistency) induce an additional guidance velocity $g_\tau(x)$ in the reverse ODE that converges to the spatial gradient of a functional. Specifically, the total effective velocity field of the EntroDiff reverse sampler is*
$$v_\tau(x) = -a(\tau)\Big[\nabla_x \log p_\tau(x) + \nabla_x V(x) + \lambda\, \nabla_x \frac{\delta\mathcal{R}}{\delta\rho}(p_\tau)(x)\Big], \tag{159}$$
where V is the potential of the target distribution ($\rho^\star \propto e^{-V}$). The guidance contributions ∇V and $\lambda\nabla(\delta\mathcal{R}/\delta\rho)$ arise from the gradient of the PDE losses with respect to the sample trajectory, which by design converge to these functional-gradient limits when the losses are driven to zero in training.

Then the density evolution of the reverse sampler satisfies

$$\partial_\tau p_\tau = \nabla \cdot \Big(a(\tau)\, p_\tau\, \nabla \frac{\delta\mathcal{F}}{\delta\rho}(p_\tau)\Big). \tag{160}$$

Equivalently, under the time reparameterization $\theta(\tau) = \int_0^\tau a(r)\, dr$, we have

$$\partial_\theta p_\theta = \nabla \cdot \Big(p_\theta\, \nabla \frac{\delta\mathcal{F}}{\delta p}(p_\theta)\Big), \tag{161}$$

which is precisely the W_2-gradient flow of $\mathcal{F}$. Consequently, by the JKO theory, a single time-discrete step takes the variational form

$$p^{n+1} = \arg\min_{p \in \mathcal{P}_2(\mathbb{R}^N)} \Big\{\mathcal{F}(p) + \frac{1}{2\Delta\theta}\, W_2^2(p, p^n)\Big\}. \tag{162}$$

Proof. **Step 1: Reverse sampler as a continuity equation.**

Let X_τ denote the particle trajectory of the reverse sampler. If its continuous-time limit satisfies the ODE

$$\frac{dX_\tau}{d\tau} = v_\tau(X_\tau), \tag{163}$$

then by the standard pushforward formula, the particle distribution $p_\tau = \mathrm{Law}(X_\tau)$ satisfies the continuity equation

$$\partial_\tau p_\tau + \nabla \cdot (p_\tau\, v_\tau) = 0. \tag{164}$$

Thus, identifying the velocity field v_τ is the central task.

For any test function $\varphi \in C_c^\infty(\mathbb{R}^N)$, the identity follows from the chain rule:

$$\begin{aligned}\frac{d}{d\tau}\int_{\mathbb{R}^N} \varphi(x)\, p_\tau(x)\, dx &= \frac{d}{d\tau}\,\mathbb{E}[\varphi(X_\tau)] \\ &= \mathbb{E}[\nabla\varphi(X_\tau)\cdot v_\tau(X_\tau)] \\ &= \int \nabla\varphi(x)\cdot v_\tau(x)\, p_\tau(x)\, dx.\end{aligned} \tag{165}$$

Integration by parts of the right-hand side yields

$$\frac{d}{d\tau}\int \varphi\, p_\tau\, dx = -\int \varphi(x)\, \nabla\cdot(p_\tau\, v_\tau)(x)\, dx, \tag{166}$$

and since φ is arbitrary, the continuity equation (164) holds in the sense of distributions (and pointwise under the smoothness of p_τ and v_τ).

Step 2: Score drift and PDE guidance — the total effective velocity.

We identify v_τ as the sum of two contributions.

(a) Score drift (from denoising score matching). Under Hypothesis 1 (exact score) and Hypothesis 4 (rigid $\mathcal{L}_{\mathrm{Burg}}$), the network score converges to the true score:

$$s_\theta(x,\tau) = \nabla_x \log p_\tau(x). \tag{167}$$

The probability flow ODE induced by score matching alone has the velocity

$$v_\tau^{\mathrm{score}}(x) = -a(\tau)\, s_\theta(x,\tau) = -a(\tau)\, \nabla_x \log p_\tau(x). \tag{168}$$

(b) PDE guidance drift (from constraint losses). The EntroDiff loss family includes PDE-constraint terms beyond denoising score matching: $\mathcal{L}_{\mathrm{ent}}$ (Kruzhkov entropy penalty) and $\mathcal{L}_{\mathrm{Burg}}$ (Burgers consistency). These losses are functionals of the generated trajectory that penalize deviations from physical admissibility. In the continuous-time and infinite-data limit, their gradients with respect to the sample position induce an additional drift $g_\tau(x)$.

By Hypothesis 5, as the constraint losses are driven to zero, this guidance drift converges to the gradient of the potential that defines the target distribution plus the variational gradient of the PDE-constraint functional:

$$g_\tau(x) = -\Big(\nabla_x V(x) + \lambda\, \nabla_x \frac{\delta\mathcal{R}}{\delta\rho}(p_\tau)(x)\Big). \tag{169}$$

Here $\nabla V = -\nabla \log \rho^\star$ is the drift toward the target $\rho^\star \propto e^{-V}$, and $\lambda\nabla(\delta\mathcal{R}/\delta\rho)$ is the PDE-constraint drift (see Step 4 for the construction of $\mathcal{R}$).

(c) Total effective velocity. Summing (168) and (169), and absorbing the common factor $-a(\tau)$,

$$v_\tau(x) = -a(\tau)\Big[\nabla_x \log p_\tau(x) + \nabla_x V(x) + \lambda\, \nabla_x \frac{\delta\mathcal{R}}{\delta\rho}(p_\tau)(x)\Big]. \tag{170}$$

Substituting into the continuity equation gives

$$\partial_\tau p_\tau = \nabla\cdot\Big(a(\tau)\, p_\tau\, \big[\nabla\log p_\tau + \nabla V + \lambda\nabla\tfrac{\delta\mathcal{R}}{\delta\rho}\big]\Big). \tag{171}$$

We now show that the bracketed quantity in (170) is exactly $\nabla(\delta\mathcal{F}/\delta\rho)$.

Step 3: First variation of the entropy term.

Consider the relative entropy

$$\mathrm{Ent}(p \mid \rho^\star) = \int p \log \frac{p}{\rho^\star}\, dx. \tag{172}$$

Take a perturbation $p_\varepsilon = p + \varepsilon\eta$ with $\int \eta\, dx = 0$. A first-order expansion gives

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathrm{Ent}(p_\varepsilon \mid \rho^\star) = \int \eta(x)\Big(\log \frac{p(x)}{\rho^\star(x)} + 1\Big)\, dx. \tag{173}$$

Hence the L^2 first variation is

$$\frac{\delta}{\delta p}\,\mathrm{Ent}(p \mid \rho^\star) = \log \frac{p}{\rho^\star} + 1 \tag{174}$$

(the additive constant does not affect the gradient flow). Taking the spatial gradient,

$$\nabla \frac{\delta}{\delta p}\,\mathrm{Ent}(p \mid \rho^\star) = \nabla \log p - \nabla \log \rho^\star. \tag{175}$$

Since $\rho^\star \propto e^{-V}$, we have $-\nabla \log \rho^\star = \nabla V$, and therefore

$$\nabla \frac{\delta}{\delta p}\,\mathrm{Ent}(p \mid \rho^\star) = \nabla \log p + \nabla V. \tag{176}$$

Step 4: PDE constraint as a differentiable functional.

The PDE constraint (beyond $\mathcal{L}_{\mathrm{Burg}}$) must be expressible as a density functional $\mathcal{R}(\rho)$. Under Hypothesis 3, $\mathcal{R}$ is Fréchet-differentiable on $\mathcal{P}_2$, so its first variation $\frac{\delta\mathcal{R}}{\delta\rho}$ exists and satisfies

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathcal{R}(\rho_\varepsilon) = \int \frac{\delta\mathcal{R}}{\delta\rho}(\rho)\, \eta\, dx \tag{177}$$

for any zero-mass perturbation η.

In the Wasserstein setting, perturbations are generated by a transport field ξ via the pushforward:

$$T_\varepsilon(x) = x + \varepsilon\, \xi(x), \qquad \rho_\varepsilon = (T_\varepsilon)_{\#}\, \rho. \tag{178}$$

The corresponding variational formula is

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathcal{R}\big((\mathrm{Id} + \varepsilon\xi)_{\#}\rho\big) = \int \nabla \frac{\delta\mathcal{R}}{\delta\rho}(\rho) \cdot \xi\, \rho\, dx. \tag{179}$$

Thus, the $\mathcal{R}$ gradient flow term contributes a drift $-\nabla(\delta\mathcal{R}/\delta\rho)$ in the velocity field (see (169)).

Step 5: Identification with the free energy gradient.

Combining the entropy and PDE-constraint terms, the free energy is

$$\mathcal{F}(\rho) = \mathrm{Ent}(\rho \mid \rho^\star) + \lambda\, \mathcal{R}(\rho). \tag{180}$$

From (176) and Step 4, its spatial variational gradient is

$$\nabla \frac{\delta\mathcal{F}}{\delta\rho} = \nabla \frac{\delta}{\delta\rho}\,\mathrm{Ent}(\rho \mid \rho^\star) + \lambda\, \nabla \frac{\delta\mathcal{R}}{\delta\rho} = \nabla \log \rho + \nabla V + \lambda\, \nabla \frac{\delta\mathcal{R}}{\delta\rho}. \tag{181}$$

Comparing (170) with (181), the total effective velocity is exactly

$$v_\tau(x) = -a(\tau)\, \nabla_x \frac{\delta\mathcal{F}}{\delta\rho}(p_\tau)(x). \tag{182}$$

Substituting into the continuity equation (Step 1),

$$\partial_\tau p_\tau + \nabla \cdot \Big(p_\tau \Big[-a(\tau)\, \nabla \frac{\delta\mathcal{F}}{\delta\rho}(p_\tau)\Big]\Big) = 0, \tag{183}$$

i.e.,

$$\partial_\tau p_\tau = \nabla \cdot \Big(a(\tau)\, p_\tau\, \nabla \frac{\delta \mathcal{F}}{\delta \rho}(p_\tau) \Big). \tag{184}$$

To eliminate the time-weight $a(\tau)$, define the reparameterized time

$$\theta(\tau) = \int_0^\tau a(r)\, dr. \tag{185}$$

Since $a(\tau) > 0$, θ is strictly increasing and locally invertible. By the chain rule,

$$\partial_\tau p_\tau = \frac{d\theta}{d\tau}\, \partial_\theta p_\theta = a(\tau)\, \partial_\theta p_\theta. \tag{186}$$

Cancelling $a(\tau)$ yields the canonical Wasserstein-2 gradient flow form

$$\partial_\theta p_\theta = \nabla \cdot \Big(p_\theta\, \nabla \frac{\delta \mathcal{F}}{\delta p}(p_\theta) \Big). \tag{187}$$

This completes the bridge from the EntroDiff reverse sampler to an exact Wasserstein gradient flow. □

Step 6: JKO discretization from the Wasserstein gradient flow.

We now show that the implicit Euler discretization of the above gradient flow is equivalent to the JKO variational scheme.

6.1 Implicit Euler discretization. Consider a uniform time grid with step $\Delta\theta$ (in reparameterized time). Given $p^n \approx p_{\theta_n}$, the implicit Euler discretization of the gradient flow is

$$\frac{p^{n+1} - p^n}{\Delta\theta} = \nabla \cdot \Big(p^{n+1}\, \nabla \frac{\delta \mathcal{F}}{\delta \rho}(p^{n+1}) \Big). \tag{188}$$

6.2 Proximal step analogy. In Euclidean space, the implicit Euler step for $\dot{x} = -\nabla F(x)$ is equivalent to the proximal minimization

$$x^{n+1} = \arg\min_x \Big\{ F(x) + \frac{1}{2\Delta t}\, \|x - x^n\|^2 \Big\}. \tag{189}$$

The Wasserstein analogue replaces the Euclidean distance with W_2 and the Euclidean gradient with the continuity equation structure. The claim is that

$$p^{n+1} = \arg\min_{p \in \mathcal{P}_2(\mathbb{R}^N)} \Big\{ \mathcal{F}(p) + \frac{1}{2\Delta\theta}\, W_2^2(p, p^n) \Big\} \tag{190}$$

is the JKO step corresponding to the implicit Euler discretization above.

6.3 Variational derivation. Define the augmented functional

$$\mathcal{J}(p) = \mathcal{F}(p) + \frac{1}{2\Delta\theta}\, W_2^2(p, p^n), \tag{191}$$

and let p^{n+1} be a minimizer. In Wasserstein space, admissible variations are generated by transport maps rather than additive perturbations. Define the perturbation

$$T_\varepsilon(x) = x + \varepsilon\, \xi(x), \qquad p_\varepsilon = (T_\varepsilon)_\#\, p^{n+1}, \tag{192}$$

where ξ is an arbitrary smooth, compactly supported vector field. Optimality requires

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathcal{J}(p_\varepsilon) = 0. \tag{193}$$

Variation of $\mathcal{F}$. By the Wasserstein variational formula,

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathcal{F}(p_\varepsilon) = \int \nabla \frac{\delta \mathcal{F}}{\delta \rho}(p^{n+1}) \cdot \xi\, p^{n+1}\, dx. \tag{194}$$

Variation of the W_2^2 term. Let π be the optimal transport plan from p^{n+1} to p^n, with Kantorovich potential ϕ. A classical result of optimal transport theory gives

$$\frac{\delta}{\delta\rho}\Big(\frac{1}{2}\,W_2^2(\rho, p^n)\Big) = \phi, \tag{195}$$

and therefore

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0}\frac{1}{2}\,W_2^2(p_\varepsilon, p^n) = \int \nabla\phi\cdot\xi\; p^{n+1}\,dx. \tag{196}$$

First-order condition. Combining both terms,

$$\int\Big[\nabla\frac{\delta\mathcal{F}}{\delta\rho}(p^{n+1}) + \frac{1}{\Delta\theta}\,\nabla\phi\Big]\cdot\xi\; p^{n+1}\,dx = 0. \tag{197}$$

Since ξ is arbitrary, the integrand must vanish p^{n+1}-almost everywhere, yielding

$$\nabla\phi = -\Delta\theta\,\nabla\frac{\delta\mathcal{F}}{\delta\rho}(p^{n+1}). \tag{198}$$

Connection to the optimal transport map. The Brenier theorem gives the optimal transport map T from p^{n+1} to p^n as $T(x) = x - \nabla\phi(x)$. Substituting the expression for $\nabla\phi$,

$$T(x) = x + \Delta\theta\,\nabla\frac{\delta\mathcal{F}}{\delta\rho}(p^{n+1}). \tag{199}$$

Deriving the discrete continuity equation. By the pushforward relation $p^n = T_\# p^{n+1}$, we expand for small $\Delta\theta$ (using the Jacobian expansion of the pushforward):

$$p^n(x) = p^{n+1}(x) - \Delta\theta\,\nabla\cdot\Big(p^{n+1}\,\nabla\frac{\delta\mathcal{F}}{\delta\rho}(p^{n+1})\Big) + o(\Delta\theta). \tag{200}$$

Rearranging,

$$\frac{p^{n+1}-p^n}{\Delta\theta} = \nabla\cdot\Big(p^{n+1}\,\nabla\frac{\delta\mathcal{F}}{\delta\rho}(p^{n+1})\Big) + o(1). \tag{201}$$

Taking $\Delta\theta\to 0$, we recover exactly the implicit Euler discretization (188) of the Wasserstein gradient flow.

6.4 Conclusion of the JKO derivation. Thus, a minimizer p^{n+1} of the JKO functional satisfies the implicit Euler discretization of the Wasserstein gradient flow. The reverse sampler, whose density evolution was established in Steps 1–5 as the gradient flow of $\mathcal{F}$, therefore admits the JKO variational characterization in the discrete-time limit $\Delta\tau\to 0$. This completes the proof of Theorem 5. □

Compact Fokker–Planck expansion.

For completeness, expanding the gradient flow structurally: using $\nabla\log p_\theta = \nabla p_\theta/p_\theta$ and the explicit form of the entropy variation from Step 3, the gradient flow PDE becomes

$$\partial_\theta p_\theta = \Delta p_\theta + \nabla\cdot(p_\theta\nabla V) + \lambda\,\nabla\cdot\Big(p_\theta\,\nabla\frac{\delta\mathcal{R}}{\delta\rho}\Big), \tag{202}$$

exhibiting the standard decomposition into a diffusion term (Δp_θ), an external potential drift ($\nabla\cdot(p_\theta\nabla V)$), and a PDE-constraint transport term.

Remark (Relation to the Obukhov–Fokker–Planck correspondence).

The identification (182) can be understood through the lens of entropy-dissipation balances. The pure reverse probability-flow ODE ($v = -a\nabla\log p$) is the Wasserstein gradient flow of the negative entropy $H(p) = \int p\log p$. Adding the PDE guidance terms $\nabla V + \lambda\nabla(\delta\mathcal{R}/\delta\rho)$ shifts the stationary point from the standard Gaussian to the target $\rho^\star$ while incorporating the physical constraint $\mathcal{R}$. The resulting free energy $\mathcal{F} = \mathrm{Ent}(\cdot\mid\rho^\star) + \lambda\mathcal{R}$ is the natural Lyapunov functional for the guided dynamics, and the JKO scheme is its canonical time discretization [**?**].

B Additional Experiments

B.1 E4 · 2D Shallow-water (rebuttal)

TODO: If time permits before rebuttal: hydraulic jump benchmark.

B.2 E5 · 1D Vlasov–Poisson (rebuttal)

TODO: Cold-beam instability with weak-solution filamentation. Compare against Deep Kinetic JKO [14].

B.3 Extended ablations

TODO: Ablate noise schedule, parameterization, loss-family components, and λ hyperparameters individually.

C Implementation Details

C.1 Network architecture

TODO: Detail ϕ^{sm}, ϕ^{sh}, κ networks. Sizes, activations, normalizations.

C.2 Training hyperparameters

TODO: Optimizer (AdamW), batch sizes, learning rates, EMA, training horizon, λ_{ent}, λ_{BV}, λ_{Burg}.

C.3 Sampling hyperparameters

TODO: Number of Heun steps, ζ_{obs}, ζ_{PDE} schedules.

C.4 Compute and reproducibility

TODO: GPU type, runtime per benchmark, anonymized code repository link in the supplementary material.

NeurIPS Paper Checklist

The checklist is designed to encourage best practices for responsible machine learning research, addressing issues of reproducibility, transparency, research ethics, and societal impact. Do not remove the checklist: **The papers not including the checklist will be desk rejected.** The checklist should follow the references and follow the (optional) supplemental material. The checklist does NOT count towards the page limit.

Please read the checklist guidelines carefully for information on how to answer these questions. For each question in the checklist:

- You should answer [Yes], [No], or [N/A].
- [N/A] means either that the question is Not Applicable for that particular paper or the relevant information is Not Available.
- Please provide a short (1–2 sentence) justification right after your answer (even for [N/A]).

The checklist answers are an integral part of your paper submission. They are visible to the reviewers, area chairs, senior area chairs, and ethics reviewers. You will also be asked to include it (after eventual revisions) with the final version of your paper, and its final version will be published with the paper.

The reviewers of your paper will be asked to use the checklist as one of the factors in their evaluation. While [Yes] is generally preferable to [No], it is perfectly acceptable to answer [No] provided a proper justification is given (e.g., error bars are not reported because it would be too computationally expensive" or "we were unable to find the license for the dataset we used"). In general, answering [No] or [N/A] is not grounds for rejection. While the questions are phrased in a binary way, we acknowledge that the true answer is often more nuanced, so please just use your best judgment and write a justification to elaborate. All supporting evidence can appear either in the main paper or the supplemental material, provided in appendix. If you answer [Yes] to a question, in the justification please point to the section(s) where related material for the question can be found.

IMPORTANT, please:

- **Delete this instruction block, but keep the section heading "NeurIPS Paper Checklist"**,
- **Keep the checklist subsection headings, questions/answers and guidelines below.**
- **Do not modify the questions and only use the provided macros for your answers**.

1. **Claims**

   Question: Do the main claims made in the abstract and introduction accurately reflect the paper's contributions and scope?

   Answer: **[TODO]**

   Justification: **[TODO]**

   Guidelines:

   - The answer [N/A] means that the abstract and introduction do not include the claims made in the paper.
   - The abstract and/or introduction should clearly state the claims made, including the contributions made in the paper and important assumptions and limitations. A [No] or [N/A] answer to this question will not be perceived well by the reviewers.
   - The claims made should match theoretical and experimental results, and reflect how much the results can be expected to generalize to other settings.
   - It is fine to include aspirational goals as motivation as long as it is clear that these goals are not attained by the paper.

2. **Limitations**

   Question: Does the paper discuss the limitations of the work performed by the authors?

   Answer: **[TODO]**

   Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper has no limitation while the answer [No] means that the paper has limitations, but those are not discussed in the paper.
- The authors are encouraged to create a separate "Limitations" section in their paper.
- The paper should point out any strong assumptions and how robust the results are to violations of these assumptions (e.g., independence assumptions, noiseless settings, model well-specification, asymptotic approximations only holding locally). The authors should reflect on how these assumptions might be violated in practice and what the implications would be.
- The authors should reflect on the scope of the claims made, e.g., if the approach was only tested on a few datasets or with a few runs. In general, empirical results often depend on implicit assumptions, which should be articulated.
- The authors should reflect on the factors that influence the performance of the approach. For example, a facial recognition algorithm may perform poorly when image resolution is low or images are taken in low lighting. Or a speech-to-text system might not be used reliably to provide closed captions for online lectures because it fails to handle technical jargon.
- The authors should discuss the computational efficiency of the proposed algorithms and how they scale with dataset size.
- If applicable, the authors should discuss possible limitations of their approach to address problems of privacy and fairness.
- While the authors might fear that complete honesty about limitations might be used by reviewers as grounds for rejection, a worse outcome might be that reviewers discover limitations that aren't acknowledged in the paper. The authors should use their best judgment and recognize that individual actions in favor of transparency play an important role in developing norms that preserve the integrity of the community. Reviewers will be specifically instructed to not penalize honesty concerning limitations.

3. **Theory assumptions and proofs**

Question: For each theoretical result, does the paper provide the full set of assumptions and a complete (and correct) proof?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper does not include theoretical results.
- All the theorems, formulas, and proofs in the paper should be numbered and cross-referenced.
- All assumptions should be clearly stated or referenced in the statement of any theorems.
- The proofs can either appear in the main paper or the supplemental material, but if they appear in the supplemental material, the authors are encouraged to provide a short proof sketch to provide intuition.
- Inversely, any informal proof provided in the core of the paper should be complemented by formal proofs provided in appendix or supplemental material.
- Theorems and Lemmas that the proof relies upon should be properly referenced.

4. **Experimental result reproducibility**

Question: Does the paper fully disclose all the information needed to reproduce the main experimental results of the paper to the extent that it affects the main claims and/or conclusions of the paper (regardless of whether the code and data are provided or not)?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper does not include experiments.

- If the paper includes experiments, a [No] answer to this question will not be perceived well by the reviewers: Making the paper reproducible is important, regardless of whether the code and data are provided or not.
- If the contribution is a dataset and/or model, the authors should describe the steps taken to make their results reproducible or verifiable.
- Depending on the contribution, reproducibility can be accomplished in various ways. For example, if the contribution is a novel architecture, describing the architecture fully might suffice, or if the contribution is a specific model and empirical evaluation, it may be necessary to either make it possible for others to replicate the model with the same dataset, or provide access to the model. In general. releasing code and data is often one good way to accomplish this, but reproducibility can also be provided via detailed instructions for how to replicate the results, access to a hosted model (e.g., in the case of a large language model), releasing of a model checkpoint, or other means that are appropriate to the research performed.
- While NeurIPS does not require releasing code, the conference does require all submissions to provide some reasonable avenue for reproducibility, which may depend on the nature of the contribution. For example
  (a) If the contribution is primarily a new algorithm, the paper should make it clear how to reproduce that algorithm.
  (b) If the contribution is primarily a new model architecture, the paper should describe the architecture clearly and fully.
  (c) If the contribution is a new model (e.g., a large language model), then there should either be a way to access this model for reproducing the results or a way to reproduce the model (e.g., with an open-source dataset or instructions for how to construct the dataset).
  (d) We recognize that reproducibility may be tricky in some cases, in which case authors are welcome to describe the particular way they provide for reproducibility. In the case of closed-source models, it may be that access to the model is limited in some way (e.g., to registered users), but it should be possible for other researchers to have some path to reproducing or verifying the results.

5. **Open access to data and code**

Question: Does the paper provide open access to the data and code, with sufficient instructions to faithfully reproduce the main experimental results, as described in supplemental material?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that paper does not include experiments requiring code.
- Please see the NeurIPS code and data submission guidelines (`https://neurips.cc/public/guides/CodeSubmissionPolicy`) for more details.
- While we encourage the release of code and data, we understand that this might not be possible, so [No] is an acceptable answer. Papers cannot be rejected simply for not including code, unless this is central to the contribution (e.g., for a new open-source benchmark).
- The instructions should contain the exact command and environment needed to run to reproduce the results. See the NeurIPS code and data submission guidelines (`https://neurips.cc/public/guides/CodeSubmissionPolicy`) for more details.
- The authors should provide instructions on data access and preparation, including how to access the raw data, preprocessed data, intermediate data, and generated data, etc.
- The authors should provide scripts to reproduce all experimental results for the new proposed method and baselines. If only a subset of experiments are reproducible, they should state which ones are omitted from the script and why.
- At submission time, to preserve anonymity, the authors should release anonymized versions (if applicable).

- Providing as much information as possible in supplemental material (appended to the paper) is recommended, but including URLs to data and code is permitted.

6. **Experimental setting/details**

   Question: Does the paper specify all the training and test details (e.g., data splits, hyperparameters, how they were chosen, type of optimizer) necessary to understand the results?

   Answer: **[TODO]**

   Justification: **[TODO]**

   Guidelines:

   - The answer [N/A] means that the paper does not include experiments.
   - The experimental setting should be presented in the core of the paper to a level of detail that is necessary to appreciate the results and make sense of them.
   - The full details can be provided either with the code, in appendix, or as supplemental material.

7. **Experiment statistical significance**

   Question: Does the paper report error bars suitably and correctly defined or other appropriate information about the statistical significance of the experiments?

   Answer: **[TODO]**

   Justification: **[TODO]**

   Guidelines:

   - The answer [N/A] means that the paper does not include experiments.
   - The authors should answer [Yes] if the results are accompanied by error bars, confidence intervals, or statistical significance tests, at least for the experiments that support the main claims of the paper.
   - The factors of variability that the error bars are capturing should be clearly stated (for example, train/test split, initialization, random drawing of some parameter, or overall run with given experimental conditions).
   - The method for calculating the error bars should be explained (closed form formula, call to a library function, bootstrap, etc.)
   - The assumptions made should be given (e.g., Normally distributed errors).
   - It should be clear whether the error bar is the standard deviation or the standard error of the mean.
   - It is OK to report 1-sigma error bars, but one should state it. The authors should preferably report a 2-sigma error bar than state that they have a 96% CI, if the hypothesis of Normality of errors is not verified.
   - For asymmetric distributions, the authors should be careful not to show in tables or figures symmetric error bars that would yield results that are out of range (e.g., negative error rates).
   - If error bars are reported in tables or plots, the authors should explain in the text how they were calculated and reference the corresponding figures or tables in the text.

8. **Experiments compute resources**

   Question: For each experiment, does the paper provide sufficient information on the computer resources (type of compute workers, memory, time of execution) needed to reproduce the experiments?

   Answer: **[TODO]**

   Justification: **[TODO]**

   Guidelines:

   - The answer [N/A] means that the paper does not include experiments.
   - The paper should indicate the type of compute workers CPU or GPU, internal cluster, or cloud provider, including relevant memory and storage.
   - The paper should provide the amount of compute required for each of the individual experimental runs as well as estimate the total compute.

- The paper should disclose whether the full research project required more compute than the experiments reported in the paper (e.g., preliminary or failed experiments that didn't make it into the paper).

9. **Code of ethics**

Question: Does the research conducted in the paper conform, in every respect, with the NeurIPS Code of Ethics `https://neurips.cc/public/EthicsGuidelines`?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the authors have not reviewed the NeurIPS Code of Ethics.
- If the authors answer [No], they should explain the special circumstances that require a deviation from the Code of Ethics.
- The authors should make sure to preserve anonymity (e.g., if there is a special consideration due to laws or regulations in their jurisdiction).

10. **Broader impacts**

Question: Does the paper discuss both potential positive societal impacts and negative societal impacts of the work performed?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that there is no societal impact of the work performed.
- If the authors answer [N/A] or [No], they should explain why their work has no societal impact or why the paper does not address societal impact.
- Examples of negative societal impacts include potential malicious or unintended uses (e.g., disinformation, generating fake profiles, surveillance), fairness considerations (e.g., deployment of technologies that could make decisions that unfairly impact specific groups), privacy considerations, and security considerations.
- The conference expects that many papers will be foundational research and not tied to particular applications, let alone deployments. However, if there is a direct path to any negative applications, the authors should point it out. For example, it is legitimate to point out that an improvement in the quality of generative models could be used to generate Deepfakes for disinformation. On the other hand, it is not needed to point out that a generic algorithm for optimizing neural networks could enable people to train models that generate Deepfakes faster.
- The authors should consider possible harms that could arise when the technology is being used as intended and functioning correctly, harms that could arise when the technology is being used as intended but gives incorrect results, and harms following from (intentional or unintentional) misuse of the technology.
- If there are negative societal impacts, the authors could also discuss possible mitigation strategies (e.g., gated release of models, providing defenses in addition to attacks, mechanisms for monitoring misuse, mechanisms to monitor how a system learns from feedback over time, improving the efficiency and accessibility of ML).

11. **Safeguards**

Question: Does the paper describe safeguards that have been put in place for responsible release of data or models that have a high risk for misuse (e.g., pre-trained language models, image generators, or scraped datasets)?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper poses no such risks.

- Released models that have a high risk for misuse or dual-use should be released with necessary safeguards to allow for controlled use of the model, for example by requiring that users adhere to usage guidelines or restrictions to access the model or implementing safety filters.
- Datasets that have been scraped from the Internet could pose safety risks. The authors should describe how they avoided releasing unsafe images.
- We recognize that providing effective safeguards is challenging, and many papers do not require this, but we encourage authors to take this into account and make a best faith effort.

12. **Licenses for existing assets**

    Question: Are the creators or original owners of assets (e.g., code, data, models), used in the paper, properly credited and are the license and terms of use explicitly mentioned and properly respected?

    Answer: **[TODO]**

    Justification: **[TODO]**

    Guidelines:

    - The answer [N/A] means that the paper does not use existing assets.
    - The authors should cite the original paper that produced the code package or dataset.
    - The authors should state which version of the asset is used and, if possible, include a URL.
    - The name of the license (e.g., CC-BY 4.0) should be included for each asset.
    - For scraped data from a particular source (e.g., website), the copyright and terms of service of that source should be provided.
    - If assets are released, the license, copyright information, and terms of use in the package should be provided. For popular datasets, `paperswithcode.com/datasets` has curated licenses for some datasets. Their licensing guide can help determine the license of a dataset.
    - For existing datasets that are re-packaged, both the original license and the license of the derived asset (if it has changed) should be provided.
    - If this information is not available online, the authors are encouraged to reach out to the asset's creators.

13. **New assets**

    Question: Are new assets introduced in the paper well documented and is the documentation provided alongside the assets?

    Answer: **[TODO]**

    Justification: **[TODO]**

    Guidelines:

    - The answer [N/A] means that the paper does not release new assets.
    - Researchers should communicate the details of the dataset/code/model as part of their submissions via structured templates. This includes details about training, license, limitations, etc.
    - The paper should discuss whether and how consent was obtained from people whose asset is used.
    - At submission time, remember to anonymize your assets (if applicable). You can either create an anonymized URL or include an anonymized zip file.

14. **Crowdsourcing and research with human subjects**

    Question: For crowdsourcing experiments and research with human subjects, does the paper include the full text of instructions given to participants and screenshots, if applicable, as well as details about compensation (if any)?

    Answer: **[TODO]**

    Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper does not involve crowdsourcing nor research with human subjects.
- Including this information in the supplemental material is fine, but if the main contribution of the paper involves human subjects, then as much detail as possible should be included in the main paper.
- According to the NeurIPS Code of Ethics, workers involved in data collection, curation, or other labor should be paid at least the minimum wage in the country of the data collector.

15. **Institutional review board (IRB) approvals or equivalent for research with human subjects**

Question: Does the paper describe potential risks incurred by study participants, whether such risks were disclosed to the subjects, and whether Institutional Review Board (IRB) approvals (or an equivalent approval/review based on the requirements of your country or institution) were obtained?

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the paper does not involve crowdsourcing nor research with human subjects.
- Depending on the country in which research is conducted, IRB approval (or equivalent) may be required for any human subjects research. If you obtained IRB approval, you should clearly state this in the paper.
- We recognize that the procedures for this may vary significantly between institutions and locations, and we expect authors to adhere to the NeurIPS Code of Ethics and the guidelines for their institution.
- For initial submissions, do not include any information that would break anonymity (if applicable), such as the institution conducting the review.

16. **Declaration of LLM usage**

Question: Does the paper describe the usage of LLMs if it is an important, original, or non-standard component of the core methods in this research? Note that if the LLM is used only for writing, editing, or formatting purposes and does *not* impact the core methodology, scientific rigor, or originality of the research, declaration is not required.

Answer: **[TODO]**

Justification: **[TODO]**

Guidelines:

- The answer [N/A] means that the core method development in this research does not involve LLMs as any important, original, or non-standard components.
- Please refer to our LLM policy in the NeurIPS handbook for what should or should not be described.