


8.8 Compute the difference between the pure-component and partial molar enthalpies for both components at 298.15 K and various compositions in each of the following mixtures using the data in Fig. 8.1-2b.

- a. benzene- $C_6F_5H$  } by hand  
 b. benzene- $C_6F_6$  }  
 c. benzene- $C_6F_5Cl$  } by code w/ data from paper  
 d. benzene- $C_6F_5Br$  }  
 e. benzene- $C_6F_5I$  }

$$1:2 \rightarrow 0.66$$

$$2:1 \rightarrow 0.33$$

(a) bringing over the graph (similar to 8.6 figures)




now, @  $x_{C_6F_5Y} = 0.33$

@  $x_{C_6F_5Y} = 0.66$

Component	$\bar{H} - H$
$C_6H_6$	170 J/mol
$C_6F_5H$	-10 J/mol

Component	$\bar{H} - H$
$C_6H_6$	110 J/mol
$C_6F_5H$	-20 J/mol

(b)

@  $x = 0.33$ 

Component	$\bar{H} - H$
$\text{C}_6\text{H}_6$	100 J/mol
$\text{C}_6\text{F}_6$	-1050 J/mol

@  $x = 0.66$ 

Component	$\bar{H} - H$
$\text{C}_6\text{H}_6$	-1090 J/mol
$\text{C}_6\text{F}_6$	-280 J/mol

(c) now using the data / eqn. fits from the paper  
Sandler I'll do the others with code

cheg325 hw1 q2


AUTHOR  
kyle wodehouse

PUBLISHED  
February 14, 2025

grabbing the fit parameters from the paper sandler references and quickly plotting the curve


also note that the equations from the paper are given in the form

$$\Delta_{\text{mix}}H = x(1 - x) \left( h_0 + h_1(1 - 2x) + h_2(1 - 2x)^2 \right)$$



the derivative of the equations the paper provides are in this form


$$\frac{d}{dx_2} \Delta H_{\text{mix}} = (1 - 2x_2)(h_0 + h_1(1 - 2x_2) + h_2(1 - 2x_2)^2) + x_2(1 - x_2)(-2h_1 - 4h_2(1 - 2x_2))$$



Now we can do all the calculations needed for this problem by evaluating the derivatives at the desired points and use this equation

$$\bar{H}_{C_6H_6} - \underline{H}_{C_6H_6} = \Delta_{\text{mix}}\underline{H} - x_{C_6F_6Y} \frac{\partial \Delta_{\text{mix}}\underline{H}}{\partial x_{C_6F_6Y}}$$

and then this equation for  $C_6F_5Y$



benzene differences !!

df.pivot(index='composition', columns='Y')['benzene difference']

Y	Br	Cl	F	H	I
composition					
0.333	-199.082911	-17.835391	184.271225	177.887603	-473.017385
0.667	-1424.003857	-1200.267830	-836.973728	127.043932	-1994.638285

C6F5Y differences !!

df.pivot(index='composition', columns='Y')['C6F5Y difference']

Y	Br	Cl	F	H	I
composition					
0.333	-1529.168499	-1474.700279	-1275.061265	-43.700907	-1947.077871
0.667	-330.776461	-361.496370	-364.328594	-35.929454	-413.459347

# this is filler text. ignore please.