

HW1

Minh Luc, Devin Pham, Kyle Moore

Friday of Week 1, 04/01/2022

This is a template `Rmarkdown` file. Please find your partners and form a group for the assignments and final project. It is better to not change after your group is fixed.

Please read `Math189_HW_template.Rmd` carefully when you write your reports, and use the template file `Math189_HW_template.Rmd` to generate your reports. If you generate your report as a html file, please save it as a pdf file. Please submit your report to **Gradescope** before the deadline.

Question 0

Please write down you name and ID like below:

Member 1:

- Name: Minh Luc
- Student ID: A17209607

Member 2:

- Name: Kyle Moore
- Student ID: A14271413

Member 3:

- Name: Devin Pham
 - Student ID: A17198936
-

Question 1.

*Let $Z \sim N(0, 1)$.

(a) $P(Z > 1)$;

Answer: The probability can be calculated by

```
p <- pnorm(q = 1, mean = 0, sd = 1, lower = FALSE)
print(p)
```

```
## [1] 0.1586553
```

```
p3 <- round(p, digits = 3) # x rounded to 3 decimal places
print(p3)
```

```
## [1] 0.159
```

Therefore,

$$P(Z > 1) \approx 0.159.$$

(b) $P(Z < -1.06)$;

Answer: The probability can be calculated by

```
p <- pnorm(q = -1.06, mean = 0, sd = 1, lower = TRUE)
print(p)
```

```
## [1] 0.1445723
```

```
p3 <- round(p, digits = 3) # x rounded to 3 decimal places
print(p3)
```

```
## [1] 0.145
```

Therefore,

$$P(Z < -1.06) \approx 0.145.$$

(c) $P(-2.33 < Z \leq 1.06)$.

Answer: The probability can be calculated by

```
p <- pnorm(q = 1.06, mean = 0, sd = 1, lower = TRUE) - pnorm(q = -2.33, mean = 0, sd = 1, lower = TRUE)
print(p)
```

```
## [1] 0.8455246
```

```
p3 <- round(p, digits = 3) # x rounded to 3 decimal places
print(p3)
```

```
## [1] 0.846
```

Therefore,

$$P(-2.33 < Z \leq 1.06) \approx 0.846.$$

Question 2.

Let (X, Y) be a random vector whose pdf is given by

$$f_{X,Y}(x, y) = \frac{2}{3}(x + 2y), \quad 0 \leq x \leq 1, 0 \leq y \leq 1.$$

(a) Find the marginal distribution of X .

Answer: The marginal distribution of X is given by

$$\begin{aligned} f_X(x) &= \int_0^1 f_{X,Y}(x, y) dy \\ &= \int_0^1 \left(\frac{2}{3}(x + 2y) \right) dy \\ &= \frac{2}{3}x \int_0^1 dy + \frac{2}{3} \int_0^1 2y dy \\ &= \frac{2}{3}x \times 1 + \frac{2}{3} \times y^2 \Big|_0^1 \\ &= \frac{2}{3}x + \frac{2}{3}, \quad 0 \leq x \leq 1. \end{aligned}$$

Like this, if you want to split a long equation into several lines, you can use `\begin{aligned}` ... `\end{aligned}` environment or `\begin{split}` ... `\end{split}` environment inside the `$$... $$` environment. Please note that blank lines are not allowed in the `$$... $$` environment, otherwise the compile will not be successful. The double backslash `\\` at the end of each line works as a newline character. Use the ampersand character `&`, to set the points where the equations are vertically aligned.

(b) Find the marginal distribution of Y .

Answer: The marginal distribution of y is given by

$$\begin{aligned}
 f_Y(y) &= \int_0^1 f_{X,Y}(x,y) dx \\
 &= \int_0^1 \left(\frac{2}{3}(x+2y) \right) dx \\
 &= \frac{2}{3} \int_0^1 x dx + \frac{4}{3} y \int_0^1 dx \\
 &= \frac{2}{3} \times \frac{x^2}{2} \Big|_0^1 + \frac{4}{3} y \times 1 \\
 &= \frac{2}{3} \times \frac{1}{3} + \frac{4}{3} y \\
 &= \frac{1}{3} + \frac{4}{3} y, \quad 0 \leq y \leq 1.
 \end{aligned}$$

(c) Find $E(X)$, $E(Y)$, $\text{Var}(X)$ and $\text{Var}(Y)$.

$E(X)$:

Since $E(X) = \int x f_X(x) dx$, using the above marginal distribution:

```
fx = function(x){
  x*((2/3)*x + (2/3)) # x times marginal fx
}
```

```
mean_x = integrate(fx, lower=0, upper=1)
print(round(mean_x$value, digits=3))
```

```
## [1] 0.556
```

$$E(X) \approx .556$$

$\text{Var}(X)$:

Since $\text{Var}(X) = E(X^2) - E(X)^2$, using the above marginal distribution and $E(X)$:

```
fx = function(x){
  x*((2/3)*x + (2/3)) # x times marginal fx
}
```

```
fx2 = function(x){
  (x^2)*((2/3)*x + (2/3)) # x^2 times marginal fx
}
```

```
var_x = (integrate(fx2, lower=0, upper=1)$value -
  (integrate(fx, lower=0, upper=1)$value)^2)
print(round(var_x, digits=3))
```

```
## [1] 0.08
```

$$\text{Var}(X) \approx .080$$

$E(Y)$:

Since $E(Y) = \int y f_Y(y) dy$, using the above marginal distribution:

```
fy = function(y){  
  y*((1/3) +(4/3)*y) # y times marginal fy  
}
```

```
mean_y = integrate(fy, lower=0, upper=1)  
print(round(mean_y$value, digits=3))
```

```
## [1] 0.611
```

$$E(Y) \approx .611$$

$\text{Var}(Y)$:

Since $\text{Var}(Y) = E(Y^2) - E(Y)^2$, using the above marginal distribution and $E(Y)$:

```
fy = function(y){  
  y*((1/3) +(4/3)*y) # y times marginal fy  
}
```

```
fy2 = function(y){  
  (y^2)*(((1/3) +(4/3)*y)) # y^2 times marginal fy  
}
```

```
var_y = (integrate(fy2, lower=0, upper=1)$value -  
         (integrate(fy, lower=0, upper=1)$value)^2)  
print(round(var_y, digits=3))
```

```
## [1] 0.071
```

$$\text{Var}(Y) \approx .071$$

(d) Find $\text{Cov}(X, Y)$.

We will use the equation $\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$

$$\begin{aligned}
E(XY) &= \int_0^1 \int_0^1 xy f_{X,Y}(x,y) dx dy \\
&= \int_0^1 \int_0^1 xy \left(\frac{2}{3}x + \frac{4}{3}y \right) dx dy \\
&= \frac{1}{3} \int_0^1 \int_0^1 (2x^2y + 4xy^2) dx dy \\
&= \frac{1}{3} \int_0^1 2y \frac{x^3}{3} \Big|_0^1 + 4y^2 \frac{x^2}{2} \Big|_0^1 dy \\
&= \frac{1}{3} \int_0^1 2y^2 + \frac{2}{3}y dy \\
&= \frac{1}{3} \left(\frac{2}{3} \frac{y^3}{3} \Big|_0^1 + \frac{2}{3} \frac{y^3}{3} \Big|_0^1 \right) \\
&= \frac{1}{3} \left(\frac{1}{3} + \frac{2}{3} \right) \\
&= \frac{1}{3} \approx .333
\end{aligned}$$

Then plug into the equation:

$$\begin{aligned}
\text{Cov}(X,Y) &= E(XY) - E(X)E(Y) \\
&= .333 - (.556)(.611) \\
&= .333 - .339716 \\
&= .006716 \approx .007
\end{aligned}$$

(e) Find $\text{Cor}(X,Y)$.

$$\begin{aligned}
\text{Cor}(X,Y) &= \frac{\text{Cov}(X,Y)}{\sigma_X \sigma_Y} \\
&= \frac{.007}{\sqrt{\text{Var}(X)} \sqrt{\text{Var}(Y)}} \\
&= \frac{.007}{\sqrt{.080} \sqrt{.071}} \\
&= .0928804 \approx .093
\end{aligned}$$

Question 3.

(a) Let $n \geq 1$ and let X_1, \dots, X_n be a sample from $N(\mu, \sigma^2)$, where μ and σ^2 are both unknown. Provide an estimator $\hat{\mu}$ of the unknown parameter μ ?

Answer: An estimator of the population mean is given by

$$\hat{\mu} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i = \frac{1}{n} (X_1 + \dots + X_n).$$

(b) Following (a), provide an estimator $\hat{\sigma}^2$ of the unknown parameter σ^2 .

- (c) Read the [R documentation webpage](#) again, and use `rnorm` to generate $n = 10$ normal random numbers with $mean = 1$ and $sd = 3$.

```
rnorm(n = 10, mean = 1, sd = 3)
```

```
## [1] -1.1715226 1.4743347 1.6945760 -3.6123964 -0.1741923 3.5629765
## [7] -3.9003427 -3.3987796 1.4788274 3.0751290
```

However, when we re-run the line above, the result is different:

```
rnorm(n = 10, mean = 1, sd = 3)
```

```
## [1] -0.0876810 4.0887891 -3.1992997 5.3632474 -3.4938273 4.9067789
## [7] -0.7225247 5.9745749 1.4140158 5.7731955
```

To keep the results unchanged, we can set a random seed every time before we use the `rnorm` function:

```
set.seed(1)
```

```
rnorm(n = 10, mean = 1, sd = 3)
```

```
## [1] -0.87936143 1.55092997 -1.50688584 5.78584241 1.98852332 -1.46140515
## [7] 2.46228716 3.21497412 2.72734405 0.08383484
```

```
set.seed(1)
```

```
rnorm(n = 10, mean = 1, sd = 3)
```

```
## [1] -0.87936143 1.55092997 -1.50688584 5.78584241 1.98852332 -1.46140515
## [7] 2.46228716 3.21497412 2.72734405 0.08383484
```

Now, choose an arbitrary integer as your own seed, and generate $n = 5$ normal random numbers with $mean = 2$ and $sd = 1$. Based on (a), find your value of $\hat{\mu}$ using your random numbers and find the difference between $\hat{\mu}$ and the true mean μ .

- (d) Discuss how can you estimate μ more accurately, and explain your idea.