

Lecture Notes

Kyle Chui

2022-01-04

Contents

1	Lecture 1	1
1.1	Overview of This Course	1
1.2	The Beginnings	1
1.3	Egyptian Mathematics	1
1.3.1	Egyptian Number System	1
1.3.2	Egyptian Arithmetic	1
2	Lecture 2	3
2.1	Babylonian Mathematics	3
2.1.1	Babylonian Number System	3
2.1.2	Babylonian Algebra	3
2.1.3	Babylonian Geometry	3
3	Lecture 3	5
3.1	Greek Mathematics	5
3.1.1	The Theorem of Pythagoras	5
3.1.2	Rational points on the circle	6

1 Lecture 1

1.1 Overview of This Course

“Ancient” history of mathematics

- Egyptian mathematics (c. 3200 BCE – 300 BCE): Number system in base 10, fractions.
- Babylonian mathematics (c. 3300 BCE – 500 BCE): Place-value sexagesimal number system, roots of algebra.
- Greek mathematics (c. 700 BCE – 400 CE): Deduction method, geometry, roots of number theory.
- Mathematics in Asia (c. 1100 BCE – 1200 CE): Roots of number theory and algebra.
- Polynomial equations.
- Development of calculus, infinite series (16th–17th centuries CE).

1.2 The Beginnings

What is “Mathematics”?

1. Logical-deductive mathematics (initiated by the Greeks). This is not the generally accepted view of “mathematics” anymore.
2. Abstract counting problems: “mathematical” exercises without a direct practical use.
3. Practical counting problems: Bookkeeping for bureaucracy, inventories of goods and harvest, calculations involving lengths and areas for farming purposes, etc.
4. Development of a number system.

Note. We will see later that 2–4 all occur almost simultaneously.

1.3 Egyptian Mathematics

Most writing was done on papyrus, which doesn’t preserve well, so we don’t have much evidence from this period.

1.3.1 Egyptian Number System

The Egyptians had a base 10 number system, with a new Hieroglyphic symbol for each power of 10. This number system is *not* place valued. One simply adds all the symbols to get the number. For instance $\cap\cap\cap||| = 34$. By comparison, our modern number system is place-valued, which means that the position of each digit matters, i.e. $254 \neq 452$.

1.3.2 Egyptian Arithmetic

Summation is extremely easy, you just write the numbers next to each other to get their sum. Multiplication was done via consecutive doubling and adding. We do this by decomposing one number into powers of 2, and then distributing the other number across this sum. Because of this method, Egyptians had tablets with powers of 2, and tablets with consecutive doublings of many numbers.

We also have sources of:

- Fractions—The Egyptians used almost exclusively fractions of the form $\frac{1}{n}$ (aside from $\frac{2}{3}$ and $\frac{3}{4}$) and used sums of these to express more general fractions.

- Notation—Since any fraction can be written as the sum of fractions of the form $\frac{1}{n}$, any general fraction could be written by just writing $\frac{1}{n}$ fractions next to each other.
- Algebra—We have evidence that they solved some linear and second order (quadratic) equations.
- Geometry—Though limited, we have evidence that the Egyptians could calculate areas and volumes of triangles, circles, pyramids, etc.

However we see no evidence for a general theory for solving these questions, and all of the problems that we know of are elementary and mostly practical. The earliest “advanced” mathematical resources from the Egyptians are from around 1900 BCE, and are predated by Babylonian mathematics.

2 Lecture 2

2.1 Babylonian Mathematics

The Babylonians, in comparison to the Egyptians:

- had a more advanced number system,
- solved more difficult problems,
- solved more abstract (“useless”) problems.

Hence people often view the Babylonians in Mesopotamia as being the starting point of “serious” mathematics.

Mesopotamia was located between the Tigris and Euphrates rivers, and its name means “the land between the rivers”. Historical texts from Mesopotamia are unusually well-preserved, because texts were made by making impressions on clay tablets, which hardened and preserve well.

2.1.1 Babylonian Number System

The Babylonian number system was a *sexagesimal* (base 60), place-valued system.

Note.

- The number zero didn’t exist just yet! So instead of writing “20”, they would just write “2”, and figure out what number it took from context.
- They also had fractions, but didn’t have a symbol for a decimal point, so they again guessed based on context.
- Furthermore, since there is no zero, there was no way of distinguishing between $1 \cdot 60^2 + 30 \cdot 1$, or $1 \cdot 60 + 30 \cdot \frac{1}{60}$, etc.

Later on in Mesopotamia, we see the first usage of the number zero (300 BCE), but only as a place holder to solve the aforementioned issues. This was also done by the Mayans around 400 CE. It was not until 600 CE that the number zero was invented in India, and carried its full weight as a “number”.

Note. There are still remnants of the Babylonian sexagesimal number system in the modern day.

- Time: Hours and minutes are divided into 60 pieces
- Angle measurements are still done in multiples/factors of 60, i.e. 360° .

2.1.2 Babylonian Algebra

This notation was also used to solve basic algebra questions, but we don’t know how exactly these problems were solved (probably the same method as the Egyptians, guess and check). It is believed that math problems were given to students in order to train them in literacy, numeracy, and applications to administration. The Babylonians also knew about the quadratic formula, although it was described in words as opposed to being given as an equation. Furthermore, although they knew of the quadratic formula, they did not have negative numbers, and so found only one solution to quadratic equations. This is the first instance we know of where a civilization has an “algorithmic” solution to a problem.

2.1.3 Babylonian Geometry

We have some sources of Babylonian geometry that indicate that they knew of the Pythagorean Theorem. One example of this is the “IM 67118” tablet, which gives the sides of a rectangle given its area and diagonal

length. There is a tablet called “Plimpton 322” that contains Pythagorean triples, or integer triplets a, b, c satisfying $a^2 + b^2 = c^2$.

3 Lecture 3

Another tablet titled “YBC7289” contains approximations for $\sqrt{2}$ and $\frac{1}{\sqrt{2}}$ that has an error of less than 1 in 2 million.

3.1 Greek Mathematics

Note. A lot of “Greek” mathematicians did not come from what is nowadays Greece, but rather the old Hellenistic empire (which was larger).

3.1.1 The Theorem of Pythagoras

Probably the oldest “mathematical theorem”, known (in some form) to:

- Egyptians
- Babylonians
- Chinese
- Indians
- Greeks

before Pythagoras.

What we know about Pythagoras with (quasi-)certainty

- He existed (~ 580 BCE–500 BCE)
- He was born in Samos (modern day Turkey)
- He left no writings himself
- He founded a school in Croton (c. 540 BCE)
 - We say “school” but it was more like a sect, with strict rules and secrecy
 - Their central belief was “All is number”, and that every number has a meaning
 - The number 10 was a “holy number”, often drawn as a triangle
 - Theory of music based on whole fractions
 - Largely responsible for introducing “rigorous” mathematics
 - They viewed “pure” mathematics as the opposite of “applied” mathematics

Theorem — *Pythagorean Theorem*

Given a right triangle with legs a and b and hypotenuse c , we have

$$a^2 + b^2 = c^2.$$

Definition. *Pythagorean Triples*

Triples (a, b, c) that satisfy $a^2 + b^2 = c^2$ are called *Pythagorean triples*.

General Formula: For positive integers p, q, r such that $p \geq q$, we can use the following equations:

$$a = (p^2 - q^2)r$$

$$b = 2pqr$$

$$c = (p^2 + q^2)r$$

“Facts”:

- Less general formulas were known to Pythagoras’ school, Plato, Babylonians...
- The first statement *and* proof of this formula is from Euclid’s “Elements”, book 10

3.1.2 Rational points on the circle

Finding rational points on the unit circle with rational coordinates is actually very related to finding Pythagorean triples.