

Lecture Notes

Kyle Chui

2022-01-04

Contents

1	Lecture 1	1
1.1	Introduction to Dynamical Systems	1
1.1.1	Where Do “Dynamical Systems” Come From?	1
1.2	Autonomous ODEs	1
2	Lecture 2	2
2.1	Reducing ODEs to First Order Autonomous Systems	2
2.1.1	Flows on the Line	3
2.1.2	Fixed Points	3
2.1.3	Stability	4
3	Lecture 3	5
3.1	Potentials	6
4	Lecture 4	7
4.1	Impossibility of Oscillations	7
4.2	Numerical Methods	8
4.2.1	Integral Equations	8
4.2.2	Numerical Approximation	8

1 Lecture 1

1.1 Introduction to Dynamical Systems

Models of real-world phenomena can often be classified as either *static* or *dynamic*. Furthermore, these systems can either be *discrete* (x_1, x_2, \dots where $x_i \in \mathbb{R}$ for $i \geq 1$) or *continuous* ($x = x(t)$ where $t \geq 0$ and $x \in \mathbb{R}$, and $\dot{x} = f(x)$).

1.1.1 Where Do “Dynamical Systems” Come From?

1. Observed phenomena
2. Mathematical model
3. “Solve” the model
4. Make predictions

1.2 Autonomous ODEs

Definition. *Autonomous ODEs*

We say that an ordinary differential equation is *autonomous* if the right-hand side does not depend on t .

- The SIR (susceptible, infected, recovered) model is an example of a *first order* system of *autonomous* ODEs.

$$\begin{cases} \dot{x}_1 = f_1(x_1, \dots, x_n) \\ \vdots \\ \dot{x}_n = f_n(x_1, \dots, x_n) \end{cases}$$

- We will refer to n as the *dimension* of the system.

2 Lecture 2

2.1 Reducing ODEs to First Order Autonomous Systems

Consider the set of differential equations given by

$$\begin{cases} \dot{x} = -\kappa(t)xy, \\ \dot{y} = \kappa(t)xy - \delta y, \\ \dot{z} = \delta y. \end{cases}$$

Introduce a new variable, i.e. $\tau = \tau(t) = t$. Then we may rewrite the above as

$$\begin{cases} \dot{x} = -\kappa(\tau)xy, \\ \dot{y} = \kappa(\tau)xy - \delta y, \\ \dot{z} = \delta y, \\ \dot{\tau} = 1. \end{cases}$$

Note that the above system is now autonomous.

Example. The Pendulum

We can model the angle θ of a pendulum of length $L > 0$ by

$$\ddot{\theta} + \frac{g}{L} \sin \theta = 0.$$

Applying Newton's Second Law, we can get the equations

$$\begin{aligned} mL\ddot{\theta} &= -mg \sin \theta \\ \theta &= \theta(t). \end{aligned}$$

Observe that if we let $x = \theta$ and $y = \dot{\theta}$, then we get

$$\begin{cases} \dot{x} = y \\ \dot{y} = -\frac{g}{L} \sin x. \end{cases}$$

Example. Pendulum with an external force

If we add an external force to our pendulum, then we get

$$\ddot{\theta} + \frac{g}{L} \sin \theta = \frac{1}{m} F(t).$$

Thus if we let $x = \theta$, $y = \dot{\theta}$, and $z = t$, then we get

$$\begin{cases} \dot{x} = y \\ \dot{y} = -\frac{g}{L} \sin x + \frac{1}{m} F(z) \\ \dot{z} = 1. \end{cases}$$

Note. In general, higher order ODEs of the form

$$\frac{d^k x}{dt^k} = f\left(x, \frac{dx}{dt}, \frac{d^2 x}{dt^2}, \dots, \frac{d^{k-1} x}{dt^{k-1}}\right)$$

can be converted into a first order system by taking

$$z_1 = x, z_2 = \frac{dx}{dt}, \dots, z_k = \frac{d^{k-1}x}{dt^{k-1}}.$$

We get the system

$$\begin{cases} \dot{z}_1 = \frac{dx}{dt} = z_2 \\ \dot{z}_2 = \frac{d^2x}{dt^2} = z_3 \\ \vdots \\ \dot{z}_k = f(z_1, z_2, \dots, z_k) \end{cases}$$

2.1.1 Flows on the Line

We will now consider systems of the form

$$\dot{x} = f(x)$$

where $f: \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function.

Example. Consider the ODE given by

$$\dot{x} = x(x+1)(x-1)^2.$$

We could use separation of variables to solve this.

Note. Solutions to ODEs usually come in three different flavors:

- Analytic methods (separation of variables)
- Geometric methods (direction fields)
- Numerical methods (Euler's method)

Definition. *Phase Space*

To help us analyze these differential equations, we can plot \dot{x} against x on a graph, and see the behavior around zeroes. This is called a *phase space*. If some neighborhood of points around a zero x tend towards x , then x is called a *stable point*. If they tend to move away from x , then x is an *unstable point*. On a phase space graph, we denote stable points with \bullet , unstable points with \circ , and other points with a half-filled circle.

2.1.2 Fixed Points

Definition. *Fixed Point*

We say that x^* is a *fixed point* of the system

$$\dot{x} = f(x)$$

if $f(x^*) = 0$. If x^* is a fixed point then the system has a constant solution given by $x(t) = x^*$. These points are also known as equilibrium points, stationary points, rest points, critical points, and steady states.

2.1.3 Stability

Definition. *Stability*

Let x^* be a fixed point of the system

$$\dot{x} = f(x).$$

For now, we say that x^* is:

- *Stable* if solutions starting close to x^* approach x^* as $t \rightarrow \infty$.
- *Unstable* if solutions starting close to x^* diverge from x^* as $t \rightarrow \infty$.
- *Half-stable* if solutions starting close to x^* approach x^* from one side, but diverge from the other side.

3 Lecture 3

Question. Can we say more about what happens close to fixed points?

Example. Consider the equation given by

$$\dot{x} = x(x+1)(x-1)^2,$$

which has stable points at -1 , 0 , and 1 .

We will use something called the *local method*. We define a new function $\eta(t) = x(t) + 1$, so $x(t) = \eta(t) - 1$. Hence $\dot{x}(t) = \dot{\eta}(t)$. Furthermore,

$$\begin{aligned} x(x+1)(x-1)^2 &= (\eta-1)\eta(\eta-2)^2 \\ &= -4\eta + O(\eta^2), \end{aligned} \quad (\eta \rightarrow 0)$$

so $\dot{\eta} \approx -4\eta$. Near $x = -1$, $\eta = x + 1$ and it satisfies $\dot{\eta} = -4\eta$, so $\eta(t) \approx Ce^{-4t}$. We can see that this approaches 0 as $t \rightarrow \infty$, so points around $x = -1$ will approach -1 .

In general, we have the following method:

Assume that x^* is a fixed point of $\dot{x} = f(x)$, i.e. $f(x^*) = 0$. Let $\eta = x - x^*$. Then

$$\begin{aligned} \dot{\eta} &= \dot{x} \\ &= f(x) \\ &= f(x^*) + f'(x^*)\eta + \frac{1}{2}f''(x^*)\eta^2 + \dots \\ &= f'(x^*)\eta + O(\eta^2). \end{aligned} \quad (\eta \rightarrow 0)$$

Hence the equation

$$\dot{\eta} = f'(x^*)\eta$$

is the *linearization* at $x = x^*$. We know that the solution to such a differential equation is

$$\eta(t) = Ce^{f'(x^*)t} = \begin{cases} 0, & f'(x^*) < 0 \\ \pm\infty, & f'(x^*) > 0 \end{cases},$$

as $t \rightarrow \infty$. In the first case, the terms near x^* will tend towards x^* , and in the latter they will diverge from x^* .

Theorem. Suppose that x^* is a fixed point of the system $\dot{x} = f(x)$. Then if

- $f'(x^*) < 0$, the fixed point x^* is stable.
- $f'(x^*) > 0$, the fixed point x^* is unstable.

Question. What happens if $f'(x^*) = 0$? Anything can happen/the test is inconclusive.

- Consider the equation $\dot{x} = x^3$. We see that $x^* = 0$ is a critical point, and that $f'(x) = 3x^2$, so $f'(x^*) = 0$. Using our usual graphical methods, we can see that points to the left of x^* will approach x^* , and so will points to the right, and so $x^* = 0$ is a stable point.
- If we use the equation $\dot{x} = -x^3$, we get the direct opposite, that is $x^* = 0$ is unstable despite having the same critical point.
- If we look at the behavior of $\dot{x} = x^2$, then we get that the critical point at $x^* = 0$ is half-stable.
- If we consider the equation $\dot{x} = 0$, then every number on the real line is a critical point, and so the solutions don't move at all.

3.1 Potentials

- Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be smooth and consider the system

$$\dot{x} = f(x).$$

- A function $V: \mathbb{R} \rightarrow \mathbb{R}$ so that

$$f(x) = -V'(x)$$

is called a *potential* for f .

- Our system can be written as a *gradient flow*

$$\dot{x} = -V'(x).$$

Note. Potential functions are *not* unique, since you can always add a constant.

4 Lecture 4

Example. Consider the differential equation $\dot{x} = x - x^3$. Since we have $\dot{x} = -V'(x)$, we have

$$V(x) = \frac{1}{4}x^4 - \frac{1}{2}x^2 = \frac{1}{4}(x^2 - 1)^2 - \frac{1}{4}.$$

When we look at the graph of $V'(x)$, we may pretend that there is a ball rolling down a hill from every point, which tells us how to find the stability of points. Each point will settle in the first “well” that it meets.

Theorem. Let $V: \mathbb{R} \rightarrow \mathbb{R}$ be smooth and consider the system

$$\dot{x} = -V'(x).$$

Then the *potential energy* $V(x(t))$ is non-increasing (as a function of time). Furthermore, if $x(t)$ is not a fixed point for all $t \in (T_1, T_2)$, then the potential energy is strictly decreasing on (T_1, T_2) .

Proof. Observe that

$$\begin{aligned} \frac{d}{dt}V(x(t)) &= V'(x(t)) \cdot \dot{x}(t) \\ &= -V'(x(t))^2. \end{aligned}$$

Hence the potential energy is non-increasing, as its derivative is always non-positive. Thus if $V'(x_1) = 0$, then x_1 is a critical point! \square

Corollary. Let $V: \mathbb{R} \rightarrow \mathbb{R}$ be smooth and consider the system

$$\dot{x} = -V'(x).$$

If x^* is an isolated critical point of V then

- If it is a local minima of V , it is a stable fixed point.
- If it is a local maxima of V , it is an unstable fixed point.
- If it is an inflection point of V , it is a half-stable fixed point.

Proof. If we imagine the ball analogy again, we can see that if x^* is a local minima then points around it will tend towards x^* , and so it is stable. The opposite happens for divergence near a local maxima, and the analogy still holds for the half-stableness when x^* is an inflection point. \square

Note. Every one dimensional system is a gradient flow, because if f is smooth, then we can take the integral and define

$$V(x) := - \int_0^x f(s) \, ds.$$

4.1 Impossibility of Oscillations

Definition. *Periodic Functions*

If there exists a constant $p > 0$ so that for all t we have

$$x(t + p) = x(t),$$

then we say that p is *periodic*.

Note. All constant functions are periodic.

Theorem. There are no non-constant periodic solutions of the system

$$\dot{x} = f(x).$$

Proof. Suppose that x is a periodic solution, with period $p > 0$. If $0 \leq t \leq p$ then, as the potential energy is non-increasing,

$$V[x(p)] \leq V[x(t)] \leq V[x(0)].$$

Since $x(p) = x(0)$, we have $V[x(t)]$ is constant. Hence $x(t)$ is constant. \square

4.2 Numerical Methods

4.2.1 Integral Equations

We want to find a solution of the equation

$$\begin{cases} \dot{x} = f(x) \\ x(0) = x_0 \end{cases}$$

Observe that

$$\begin{aligned} \dot{x} &= f(x) \\ \int_0^t \frac{dx(s)}{ds} ds &= \int_0^t f(x(s)) ds \\ x(t) - x(0) &= \int_0^t f(x(s)) ds \\ x(t) &= x_0 + \int_0^t f(x(s)) ds. \end{aligned}$$

We call this an *integral equation*, because the unknown now appears in the integral.

Example. Write the equation

$$\begin{cases} \dot{x} = \sin x \\ x(0) = 1 \end{cases}$$

as an integral equation.

We have that

$$x(t) = 1 + \int_0^t \sin(x(s)) ds.$$

4.2.2 Numerical Approximation

Suppose we have the equation

$$\begin{cases} \dot{x} = f(x) \\ x(0) = x_0 \end{cases}$$

Let's take $\Delta t > 0$ small. Then we have

$$x(\Delta t) = x_0 + \int_0^{\Delta t} f(x(s)) ds.$$

We will approximate $x(s) \approx x_0$ on the interval $(0, \Delta t)$. Thus we have

$$\begin{aligned} x(\Delta t) &= x_0 + \int_0^{\Delta t} f(x(s)) \, ds \\ &= x_0 + \int_0^{\Delta t} f(x_0) \, ds \\ &= x_0 + f(x_0)\Delta t. \end{aligned} \qquad (x_1 := x_0 + f(x_0)\Delta t)$$

Euler's method is to repeat the above to get $x_2 \approx x(2\Delta t)$. We have

$$\begin{aligned} x_2 &= x_1 + f(x_1)\Delta t, \\ x_3 &= x_2 + f(x_2)\Delta t, \\ &\dots \\ x_{n+1} &= x_n + f(x_n)\Delta t. \end{aligned}$$