

Missing Data Mechanisms

Utrecht University Winter School: Missing Data in R



**Utrecht
University**

Kyle M. Lang

Department of Methodology & Statistics
Utrecht University

Outline

Definitions

Consequences

Testing



What are Missing Data?

Missing data are empty cells in a dataset where there should be observed values.

- The missing cells correspond to true population values, but we haven't observed those values.



What are Missing Data?

Missing data are empty cells in a dataset where there should be observed values.

- The missing cells correspond to true population values, but we haven't observed those values.

Not every empty cell is a missing datum.

- Quality-of-life ratings for dead patients in a mortality study
- Firm profitability after the company goes out of business
- Self-reported severity of menstrual cramping for men
- Empty blocks of data following “gateway” items



A Little Notation

Y := An $N \times P$ Matrix of Arbitrary Data

Y_{mis} := The *missing* part of Y

Y_{obs} := The *observed* part of Y

R := An $N \times P$ response matrix

M := An $N \times P$ missingness matrix

The R and M matrices are complementary.

- $r_{np} = 1$ means y_{np} is observed; $m_{np} = 1$ means y_{np} is missing.
- $r_{np} = 0$ means y_{np} is missing; $m_{np} = 0$ means y_{np} is observed.
- M_p is the *missingness* of Y_p .

Missing Data Mechanisms

Missing Completely at Random (MCAR)

- $P(R|Y_{mis}, Y_{obs}) = P(R)$
- Missingness is unrelated to any study variables.

Missing at Random (MAR)

- $P(R|Y_{mis}, Y_{obs}) = P(R|Y_{obs})$
- Missingness is related to only the *observed* parts of study variables.

Missing not at Random (MNAR)

- $P(R|Y_{mis}, Y_{obs}) \neq P(R|Y_{obs})$
- Missingness is related to the *unobserved* parts of study variables.



Simulate Some Toy Data

```
library(mvtnorm); library(dplyr); library(magrittr)

set.seed(235711)

nObs <- 5000 # Sample Size
pm   <- 0.3  # Proportion Missing

sigma <- matrix(c(1.0, 0.5, 0.3,
                  0.5, 1.0, 0.0,
                  0.3, 0.0, 1.0),
               ncol = 3)
dat0 <- rmvnorm(nObs, c(0, 0, 0), sigma) %>% data.frame()
colnames(dat0) <- c("x", "y", "z")

dat0 %$% cor(y, x)

[1] 0.4997145
```

MCAR Example


```
## Simulate MCAR Missingness:
m <- sample(1:nObs, size = pm * nObs)

## Impose MCAR missing on Y:
mcarData <- dat0
mcarData[m, "y"] <- NA


## Check the correlation between X & Y:
mcarData %$% cor(y, x, use = "pairwise")

[1] 0.5195767
```


MCAR Example



MCAR Example



MAR Example


```
## Simulate MAR Missingness:
m <- with(dat0, x < quantile(x, probs = pm))

## Impose MAR missing on Y:
marData      <- dat0
marData[m, "y"] <- NA


## Check the correlation between X & Y:
marData %$% cor(y, x, use = "pairwise")

[1] 0.3822143
```

MAR Example



MAR Example



MNAR Example


```
## Simulate MNAR Missingness:
m <- with(dat0, y < quantile(y, probs = pm))

## Impose MNAR missing on Y:
mnarData      <- dat0
mnarData[m, "y"] <- NA


## Check the correlation between X & Y:
mnarData %$% cor(y, x, use = "pairwise")

[1] 0.3902962
```

MNAR Example



MNAR Example



Crucial Nuance

In our previous MAR example, ignoring the predictor of missingness actually produces *Indirect MNAR*.

Crucial Nuance

In our previous MAR example, ignoring the predictor of missingness actually produces *Indirect MNAR*.

QUESTION: What happens if we ignore the predictor of missingness, but that predictor is independent of our study variables?

Crucial Nuance

In our previous MAR example, ignoring the predictor of missingness actually produces *Indirect MNAR*.



QUESTION: What happens if we ignore the predictor of missingness, but that predictor is independent of our study variables?

```
m <- with(dat0, z < quantile(z, probs = pm))  
  
mcarData2          <- dat0  
mcarData2[m, "y"] <- NA  
  
mcarData2 %$% cor(y, x, use = "pairwise")  
  
[1] 0.5118075
```

ANSWER: We get back to MCAR :)



Crucial Nuance

The missing data mechanisms are not simply characteristics of an incomplete dataset; we also need to account for the analysis.



Crucial Nuance


The missing data mechanisms are not simply characteristics of an incomplete dataset; we also need to account for the analysis.



Testing the Missing Data Mechanism

We cannot fully test the MAR or MNAR assumptions.

- To do so would require knowing the values of the missing data.
- We can find observed predictors of missingness.
 - Use classification algorithms to predict missingness from Y_{obs} .
 - We can never know that we have discovered all MAR predictors.
- In practice, MAR and MNAR live on the ends of a continuum.
 - Our missing data problem exists at some unknown point along this continuum.
 - We can do a lot to nudge our problem towards the MAR side.



Testing the Missing Data Mechanism

We can (partially) test the MCAR assumption.

- With MCAR, the missing data and the observed data should have the same distribution.
- We can test for MCAR by testing the distributions of *auxiliary variables*, \mathbf{Z} .
 - Use a t-test to compare the subset of \mathbf{Z}_p that corresponds to \mathbf{Y}_{mis} to the subset corresponding to \mathbf{Y}_{obs} .
 - The Little (1988) MCAR test is a multivariate version of this.

These procedures actually test if the data are *observed* completely at random.

References

Little, R. J. A. (1988). A test of missing completely at random for multivariate data with missing values. *Journal of the American Statistical Association*, 83(404), 1198–1202. doi: 10.1080/01621459.1988.10478722

