

CFA Identification & Estimation

Theory Construction and Statistical Modeling



**Utrecht
University**

Kyle M. Lang

Department of Methodology & Statistics
Utrecht University

Outline

Model Identification

- Under-Identified Models

- Over-Identified Models

Model Estimation

Model Fit

Model Complexity

Model Comparison

- Nested Models

- Non-Nested Models

Example



MODEL IDENTIFICATION



Identifying Constraints

We must fix some parameters to identify the model.

- For each construct, we need $df \geq 0$.
- If the construct has three or more indicators:
 - Fix one parameter in the covariance model.
 - Fix one parameter in the mean model.
- If the construct has two indicators:
 - Fix an additional parameter in the covariance model.
- If the construct has only one indicator:
 - Cannot define a latent factor.



Under-Identified: Unconstrained Model


$$\Psi = [\psi_{11}]$$

$$\Lambda = \begin{bmatrix} \lambda_{11} \\ \lambda_{21} \end{bmatrix}$$

$$\Theta = \begin{bmatrix} \theta_{11} & \\ 0 & \theta_{22} \end{bmatrix}$$

$$Q = \frac{2(2+1)}{2} = 3$$

$$\begin{aligned} df &= Q - P \\ &= 3 - 5 = -2 \end{aligned}$$




Under-Identified: Marker-Variable

$$\Psi = [\psi_{11}]$$

$$\Lambda = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\Theta = \begin{bmatrix} \theta_{11} & \\ 0 & \theta_{22} \end{bmatrix}$$

$$\begin{aligned} df &= Q - P \\ &= 3 - 3 = 0 \end{aligned}$$




Under-Identified: Fixed-Factor

$$\Psi = [1]$$

$$\Lambda = \begin{bmatrix} \lambda \\ \lambda \end{bmatrix}$$

$$\Theta = \begin{bmatrix} \theta_{11} & \\ 0 & \theta_{22} \end{bmatrix}$$

$$\begin{aligned} df &= Q - P \\ &= 3 - 3 = 0 \end{aligned}$$




Over-Identified Models

What happens when we have four (or more) indicators?

With 4 indicators, our model contains 14 parameters:

- Four factor loadings
- Four residual variances
- Four item intercepts
- One latent mean
- One latent variance

$$Q = \frac{4(4+1)}{2} + 4 = 14$$
$$df = 14 - 14 = 0$$



Over-Identified Models

With 5 indicators, we have 17 model parameters:


- Five factor loadings
- Five residual variances
- Five item intercepts
- One latent mean
- One latent variance

$$Q = \frac{5(5+1)}{2} + 5 = 20$$
$$df = 20 - 17 = 3$$

With 6 indicators, we have 20 model parameters:

- Six factor loadings
- Six residual variances
- Six item intercepts
- One latent mean
- One latent variance

$$Q = \frac{6(6+1)}{2} + 6 = 27$$
$$df = 27 - 20 = 7$$



Over-Identified Models

With four indicators, we automatically have $df \geq 0$ without imposing any scaling constraints.

- Can we directly estimate the unconstrained model?

```
mod1 <- '  
fun =~ Q77 + Q84 + Q170 + Q196  
'
```

```
fit1 <- lavaan(mod1,  
               data      = sapi,  
               auto.var  = TRUE,  
               meanstructure = TRUE,  
               int.ov.free = TRUE,  
               int.lv.free = TRUE)
```

```
Warning: lavaan->lav_model_vcov():  
  Could not compute standard errors! The information matrix could not be  
  inverted. This may be a symptom that the model is not identified.
```

Hmmm...I guess not.

Over-Identified Models

```
partSummary(fit1, 1:4)
```

```
lavaan 0.6-19 ended normally after 13 iterations
```

Estimator	ML	
Optimization method	NLMINB	
Number of model parameters	14	
	Used	Total
Number of observations	970	1000

```
Model Test User Model:
```

Test statistic	58.017
Degrees of freedom	0

Over-Identified Models

```
partSummary(fit1, 7:8)
```

Latent Variables:

	Estimate	Std.Err	z-value	P(> z)
fun =~				
Q77	0.824	NA		
Q84	0.578	NA		
Q170	0.479	NA		
Q196	0.619	NA		

Intercepts:

	Estimate	Std.Err	z-value	P(> z)
.Q77	3.574	NA		
.Q84	3.232	NA		
.Q170	3.955	NA		
.Q196	3.803	NA		
fun	0.000	NA		

Over-Identified Models

```
partSummary(fit1, 9)
```

Variances:

	Estimate	Std.Err	z-value	P(> z)
.Q77	0.491	NA		
.Q84	0.760	NA		
.Q170	0.703	NA		
.Q196	0.370	NA		
fun	1.012	NA		

Over-Identified Models

In the above example, the data contain enough information to define our model parameters, but that's not enough.

- We still need scaling constraints.
- The estimation algorithm needs an anchor point from which to extrapolate the relative values of the model parameters.
- Without any scaling constraints, an infinite number of parameter matrices will produce the same $\hat{\mu}$ and $\hat{\Sigma}$.
- An infinite number of solutions are equally good.

```
## Fit a model to get some example parameters:  
mod1 <- '  
fun    =~ Q77 + Q84 + Q170 + Q196  
liked  =~ Q44 + Q63 + Q76  + Q98  
'  
  
fit1 <- cfa(mod1, data = sapi, effect.coding = TRUE)
```

Over-Identified Models

Extract the estimated factor loadings and latent covariance matrix:

```
(lambda <- inspect(fit1, "est")$lambda)
```

```
      fun liked
Q77  1.254 0.000
Q84  0.954 0.000
Q170 0.795 0.000
Q196 0.997 0.000
Q44   0.000 0.764
Q63   0.000 1.155
Q76   0.000 1.133
Q98   0.000 0.949
```

```
(psi <- inspect(fit1, "est")$psi)
```

```
      fun liked
fun    0.399
liked 0.241 0.311
```

Over-Identified Models

```
## Extract the estimated residual variances:
```

```
(theta <- inspect(fit1, "est")$theta)
```

	Q77	Q84	Q170	Q196	Q44	Q63	Q76	Q98
Q77	0.548							
Q84	0.000	0.727						
Q170	0.000	0.000	0.687					
Q196	0.000	0.000	0.000	0.364				
Q44	0.000	0.000	0.000	0.000	0.662			
Q63	0.000	0.000	0.000	0.000	0.000	0.807		
Q76	0.000	0.000	0.000	0.000	0.000	0.000	0.966	
Q98	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.469

Over-Identified Models

```
## Manually compute the model-implied covariance matrix:
```

```
(sigma <- lambda %*% psi %*% t(lambda) + theta)
```

	Q77	Q84	Q170	Q196	Q44	Q63	Q76	Q98
Q77	1.176							
Q84	0.477	1.090						
Q170	0.398	0.303	0.940					
Q196	0.499	0.380	0.316	0.761				
Q44	0.231	0.175	0.146	0.183	0.843			
Q63	0.349	0.265	0.221	0.277	0.275	1.223		
Q76	0.342	0.260	0.217	0.272	0.269	0.407	1.365	
Q98	0.287	0.218	0.182	0.228	0.226	0.341	0.335	0.750

Over-Identified Models

```
## Randomly sample an arbitrary scaling factor:
(a <- runif(1, 1, 2))

[1] 1.524314

## Rescale all factor loadings by a factor of a:
(lambda2 <- lambda * a)

      fun liked
Q77  1.911 0.000
Q84  1.454 0.000
Q170 1.212 0.000
Q196 1.520 0.000
Q44  0.000 1.164
Q63  0.000 1.760
Q76  0.000 1.727
Q98  0.000 1.447
```

Over-Identified Models

```
## Rescale the latent covariance matrix by a factor of (1 / a^2):  
(psi2 <- psi / a^2)  
  
      fun liked  
fun    0.172  
liked 0.104 0.134  
  
## Compute the model-implied covariance matrix using the rescaled parameters:  
(sigma2 <- lambda2 %*% psi2 %*% t(lambda2) + theta)  
  
      Q77    Q84   Q170   Q196    Q44    Q63    Q76    Q98  
Q77    1.176  
Q84    0.477 1.090  
Q170   0.398 0.303 0.940  
Q196   0.499 0.380 0.316 0.761  
Q44    0.231 0.175 0.146 0.183 0.843  
Q63    0.349 0.265 0.221 0.277 0.275 1.223  
Q76    0.342 0.260 0.217 0.272 0.269 0.407 1.365  
Q98    0.287 0.218 0.182 0.228 0.226 0.341 0.335 0.750
```

Over-Identified Models

```
## Compare the two model-implied covariance matrices:
```

```
all.equal(sigma, sigma2)
```

```
[1] TRUE
```

```
## Repeat the experiment 100 times:
```

```
out <- rep(NA, 100)
```

```
for(i in 1:100) {
```

```
  a <- runif(1, 1, 2)
```

```
  lambda2 <- lambda * a
```

```
  psi2 <- psi / a^2
```

```
  sigma2 <- lambda2 %*% psi2 %*% t(lambda2) + theta
```

```
  out[i] <- all.equal(sigma, sigma2)
```

```
}
```

```
## Always the same result?
```

```
all(out)
```

```
[1] TRUE
```

MODEL ESTIMATION



Model-Implied Statistics

Most statistical estimation algorithms operate by minimizing the difference between two key reference points:

1. The *model-implied* statistics/predictions/fitted values
 - The sufficient statistics implied by the structure of your model.
 - Predicted/fitted values produced by your model.
2. The *observed* statistics/values
 - The sufficient statistics calculated from the observed data.
 - The raw outcome values from your dataset.

The predictions/implied statistics produced by a good model must be simpler than the analogous quantities in the observed data.

- A model that exactly replicates the observed data is overfitting.
- The inferences from such models won't generalize to the population.

Model-Implied Statistics

You should already be familiar with this idea from OLS regression.

- The fitted values are the model implied statistics.

$$\hat{Y}_n = \hat{\beta}_0 + \sum_{p=1}^P \hat{\beta}_p X_{n,p}$$

- The raw outcome variable, Y , contains the observed values.
- Minimize the difference between $\hat{Y}$ and Y to estimate the model.

$$RSS = \sum_{n=1}^N (Y_n - \hat{Y}_n)^2$$




Hypothesized Model

We'll use this toy model to sketch out the intuition for maximum likelihood estimation of CFA models.

$$\alpha = [\alpha_1] \quad \Psi = [\psi_{11}]$$

$$\tau = \begin{bmatrix} \tau_1 \\ \tau_2 \\ \tau_3 \end{bmatrix} \quad \Lambda = \begin{bmatrix} \lambda_{11} \\ \lambda_{21} \\ \lambda_{31} \end{bmatrix}$$

$$\Theta = \begin{bmatrix} \theta_{11} & & \\ 0 & \theta_{22} & \\ 0 & 0 & \theta_{33} \end{bmatrix}$$



Parameters → Implied Statistics

The parameter matrices define the model-implied mean vector, $\hat{\mu}$, and covariance matrix, $\hat{\Sigma}$, via the following formulas.

$$\Sigma = \Lambda\Psi\Lambda^T + \Theta$$

$$\mu = \tau + \Lambda\alpha$$

By expanding these formulas, we can see how the model reproduces each mean, variance, and covariance.

$$\hat{\mu} = \begin{bmatrix} \tau_1 + \lambda_{11}\alpha_1 \\ \tau_2 + \lambda_{22}\alpha_1 \\ \tau_3 + \lambda_{33}\alpha_1 \end{bmatrix} \quad \hat{\Sigma} = \begin{bmatrix} \lambda_{11}\psi_{11}\lambda_{11} + \theta_{11} & & \\ \lambda_{11}\psi_{11}\lambda_{21} & \lambda_{21}\psi_{11}\lambda_{21} + \theta_{22} & \\ \lambda_{11}\psi_{11}\lambda_{31} & \lambda_{21}\psi_{11}\lambda_{31} & \lambda_{31}\psi_{11}\lambda_{31} + \theta_{33} \end{bmatrix}$$

Model-Implied Statistics for Estimation

The estimating algorithm chooses values for α , τ , Ψ , and Λ that minimize the differences between the model-implied statistics, $\{\hat{\mu}, \hat{\Sigma}\}$, and the sufficient statistics calculated from the observed data, $\{\bar{\mathbf{Y}}, \mathbf{S}\}$.

$$\hat{\mu} = \begin{bmatrix} \tau_1 + \lambda_{11}\alpha_1 \\ \tau_2 + \lambda_{22}\alpha_1 \\ \tau_3 + \lambda_{33}\alpha_1 \end{bmatrix} \quad \hat{\Sigma} = \begin{bmatrix} \lambda_{11}\psi_{11}\lambda_{11} + \theta_{11} & & \\ \lambda_{11}\psi_{11}\lambda_{21} & \lambda_{21}\psi_{11}\lambda_{21} + \theta_{22} & \\ \lambda_{11}\psi_{11}\lambda_{31} & \lambda_{21}\psi_{11}\lambda_{31} & \lambda_{31}\psi_{11}\lambda_{31} + \theta_{33} \end{bmatrix}$$

$$\bar{\mathbf{Y}} = \begin{bmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \bar{y}_3 \end{bmatrix} \quad \mathbf{S} = \begin{bmatrix} \text{var}(y_1) & & \\ \text{cov}(y_2, y_1) & \text{var}(y_2) & \\ \text{cov}(y_3, y_1) & \text{cov}(y_3, y_2) & \text{var}(y_3) \end{bmatrix}$$



Optimization: Intuition

We can formulate the familiar OLS objective as an abstract optimization problem as follows.

$$f(\beta_0, \beta_1, \dots, \beta_p) = \arg \min_{\beta_0, \beta_1, \dots, \beta_p} \|Y - \hat{Y}\|$$

Applying the same idea to our CFA optimization problem, we get the following formulation.

$$f(\Lambda, \Psi) = \arg \min_{\Lambda, \Psi} \|\text{Cov}(\mathbf{Y}) - \hat{\Sigma}\|$$

$$f(\tau, \alpha) = \arg \min_{\tau, \alpha} \|\bar{\mathbf{Y}} - \hat{\mu}\|$$



Optimization: Objective Function

In OLS regression, we minimize the residual sum of squares (RSS).

$$RSS = \sum_{n=1}^N \left(Y_n - \hat{Y}_n \right)^2$$

In CFA, we minimize the *model deviance*.

$$D_M = -2 \left[\ell \left(\hat{\Sigma}, \hat{\mu} \right) - \ell \left(\mathbf{S}, \bar{\mathbf{Y}} \right) \right]$$

Where, $\ell(\cdot)$ denotes a *loglikelihood* function.

- $\ell \left(\hat{\Sigma}, \hat{\mu} \right)$ returns the loglikelihood value for our estimated model.
- $\ell \left(\mathbf{S}, \bar{\mathbf{Y}} \right)$ returns the loglikelihood value for the *saturated model*.

Optimization: Loglikelihood Functions

A loglikelihood function is a model-based distance metric.

- Derived from the assumed data distribution (i.e., multivariate normal, for the basic CFA)
- Quantifies the "chances" of your estimated model having generated the observed data

$$\ell(\hat{\Sigma}, \hat{\mu}) = -\frac{K}{2} \ln(2\pi) - \frac{1}{2} \ln |\hat{\Sigma}| - \frac{1}{2} \sum_{n=1}^N (Y_n - \hat{\mu})^T \hat{\Sigma}^{-1} (Y_n - \hat{\mu})$$

Say we fit two model, M_a and M_b , to the same dataset. If M_a 's loglikelihood is closer to zero, we know that M_a represents the data better than M_b does.

- M_a 's implied statistics are closer to the observed sufficient statistics.
- M_a 's parameters are more plausible, given the data.

MODEL FIT



Model Fit: Intuition

We want a plausible, parsimonious, and useful model.

1. Our model should make theoretical sense.
2. Our model's implied statistics should closely align with the observed sufficient statistics.
3. Our model should be no more complex than necessary.

We can use *model fit statistics/indices* to evaluate the second and third criteria.




χ^2 Statistic

The χ^2 statistic is the most fundamental fit statistic.

- Derived directly from the objective function used to estimate the model.
- Allows rigorous hypothesis tests.
- Basis for most other fit indices.

$$\chi^2 = (N - 1)D_M$$

This statistic follows a χ^2 distribution with $df = Q - P$ (i.e., the same DF as the estimated model).



Problems with χ^2



RMSEA





SRMR



MODEL COMPLEXITY



DF & Complexity



Parsimony



MODEL COMPARISON



Comparing Nested Models



Likelihood Ratio Tests



Wald Tests



Score Tests



Comparing Non-Nested Models



Information Criteria



EXAMPLE




CFA of Extraversion Items

Suppose we hypothesize two distinct dimensions of extraversion underlying 8 of the SAPI items.

1. Having Fun
2. Being Liked by Others

We'll define our measurement model as the two-factor CFA shown to the right.



Example: Marker Variable

Load the SAPI data.

```
dataDir <- "data"
sapi <- read.table(here::here(dataDir, "sapi.txt"),
                  header = TRUE,
                  na.strings = "-999")
```

Specify the **lavaan** model syntax for the SAPI extraversion CFA.

```
mod1 <- '
fun    =~ Q77 + Q84 + Q170 + Q196
liked =~ Q44 + Q63 + Q76 + Q98
'
```

Use the `cfa()` function to estimate the model.

```
library(lavaan)
out1 <- cfa(mod1, data = sapi)
```

Example: Marker Variable

```
partSummary(out1, 1:4)
```

lavaan 0.6-19 ended normally after 30 iterations

Estimator	ML
Optimization method	NLMINB
Number of model parameters	17

	Used	Total
Number of observations	959	1000

Model Test User Model:

Test statistic	130.193
Degrees of freedom	19
P-value (Chi-square)	0.000

Example: Marker Variable

```
partSummary(out1, 5:7)
```

Parameter Estimates:

Standard errors	Standard
Information	Expected
Information saturated (h1) model	Structured

Latent Variables:

	Estimate	Std.Err	z-value	P(> z)
fun =~				
Q77	1.000			
Q84	0.761	0.051	14.902	0.000
Q170	0.634	0.047	13.558	0.000
Q196	0.795	0.046	17.381	0.000
liked =~				
Q44	1.000			
Q63	1.512	0.147	10.278	0.000
Q76	1.483	0.149	9.955	0.000
Q98	1.243	0.119	10.462	0.000

Example: Marker Variable

```
partSummary(out1, 8:9)
```

Covariances:

	Estimate	Std.Err	z-value	P(> z)
fun ~~				
liked	0.231	0.025	9.234	0.000

Variances:

	Estimate	Std.Err	z-value	P(> z)
.Q77	0.548	0.038	14.389	0.000
.Q84	0.727	0.039	18.703	0.000
.Q170	0.687	0.035	19.572	0.000
.Q196	0.364	0.025	14.731	0.000
.Q44	0.662	0.034	19.291	0.000
.Q63	0.807	0.048	16.943	0.000
.Q76	0.966	0.054	17.931	0.000
.Q98	0.469	0.029	16.121	0.000
fun	0.627	0.056	11.303	0.000
liked	0.182	0.029	6.290	0.000

Example: Marker Variable

```
inspect(out1, "r2")
```

```
   Q77   Q84  Q170  Q196   Q44   Q63   Q76   Q98  
0.534 0.333 0.268 0.521 0.215 0.340 0.293 0.374
```

```
fitMeasures(out1) |> head(22) |> round(3)
```

npar	fmin	chisq	df
17.000	0.068	130.193	19.000
pvalue	baseline.chisq	baseline.df	baseline.pvalue
0.000	1574.886	28.000	0.000
cfi	tli	nnfi	rfi
0.928	0.894	0.894	0.878
nfi	pnfi	ifi	rni
0.917	0.622	0.929	0.928
logl	unrestricted.logl	aic	bic
-10147.587	-10082.491	20329.175	20411.895
ntotal	bic2		
959.000	20357.903		

Example: Marker Variable

```
fitMeasures(out1) |> tail(-22) |> round(3)
```

rmsea	rmsea.ci.lower	rmsea.ci.upper
0.078	0.066	0.091
rmsea.ci.level	rmsea.pvalue	rmsea.close.h0
0.900	0.000	0.050
rmsea.notclose.pvalue	rmsea.notclose.h0	rmr
0.421	0.080	0.042
rmr_nomean	srmr	srmr_bentler
0.042	0.043	0.043
srmr_bentler_nomean	crmr	crmr_nomean
0.043	0.049	0.049
srmr_mplus	srmr_mplus_nomean	cn_05
0.043	0.043	223.037
cn_01	gfi	agfi
267.582	0.968	0.939
pgfi	mfi	ecvi
0.511	0.944	0.171

Example: Fixed Factor

We only need to change one option to implement the fixed-factor method.

- The `std.lv = TRUE` option (i.e., standardized latent variables) applies the appropriate constraints.

```
out2 <- cfa(mod1, data = sapi, std.lv = TRUE)
```

Example: Fixed Factor

```
partSummary(out2, 1:4)
```

lavaan 0.6-19 ended normally after 17 iterations

Estimator	ML	
Optimization method	NLMINB	
Number of model parameters	17	
	Used	Total
Number of observations	959	1000

Model Test User Model:

Test statistic	130.193
Degrees of freedom	19
P-value (Chi-square)	0.000

Example: Fixed Factor

```
partSummary(out2, 5:7)
```

Parameter Estimates:

Standard errors	Standard
Information	Expected
Information saturated (h1) model	Structured

Latent Variables:

	Estimate	Std.Err	z-value	P(> z)
fun =~				
Q77	0.792	0.035	22.606	0.000
Q84	0.603	0.035	17.193	0.000
Q170	0.502	0.033	15.180	0.000
Q196	0.630	0.028	22.308	0.000
liked =~				
Q44	0.426	0.034	12.580	0.000
Q63	0.644	0.040	16.071	0.000
Q76	0.632	0.043	14.845	0.000
Q98	0.530	0.031	16.912	0.000

Example: Fixed Factor

```
partSummary(out2, 8:9)
```

Covariances:

	Estimate	Std.Err	z-value	P(> z)
fun ~~				
liked	0.683	0.033	20.483	0.000

Variances:

	Estimate	Std.Err	z-value	P(> z)
.Q77	0.548	0.038	14.389	0.000
.Q84	0.727	0.039	18.703	0.000
.Q170	0.687	0.035	19.572	0.000
.Q196	0.364	0.025	14.731	0.000
.Q44	0.662	0.034	19.291	0.000
.Q63	0.807	0.048	16.943	0.000
.Q76	0.966	0.054	17.931	0.000
.Q98	0.469	0.029	16.121	0.000
fun	1.000			
liked	1.000			

Example: Fixed Factor

```
inspect(out2, "r2")
```

```
      Q77   Q84  Q170  Q196   Q44   Q63   Q76   Q98  
0.534 0.333 0.268 0.521 0.215 0.340 0.293 0.374
```

```
fitMeasures(out2) |> head(22) |> round(3)
```

npar	fmin	chisq	df
17.000	0.068	130.193	19.000
pvalue	baseline.chisq	baseline.df	baseline.pvalue
0.000	1574.886	28.000	0.000
cfi	tli	nnfi	rfi
0.928	0.894	0.894	0.878
nfi	pnfi	ifi	rni
0.917	0.622	0.929	0.928
logl	unrestricted.logl	aic	bic
-10147.587	-10082.491	20329.175	20411.895
ntotal	bic2		
959.000	20357.903		

Example: Fixed Factor

```
fitMeasures(out2) |> tail(-22) |> round(3)
```

rmsea	rmsea.ci.lower	rmsea.ci.upper
0.078	0.066	0.091
rmsea.ci.level	rmsea.pvalue	rmsea.close.h0
0.900	0.000	0.050
rmsea.notclose.pvalue	rmsea.notclose.h0	rmr
0.421	0.080	0.042
rmr_nomean	srmr	srmr_bentler
0.042	0.043	0.043
srmr_bentler_nomean	crmr	crmr_nomean
0.043	0.049	0.049
srmr_mplus	srmr_mplus_nomean	cn_05
0.043	0.043	223.037
cn_01	gfi	agfi
267.582	0.968	0.939
pgfi	mfi	ecvi
0.511	0.944	0.171

Compare Results

```
inspect(out1, "est")$lambda - inspect(out2, "est")$lambda
```

```
      fun liked  
Q77  0.208 0.000  
Q84  0.158 0.000  
Q170 0.132 0.000  
Q196 0.165 0.000  
Q44  0.000 0.574  
Q63  0.000 0.868  
Q76  0.000 0.851  
Q98  0.000 0.713
```

```
inspect(out1, "est")$psi - inspect(out2, "est")$psi
```

```
      fun liked  
fun    -0.373  
liked -0.453 -0.818
```

Compare Results

```
all.equal(fitMeasures(out1), fitMeasures(out2))
```

```
[1] TRUE
```

```
inspect(out1, "r2") - inspect(out2, "r2")
```

Q77	Q84	Q170	Q196	Q44	Q63	Q76	Q98
0	0	0	0	0	0	0	0

```
inspect(out1, "est")$theta - inspect(out2, "est")$theta
```

	Q77	Q84	Q170	Q196	Q44	Q63	Q76	Q98
Q77	0							
Q84	0	0						
Q170	0	0	0					
Q196	0	0	0	0				
Q44	0	0	0	0	0			
Q63	0	0	0	0	0	0		
Q76	0	0	0	0	0	0	0	
Q98	0	0	0	0	0	0	0	0