# Magic Gems: A Polyhedral Framework for Magic Squares

Kyle Elliott Mathewson
*Faculty of Science, University of Alberta*
kyle.mathewson@ualberta.ca

**Abstract**

A magic square can be read as an arrangement of vectors from the centroid whose vertical components cancel along every row, column and both diagonals. The *Magic Gem* of an $n \times n$ magic square is the convex hull of the tips $\mathbf{p}_{ij} = (x_{ij}, y_{ij}, z_{ij})$ of these vectors, where $(x_{ij}, y_{ij})$ is the centred cell and $z_{ij}$ the centred entry. We show that $\sum \mathbf{p}_{ij} = \mathbf{0}$ for every arrangement, so unweighted vector balance is not specific to magic squares; the magic-specific consequences are the height-weighted moments $\sum z\,\varphi(x) = \sum z\,\varphi(y) = 0$, which include Loly's invariant, while $\sum z\,xy$ is not forced. For order four we compute the 880 gems exactly (8 to 16 vertices, 39 $f$-vectors, 352 combinatorial types, 155 volumes, 616 congruence classes) and, by an exact computer-assisted calculation, determine all their isometry groups: the 48 associative squares give centrally symmetric gems, the 304 remaining self-complementary squares give gems with exactly one half-turn symmetry, and the other 528 gems have none; every gem is chiral except the 48 centrally symmetric ones. We also prove, with computer assistance, that the Siamese square of every odd order $n \geq 5$ has a gem with exactly twelve vertices.

**Keywords:** magic squares, convex polyhedra, symmetry, combinatorial geometry, moment of inertia

## 1 Introduction

Magic squares are among the oldest objects of recreational mathematics: the $3 \times 3$ Lo Shu belongs to Chinese tradition, a $4 \times 4$ square is engraved in Dürer's *Melencolia I* (1514), and constructions for every order have been known for centuries [2, 17]. The modern literature treats them from several directions: enumeration and the geometry of the polytopes whose lattice points are the squares [20, 8, 1, 4, 3], spectral and matrix properties [13, 15, 16, 5], and, in work of Loly and collaborators, physical invariants such as the moment of inertia and the multipole moments [11, 18, 19].

**Definition 1.1** (Magic square)**.** An *arrangement* of order $n$ is a bijection $S = (s_{ij})$ from the cells of the $n \times n$ grid onto $\{1, \ldots, n^2\}$. It is a *magic square* if every row, every column and both main diagonals sum to the magic constant $M(n) = n(n^2 + 1)/2$, and *semi-magic* if only the rows and columns do.

**The picture.** Centre the grid at the origin and attach to the cell $(i, j)$ the vector

$$\mathbf{p}_{ij} = (x_{ij}, y_{ij}, z_{ij}),$$

whose horizontal part $(x_{ij}, y_{ij})$ is the position of the cell and whose vertical part is the *centred entry* $z_{ij} = s_{ij} - \frac{n^2+1}{2}$. Then $S$ is magic exactly when the vertical components of these vectors cancel along every row, column and both diagonals (Lemma 2.2; Figure 1, Figure 2). The *Magic Gem* $\mathcal{G}(S)$ is the convex hull of the tips of the $n^2$ vectors $\mathbf{p}_{ij}$ (Section 2). Figure 1 shows the vectors of Dürer's square and the gem they span; the tips fill the gem's 12 vertices and four interior points, and $\sum \mathbf{p}_{ij} = \mathbf{0}$.

An interactive display of the 880 order-four gems is available at https://kylemath.github.io/MagicGemWebpage/.





Figure 1: Dürer's square as vectors from the centroid $O$. (a) The square as a heat map of the heights $z_{ij} = s_{ij} - 8.5$ (orange: above $O$, blue: below), with the entries $s_{ij}$ printed in the cells; every row, column and both diagonals sum to 34. (b) The 16 vectors $\mathbf{p}_{ij} = (x_{ij}, y_{ij}, z_{ij})$: each points from $O$ over the cell $(x_{ij}, y_{ij})$ of the floor grid (dotted drop lines) to a tip at height $z_{ij}$, labelled by the entry $s_{ij}$. (c) The same vectors inside their convex hull, the Magic Gem: 12 tips are vertices (filled dots) and 4 lie in the interior (open dots). The colour bar is shared by (a) and (b). The vectors sum to $\mathbf{0}$ for every arrangement of $1, \ldots, 16$ (Proposition 3.1); what singles out magic squares is how the heights cancel along lines (Figure 2).

**Results.**

1. *Balance* (Section 3). For every arrangement, magic or not, $\sum_{i,j} \mathbf{p}_{ij} = \mathbf{0}$ (Proposition 3.1), so the unweighted statement "the vectors balance about the centroid" carries no information about $S$; it concerns all $n^2$ points, interior ones included, not the solid hull. The magic-specific statements weight each vector by its height: $\sum z\,\varphi(x) = \sum z\,\varphi(y) = 0$ for every function $\varphi$ when $S$ is semi-magic, which contains the vanishing of the height-weighted centroid and Loly's invariant, while the mixed moment $\sum z\,xy$ is not forced (Proposition 3.2, Corollary 3.3).

2. *Symmetry of the order-four gems* (Section 4). The complement $S \mapsto 17 - S$ reflects the gem in the plane $z = 0$. We prove that a self-complementary square yields a gem with an explicit isometry (a point reflection if $S$ is associative, a half-turn about a horizontal axis otherwise), and determine by exact computation the complete isometry group of each of the 880 gems: 48 gems are centrally symmetric, 304 have exactly one half-turn symmetry, and 528 have no symmetry (Theorem 4.3). Facet orbits are classified (Proposition 4.4).

3. *Geometry of the* 880 *gems* (Section 5.1). Vertex counts, $f$-vectors, combinatorial types, volumes and congruence classes; the congruence classes are exactly the orbits of the complement map, and no two gems have the same vertex set.

4. *Siamese gems* (Proposition 5.2). The Siamese square of every odd order $n \geq 5$ has a gem with exactly twelve vertices, 26 edges and 16 facets (computer-assisted proof, Section B); the fraction of points that are vertices tends to zero.

**What is and is not new.** The polyhedral geometry of magic squares is classical: nonnegative integer matrices with all line sums equal to $t$ are the lattice points of the $t$-th dilate of the Birkhoff polytope, and their number is a polynomial in $t$ [20, 8, 22]; the count of magic squares with distinct entries is an Ehrhart quasi-polynomial of an inside-out polytope [4, 3]; the cones of magic squares

and cubes have Hilbert-basis descriptions [1]; contingency tables with fixed margins are the same lattice points seen as a sample space [6]. In all of these the magic squares are the *points* of a polytope, whereas the Magic Gem attaches a polyhedron to a *single* square, in the spirit of the physical invariants of Loly and Rogers [11, 18, 19] (see also [12] for Franklin squares). The balance statements of Section 3 are elementary and implicit in that work; we state and prove them in the gem's coordinates because the geometric picture invites misreadings of them. The symmetry classification, the exact geometry of the 880 gems and the Siamese theorem are new. The reading of the line conditions as orthogonality conditions on the height function, the hierarchy of moment conditions, the statistics of random arrangements and the enumeration of low-energy squares are developed in the companion paper [14]; none of its results is used here.

Section 2 defines the gem, Section 3 treats balance, Section 4 symmetry, Section 5 the exact geometry of the order-four and Siamese gems, Section 6 open questions, and the appendices contain the computational methods and the remaining steps of the Siamese proof.

## 2 The Magic Gem

Let $S = (s_{ij})$ be an arrangement of order $n$, with cells indexed by $i, j \in \{0, \dots, n-1\}$ ($i$ the row from the top, $j$ the column from the left), and write $N = n^2$.

**Definition 2.1** (Gem coordinates and Magic Gem)**.** To the cell $(i, j)$ assign the point $\mathbf{p}_{ij} = (x_{ij}, y_{ij}, z_{ij}) \in \mathbb{R}^3$ with

$$x_{ij} = j - \frac{n-1}{2}, \qquad y_{ij} = \frac{n-1}{2} - i, \qquad z_{ij} = s_{ij} - \frac{N+1}{2}. \tag{1}$$

The *Magic Gem* of $S$ is the convex hull $\mathcal{G}(S) = \operatorname{conv}\{\mathbf{p}_{ij}\}$. The function $z = (z_{ij})$ is the *height function* of $S$, and each $\mathbf{p}_{ij}$ is read as a vector from the origin. Every centroid or moment statement in this paper refers to the full set of $N$ points $\mathbf{p}_{ij}$, interior points included, and not to the solid $\mathcal{G}(S)$.

The $x$-axis runs along the columns and the $y$-axis along the rows (top row at positive $y$); the centred entries have mean zero because the mean entry is $(N+1)/2$. Let $x_j = j - \frac{n-1}{2}$ and $y_i = \frac{n-1}{2} - i$ denote the $n$ values taken by $x$ and $y$. A *line* is one of the $n$ rows, the $n$ columns, the main diagonal $\{i = j\}$ or the anti-diagonal $\{i + j = n-1\}$; each contains $n$ cells. For a line $L$ let $\delta_L(S) = \sum_{c \in L} s_c - M(n)$ be its *deviation*, so $S$ is magic iff $\delta_L(S) = 0$ for all $2n+2$ lines.

**Lemma 2.2** (Means, variances, line vector sums)**.** *For every arrangement $S$ of order $n$:*

1. $\sum_{i,j} x_{ij} = \sum_{i,j} y_{ij} = \sum_{i,j} z_{ij} = 0$ *and* $\sum_{i,j} x_{ij} y_{ij} = 0$*;*
2. *the empirical variances are* $\operatorname{Var}(x) = \operatorname{Var}(y) = \frac{n^2-1}{12}$ *and* $\operatorname{Var}(z) = \frac{n^4-1}{12}$*;*
3. *for every line $L$, $\delta_L(S) = \sum_{c \in L} z_c$, and*

$$\sum_{c \in L} \mathbf{p}_c = \big(n\,\bar{x}_L,\ n\,\bar{y}_L,\ \delta_L(S)\big),$$

*where $(\bar{x}_L, \bar{y}_L)$ is the centre of the line (the mean of its cell positions): $(0, y_i)$ for the row $i$, $(x_j, 0)$ for the column $j$, and $(0, 0)$ for both diagonals.*

*In particular $S$ is magic if and only if the vector sum of the $\mathbf{p}_c$ along every line is horizontal, i.e. the vertical components cancel along every row, column and diagonal.*

*Proof.* (1) $\sum_{i,j} x_{ij} = n \sum_j x_j = 0$, likewise for $y$; $\sum z = \sum_{v=1}^{N} v - N\frac{N+1}{2} = 0$; and $\sum xy = (\sum_j x_j)(\sum_i y_i) = 0$. (2) $\operatorname{Var}(x) = \frac{1}{n}\sum_j x_j^2$ is the variance of $n$ equally spaced points with unit spacing, $(n^2-1)/12$; the same holds for $y$, and for $z$ with $N$ points. (3) Since $M(n) = n\frac{N+1}{2}$, $\sum_{c \in L} z_c = \sum_{c \in L} s_c - M(n) = \delta_L$. The cells of a row $i$ have $y = y_i$ and $x$ running over all $x_j$ (sum 0); for a column the roles are exchanged; each diagonal is invariant under $(x, y) \mapsto (-x, -y)$, so its position sums vanish. □

Figure 2 illustrates Lemma 2.2(3) for Dürer's square: along each of four lines the heights stack head to tail and return to zero, so the vector sum of the line is horizontal, $(0,6,0) = 4\,(0,\tfrac{3}{2},0)$ for the top row; after swapping two entries the stack of the affected lines fails to close, by the deviation $\delta_L$.





Figure 2: Magic means that heights cancel along every line. *Top:* for Dürer's square, the four vectors of the top row, left column, main diagonal and anti-diagonal (coloured, with their entries; the other twelve in grey; the cells of the line are shaded on the floor) and below each the heights $z = s - 8.5$ of the line stacked head to tail in line order; the running total returns to 0, so $\sum_{c\in L} \mathbf{p}_c$ is the horizontal vector $(n\bar{x}_L, n\bar{y}_L, 0)$ of Lemma 2.2(3), here $(0,6,0)$, $(-6,0,0)$, $\mathbf{0}$, $\mathbf{0}$. *Bottom:* the same after swapping the entries 3 and 5, which is not magic: the top row and the left column no longer close (deviations $\delta_L = +2$ and $-2$, in magenta) and their vector sums acquire the vertical component $\delta_L$; the diagonals do not contain the swapped cells and still close.

**Example 2.3** (Lo Shu)**.** For the Lo Shu square $S = \begin{pmatrix} 2 & 7 & 6 \\ 9 & 5 & 1 \\ 4 & 3 & 8 \end{pmatrix}$, $n = 3$ and the centre value is 5,

so the nine points are

$$\begin{array}{lll} \mathbf{p}_{00}=(-1,1,-3), & \mathbf{p}_{01}=(0,1,2), & \mathbf{p}_{02}=(1,1,1),\\ \mathbf{p}_{10}=(-1,0,4), & \mathbf{p}_{11}=(0,0,0), & \mathbf{p}_{12}=(1,0,-4),\\ \mathbf{p}_{20}=(-1,-1,-1), & \mathbf{p}_{21}=(0,-1,-2), & \mathbf{p}_{22}=(1,-1,3). \end{array}$$

The top row has heights $-3+2+1=0$ and horizontal vector sum $(0,3,0)$. The centre cell maps to the origin and is the only point not on the hull; the gem has $f$-vector $(8,16,10)$ and volume 18 (Figure 3).

**Proposition 2.4** ($D_4$-invariance)**.** *Let $g \in D_4$ act on the grid and let $gS$ be the arrangement with $(gS)_c = s_{g^{-1}c}$. Then $\mathcal{G}(gS) = \hat{g}\,\mathcal{G}(S)$, where $\hat{g}(x,y,z) = (g(x,y),z)$ is the isometry of $\mathbb{R}^3$ that applies the orthogonal map $g$ of the $(x,y)$-plane and fixes $z$;* $\det \hat{g} = \det g$. *In particular $\mathcal{G}(gS)$ and $\mathcal{G}(S)$ are congruent, and the gem is an invariant of the $D_4$-orbit of $S$.*

*Proof.* In the centred coordinates (1), each $g \in D_4$ acts on the cell positions as an orthogonal map of the plane (a rotation by a multiple of $90^\circ$ or a reflection in a coordinate axis or a diagonal), and $(x_{gc}, y_{gc}) = g(x_c, y_c)$. The point of $gS$ over the cell $gc$ has height $(gS)_{gc} - \frac{N+1}{2} = z_c$. Hence the point set of $gS$ is $\{(g(x_c,y_c),z_c)\}_c = \hat{g}\{\mathbf{p}_c\}_c$, and an affine bijection maps the convex hull of a set onto the convex hull of its image. □

Magic squares are mapped to magic squares by $D_4$ (lines go to lines), so every *essentially different* magic square, that is, every $D_4$-orbit, has a single gem up to congruence.

## 3 Balance: what the vectors from the centroid do and do not say

**Unweighted balance holds for every arrangement.**

**Proposition 3.1** (Centroid)**.** *For every arrangement $S$ of order $n$, $\sum_{i,j} \mathbf{p}_{ij} = \mathbf{0}$: the origin is the centroid of the $N$ points, each of which is a vector from it. Moreover $\mathbf{0}$ lies in the interior of $\mathcal{G}(S)$.*

*Proof.* The sum vanishes by Lemma 2.2(1). The centroid of a finite point set that is not contained in a plane is an interior point of its convex hull, as a convex combination with all weights positive; the points $\mathbf{p}_{ij}$ span $\mathbb{R}^3$ since the positions $(x_{ij}, y_{ij})$ do and the heights are not all equal for $n \geq 2$. □

This is a consequence of centring the coordinates and carries no information about $S$; in particular it does not distinguish magic squares from arbitrary arrangements. It is a statement about the point set with unit weights, and not about the solid: the hull depends only on its vertex set, interior points do not affect it, and the centre of mass of the solid $\mathcal{G}(S)$ need not be the origin (for 829 of the 880 gems of order four it is not). The hull records the extent of the point set (extreme points, faces, volume), not how the heights are distributed over the cells.

**Height-weighted balance is what magic means.** Take the *signed mass* $m_{ij} = z_{ij}$ at the planar position $(x_{ij}, y_{ij})$, with total mass 0. Write $\rho_i = \sum_j z_{ij}$ and $\gamma_j = \sum_i z_{ij}$ for the row and column mass sums; by Lemma 2.2(3) they are the row and column deviations of $S$.

**Proposition 3.2** (Height-weighted moments)**.** *Let $S$ be an arrangement of order $n \geq 3$.*

1. *$S$ is semi-magic if and only if $\rho_i = 0$ for every row and $\gamma_j = 0$ for every column.*
2. *For every function $\varphi\colon \mathbb{R} \to \mathbb{R}$,*

$$\sum_{i,j} z_{ij}\,\varphi(x_{ij}) = \sum_j \varphi(x_j)\gamma_j, \qquad \sum_{i,j} z_{ij}\,\varphi(y_{ij}) = \sum_i \varphi(y_i)\rho_i.$$

*Hence if $S$ is semi-magic, $\sum z\,\varphi(x) = \sum z\,\varphi(y) = 0$ for every $\varphi$ (for $\varphi(t) = t^k$, all $k \geq 0$; for $k = 1$ the height-weighted centroid is the origin). Conversely, if $\sum z\,\varphi(x) = \sum z\,\varphi(y) = 0$ for all $\varphi$, then $S$ is semi-magic.*





Figure 3: Construction of the Magic Gem and its $D_4$-invariance. Top row: (a) the Lo Shu square $S$; (b) the centred cells $(x_{ij}, y_{ij})$ with the heights $z_{ij} = s_{ij} - 5$; (c) the nine points $\mathbf{p}_{ij}$ over the grid, labelled by the entries $s_{ij}$; (d) their convex hull, the gem of Example 2.3, with $f$-vector $(8, 16, 10)$; the centre point $(0, 0, 0)$ is the only interior point. Rows two and three (e–l): the eight images of the Lo Shu under $D_4$—the identity $e$, the rotations $r, r^2, r^3$ by $90°, 180°, 270°$, the left–right and top–bottom reflections $h, v$, and the two diagonal reflections (transpose $t$ and anti-transpose $tr^2$)—each printed row by row beneath its gem. All eight gems are drawn from the same viewpoint; by Proposition 2.4 they are the images of one another under the isometries $\hat{g}$, so they are congruent, and all have $f$-vector $(8, 16, 10)$ and volume 18.

*3. The mixed moment is not forced: for $n = 4$ there are magic squares with $\sum z\,xy \neq 0$.*

*Proof.* (1) By Lemma 2.2(3), a row sums to $M(n)$ if and only if its heights sum to 0; likewise for columns. (2) Grouping the cells by column, $\sum_{i,j} z_{ij}\varphi(x_j) = \sum_j \varphi(x_j)\gamma_j$; the row identity is the same with $y_i, \rho_i$. If $\gamma = \rho = 0$ both vanish. Conversely take $\varphi$ to be the indicator of a single value $x_{j_0}$ (the $x_j$ are distinct): then $\gamma_{j_0} = 0$; likewise $\rho_{i_0} = 0$, and (1) applies. (3) Since $\sum xy = 0$, $\sum z\,xy = \sum s\,xy = -P_{xy}$, the product of inertia of Loly's value-weighted planar body. Exact enumeration over the 880 essentially different magic squares of order 4 gives $|P_{xy}| \in \{0, \ldots, 60\}$ with 48 distinct values; it vanishes for 56 classes and is non-zero for the other 824. □

Figure 4 shows both halves of Proposition 3.2. In panel a the entries themselves are piles of checkers on a checkerboard, $s_{ij}$ checkers on cell $(i, j)$: every row and column carries 34 checkers, so the horizontal centre of mass of all the checkers lies over the centre $O$ ($\sum s\,x = \sum s\,y = 0$, which for $n = 4$ is the height-weighted statement because $\sum x = \sum y = 0$). Hence each column of heights has zero total mass and the lever of signed masses balances, while the mixed moment $\sum z\,xy$ takes the value 0 for Dürer's square and +60 for another magic square. The same weighted balance in

vector form is Figure 5.





Figure 4: Height-weighted moments, order $n = 4$. (a) Dürer's square as checkers on a $4 \times 4$ checkerboard, seen from the front right: the pile on cell $(i, j)$ has $s_{ij}$ checkers (labelled, and coloured by the entry $s_{ij}$ on the sequential scale shown); every row and column carries 34 checkers, and the centre of mass of all the checkers lies over the board centre $O$ (red), $\sum s\,x = \sum s\,y = 0$. (b) The heights with row sums $\rho_i$ and column sums $\gamma_j$, all zero. (c) Reading $z_{ij}$ as a force at position $x_j$ on a lever (up +, down −): every column has net force $\gamma_j = 0$, hence $\sum z\,x = \sum_j x_j \gamma_j = 0$ (Proposition 3.2(2)); the same holds for $y$ and the rows. (d) The mixed moment is not forced: for two magic squares, the heights $z$ and the cell contributions $z\,x\,y$ (rounded to integers), which sum to 0 for Dürer's square and to +60 for the square $\begin{smallmatrix} 1 & 6 & 11 & 16 \\ 8 & 15 & 2 & 9 \\ 12 & 3 & 14 & 5 \\ 13 & 10 & 7 & 4 \end{smallmatrix}$ (so $P_{xy} = -60$).

Thus the height-weighted statements guaranteed by the line sums are exactly the pure axial ones: the vanishing of the first moment along each axis and Loly's invariant below. The mixed moment can vanish for structural reasons: if $S$ is associative, $z$ is odd under $(x, y) \mapsto (-x, -y)$ (Section 4) and $xy$ is even, so $\sum z\,xy = 0$ (this accounts for the order-3 square and for 48 of the 56 vanishing classes of order 4).

**Corollary 3.3** (Loly's moment of inertia)**.** *For every arrangement,* $I(S) := \sum_{i,j} s_{ij}(x_{ij}^2 + y_{ij}^2) = \sum_j x_j^2 \gamma_j + \sum_i y_i^2 \rho_i + \frac{n^2(n^4-1)}{12}$. *In particular* $I(S) = n^2(n^4 - 1)/12$ *for every semi-magic square.*





Figure 5: The height-weighted vector sum $\sum_{i,j} z_{ij}\,(x_{ij}, y_{ij}) = (0, 0)$ for two magic squares: Dürer's square and square #45 of the 880 (the second square of Figure 4d). The 16 moment arms $z_{ij}(x_{ij}, y_{ij})$ (orange for $z > 0$, blue for $z < 0$) are laid head to tail in order of direction; the chain closes at $O$ in both cases, which is $\sum z\,x = \sum z\,y = 0$ (Proposition 3.2(2)). The right-hand tables give $\sum z\,xy$ for the four quadrants of the grid; they add to 0 for Dürer's square and to +60 for square #45, so the weighted balance does not determine the mixed moment (Proposition 3.2(3)).

*Proof.* Substitute $s_{ij} = z_{ij} + \frac{N+1}{2}$. The first part is $\sum z\,(x^2 + y^2)$, equal to $\sum_j x_j^2\gamma_j + \sum_i y_i^2\rho_i$ by Proposition 3.2(2); the second is $\frac{N+1}{2}\sum(x^2+y^2) = \frac{N+1}{2}\cdot 2N\frac{n^2-1}{12} = \frac{n^2(n^4-1)}{12}$ by Lemma 2.2(2). □

This is Loly's invariance [11], which he derived from the row and column sums. For $n = 3, \ldots, 9$ the value is $60, 340, 1300, 3885, 9800, 21840, 44280$.

*Remark* 3.4 (Inertia tensor of the unit-mass point set). With unit masses at the $N$ points, the centre of mass is the origin and the inertia tensor about it is $I_{jk} = N(\delta_{jk}\,\mathrm{tr}\,\Sigma - \Sigma_{jk})$, where $\Sigma$ is the covariance matrix of the points [10]. By Lemma 2.2 and Proposition 3.2(2), $I_{xx} = I_{yy} = n^2(n^2-1)(n^2+2)/12$, $I_{zz} = n^2(n^2-1)/6$, $I_{xy} = 0$, $I_{xz} = -\sum_j x_j\gamma_j$ and $I_{yz} = -\sum_i y_i\rho_i$. The diagonal entries are the same for all $N!$ arrangements and the tensor is diagonal for every semi-magic square, so the inertia of the unit-mass point set does not depend on the square. The informative planar object is Loly's value-weighted body: its diagonal entries are fixed by the row and column sums (Corollary 3.3), whereas the product of inertia $P_{xy} = -\sum s_{ij}x_{ij}y_{ij}$ is a mixed moment, fixed by no line sum; $D_4$ changes at most its sign.

## 4 Symmetry of the gems

Let $\rho(x, y, z) = (x, y, -z)$ be the reflection in the plane $z = 0$. The *complement* of an arrangement is $\overline{S} = N + 1 - S$ (so $\overline{S}$ has height function $-z$), and $S$ is *associative* if $S + r^2S = N + 1$, where $r^2$ is the half-turn of the grid; $S$ is *self-complementary* if $\overline{S} = gS$ for some $g \in D_4$. Complements of magic squares are magic.

**Lemma 4.1** (Complement is a mirror image)**.** *$\mathcal{G}(\overline{S}) = \rho\,\mathcal{G}(S)$. If $\overline{S} = gS$ with $g \in D_4$, then $\iota_g := \hat{g}^{-1}\rho$, i.e. $\iota_g(x, y, z) = (g^{-1}(x, y), -z)$, is an isometry of $\mathbb{R}^3$ mapping $\mathcal{G}(S)$ onto itself, with* $\det \iota_g = -\det g$.

*Proof.* The points of $\overline{S}$ are $(x_c, y_c, -z_c) = \rho\,\mathbf{p}_c$. If $\overline{S} = gS$ then $\rho\,\mathcal{G}(S) = \mathcal{G}(\overline{S}) = \mathcal{G}(gS) = \hat{g}\,\mathcal{G}(S)$ by Proposition 2.4, so $\hat{g}^{-1}\rho\,\mathcal{G}(S) = \mathcal{G}(S)$. □

**Proposition 4.2** (Which elements of $D_4$ can complement)**.** *Let $\overline{S} = gS$ for an arrangement $S$ of order $n \geq 2$. Then $g^2 = \mathrm{id}$ and $g$ has no fixed cell other than possibly the centre cell of an odd grid, whose entry is then $(N+1)/2$. Consequently:*

1. *for $n$ odd, $g = r^2$, so $S$ is associative;*
2. *for $n$ even, $g \in \{r^2, h, v\}$ with $h, v$ the left–right and top–bottom reflections; the diagonal reflections and the quarter-turns never occur.*

*The corresponding isometries of $\mathcal{G}(S)$ are $\iota_{r^2}(\mathbf{p}) = -\mathbf{p}$ (point reflection, improper), $\iota_h(x, y, z) = (-x, y, -z)$ (half-turn about the $y$-axis, proper) and $\iota_v(x, y, z) = (x, -y, -z)$ (half-turn about the $x$-axis, proper).*

*Proof.* Applying $g$ to $\overline{S} = gS$ gives $g\overline{S} = \overline{gS} = S$, so $g^2 S = \overline{\overline{S}} = S$; since the entries are distinct, $g^2$ fixes every cell, hence $g^2 = \mathrm{id}$ ($D_4$ acts faithfully on the cells for $n \geq 2$). If $g$ fixes a cell $c$ then $s_c = N + 1 - s_c$. The involutions of $D_4$ are $r^2$, $h$, $v$ and the two diagonal reflections. The diagonal reflections fix $n \geq 2$ cells and are excluded; $h$ and $v$ fix the $n$ cells of the middle column (row) when $n$ is odd, and none when $n$ is even; $r^2$ fixes only the centre cell when $n$ is odd. For $n$ odd the centre cell is the only cell that can have the value $(N+1)/2$, so $g = r^2$. The formulas for $\iota_g$ follow from Lemma 4.1, and the determinants are $-\det r^2 = -1$, $-\det h = -\det v = +1$. □

By Proposition 4.2, associative squares give gems with a point reflection: *centrally symmetric* gems. (Every Siamese square is associative, which explains the symmetric vertex list in Proposition 5.2; Dürer's square is associative and its gem is centrally symmetric, Figure 6a, Figure 7a.) Self-complementary squares that are not associative give gems with a half-turn symmetry. For order four we determine the complete isometry group of every gem. For a finite set $X \subset \mathbb{R}^3$ let $\mathrm{Iso}(X)$ be the group of isometries of $\mathbb{R}^3$ mapping $X$ to itself.

**Theorem 4.3** (Isometry groups of the order-four gems; computer-assisted)**.** *Let $S$ be one of the* 880 *essentially different magic squares of order* 4 *and $\mathcal{G} = \mathcal{G}(S)$. Then $\mathrm{Iso}(\mathcal{G}) = \mathrm{Iso}(V) = \mathrm{Iso}(\{\mathbf{p}_{ij}\})$, where $V$ is the vertex set, and:*

1. *$S$ is associative (*48 *squares, Dudeney's group III) if and only if $\mathrm{Iso}(\mathcal{G}) = \{\mathrm{id}, -\mathrm{id}\}$ (the group $C_i$);*
2. *$S$ is self-complementary but not associative (*304 *squares, Dudeney's group VI) if and only if $\mathrm{Iso}(\mathcal{G}) = \{\mathrm{id}, \iota\}$ with $\iota$ a half-turn about a horizontal coordinate axis (the group $C_2$);*
3. *for the remaining* 528 *squares $\mathrm{Iso}(\mathcal{G})$ is trivial.*

*Consequently $\mathcal{G}(S)$ has an improper isometry if and only if $S$ is associative; the* 832 *gems of the cases (2) and (3) are chiral; and the congruence classes of the* 880 *gems number* 48+304+264 = 616*, the* 528 *gems of case (3) forming* 264 *pairs $\{\mathcal{G}(S), \mathcal{G}(\overline{S})\}$ of mirror images in the plane $z = 0$.*

*Proof.* The inclusions "$\supseteq$" and the identification of the generators for the associative and self-complementary squares are Lemma 4.1 and Proposition 4.2; the classes (1), (2) are exactly the self-complementary ones, which were counted as 48 and 304 (all of the form $h$ or $v$ after Proposition 4.2) by exact comparison of the 8 images of $S$ with $\overline{S}$. For the reverse inclusions we computed, for each of the 880 squares, *all* isometries of its vertex set and of its 16-point set, in exact integer arithmetic: the vertex set is centred at its centroid (scaled to integers), the images of three linearly independent vertices are enumerated among the vertices with the same Gram matrix entries, each candidate orthogonal map is accepted only if the induced bijection of the vertices preserves all pairwise integer squared distances (Section A). This yields the group orders $2, 2, 1$ in the three cases, equal for the vertex set and the 16-point set; since $\mathcal{G}$ is the convex hull of $V$, $\mathrm{Iso}(\mathcal{G}) = \mathrm{Iso}(V)$ (an isometry preserves extreme points). A gem is chiral if it has no improper isometry; the group $C_2$ consists of proper maps, which gives the chirality statement. Finally, congruent gems come from the same complement orbit (Proposition 5.1), and in case (3) the classes of $S$ and $\overline{S}$ are distinct, giving 264 pairs. □

Table 1: Symmetry types of the 880 order-four gems. $V$: number of hull vertices; "classes" counts congruence classes of gems.

| type of $S$ | Dudeney | squares | $\mathrm{Iso}(\mathcal{G})$ (order) | classes | $V$ |
|---|---|---|---|---|---|
| associative | III | 48 | $C_i$, inversion (2) | 48 | $10, 12, 14, 16$ |
| self-complementary, not associative | VI | 304 | $C_2$, half-turn (2) | 304 | $8, 10, 12, 14, 16$ |
| neither | others | 528 | trivial (1) | 264 | $8, \ldots, 16$ |
| total | | 880 | | 616 | |

**Proposition 4.4** (Facet orbits)**.** *Let $S$ be one of the* 880 *squares and let* $\mathrm{Iso}(\mathcal{G})$ *act on the facets of $\mathcal{G}$.*

1. *If $S$ is associative, every facet $F$ has a distinct antipodal partner $-F$, a facet in the parallel plane on the other side of the origin; so there are no fixed facets and the facets form $F/2$ antipodal pairs. Here $F \in \{12, 14, 16, 18, 20, 22\}$ with $1, 12, 12, 17, 4, 2$ gems respectively, and $V \in \{10, 12, 14, 16\}$ with $13, 28, 4, 3$ gems. The partner $-F$ has the opposite outward normal; as plane polygons $F$ and $-F$ are congruent by a rotation through $\pi$, although the point reflection carrying $F$ to $-F$ is improper.*

2. *If $S$ is self-complementary and not associative, with half-turn $\iota$ about the $y$-axis (say), the facet orbits have size* 1 *or* 2*. A facet is fixed if and only if it lies in $y = \pm\frac{3}{2}$; such a facet is a parallelogram (the hull of the top or bottom row), and exists unless the four points of that row are collinear. There are two fixed facets for* 282 *gems, one for* 18 *and none for* 4*.*

3. *The vertices of a gem with a symmetry fall into orbits of size* 2*, so $V$ is even for the* 352 *self-complementary squares.*

*Proof.* Both nontrivial isometries fix the origin, which is an interior point of $\mathcal{G}$ (Proposition 3.1), so no facet plane $\{\mathbf{n} \cdot \mathbf{p} = d\}$ has $d = 0$. (1) $-F \neq F$ because $-$ maps the facet plane $\mathbf{n} \cdot \mathbf{p} = d$ to $\mathbf{n} \cdot \mathbf{p} = -d \neq d$. (2) The half-turn $\iota = (x, y, z) \mapsto (-x, y, -z)$ maps the plane $\mathbf{n} \cdot \mathbf{p} = d$ to $\iota\mathbf{n} \cdot \mathbf{p} = d$, which is the same plane iff $\iota\mathbf{n} = \mathbf{n}$, i.e. $\mathbf{n} \parallel (0, 1, 0)$; the facet planes with this normal are $y = \pm\frac{3}{2}$. The hull of the top row (the points with $y = \frac{3}{2}$) is $\iota$-invariant; its vertices come in pairs $\pm(x, z)$ and a convex polygon with at most four vertices and a centre of symmetry is a segment or a parallelogram. The counts are those of the exact computation. (3) $\iota$ and $-\mathrm{id}$ fix no vertex: the half-turn fixes only points with $x = z = 0$, while $x$ is a half-odd integer; the point reflection fixes only the origin. □

Figure 7 shows the same three squares as point sets in the height picture: for the associative square the 16 points form 8 antipodal pairs $\{\mathbf{p}, -\mathbf{p}\}$ whose segments all pass through $O$; for the self-complementary square the half-turn about the dashed horizontal axis exchanges the 8 pairs, each segment meeting the axis perpendicularly at its midpoint; the third square has no such pairing, in agreement with Theorem 4.3.

# 5 Exact geometry: the order-four gems and the Siamese gems

All hull computations in this section are exact (integer or rational arithmetic); the methods are in Section A, the scripts and data in the ancillary directory `anc/`.

## 5.1 Geometry of the 880 order-4 gems

There are 8 magic squares of order 3, all images of the Lo Shu under $D_4$, and 7,040 of order 4, forming 880 $D_4$-orbits; the order-four squares are attributed to Frénicle de Bessy (his list was published in 1693) and have been recounted by computer [3]. Dudeney sorted them into twelve groups according to the pattern of complementary pairs $(a, 17 - a)$ [7]; we use his groups only as a





Figure 6: Symmetry types of order-four gems (same viewpoint; interior points in red; the square is printed above each gem). (a) Dürer's square: a centrally symmetric gem; an antipodal pair of facets $F$ (orange) and $-F$ (green) and the segment joining their centroids through the centre $\mathbf{0}$ (dashed). (b) A self-complementary, non-associative square (square #124): the half-turn $\iota$ about the dashed axis swaps the highlighted pair of facets and has no other symmetry. (c) A square with no symmetry (square #102): the gem is chiral, and its mirror image in $z = 0$ is the gem of the complement $17 - S$.





Figure 7: The 16 points $\mathbf{p}_{ij}$ as spheres on stems over the floor grid, coloured by height $z$ (same squares and viewpoint as Figure 6; $O$ red). (a) Dürer's square: the points form 8 antipodal pairs $\{\mathbf{p}, -\mathbf{p}\}$, joined by segments through $O$. (b) Square #124: the half-turn $(x, y, z) \mapsto (x, -y, -z)$ about the dashed $x$-axis exchanges 8 pairs of points, each pair joined by a segment perpendicular to the axis; this gem is chiral. (c) Square #102: no non-trivial isometry preserves the point set.

classical reference partition. The number 275,305,224 of essentially different order-five squares is due to Schroeppel (1973) [9]; for $n \geq 6$ exact counts are, to the author's knowledge, not available (see Trump's compilation [21]).

For each of the 880 essentially different order-4 magic squares we computed the Magic Gem $\mathcal{G}(S)$ exactly (integer facet planes, merged coplanar triangles, rational volumes; Section A); Euler's relation holds for all 880 gems. Figure 8 and Table 2 summarise the results.

Table 2: Face structure of the 880 order-4 gems by number of hull vertices $V$: number of squares, of distinct combinatorial types (face lattices up to isomorphism), of distinct exact volumes, and of distinct $f$-vectors $(V, E, F)$. The full list of the 39 $f$-vectors with multiplicities is in the ancillary file `anc/results_geometry.json`.

| $V$ | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | all |
|---|---|---|---|---|---|---|---|---|---|---|
| squares | 7 | 10 | 103 | 80 | 380 | 106 | 162 | 14 | 18 | 880 |
| types | 4 | 1 | 27 | 32 | 129 | 52 | 85 | 7 | 15 | 352 |
| volumes | 4 | 5 | 51 | 33 | 111 | 40 | 65 | 6 | 14 | 155 |
| $f$-vectors | 4 | 1 | 6 | 5 | 7 | 5 | 5 | 3 | 3 | 39 |

**Vertex counts, $f$-vectors, volumes.** Between 8 and all 16 of the points are hull vertices; $V = 12$ is modal (380/880). There are 39 distinct $f$-vectors, from $(8, 14, 8)$ (one gem, square #54) and $(8, 16, 10)$ (one gem) to $(16, 36, 22)$ (nine gems), the modal one being $(12, 26, 16)$ (164 gems); the multiset of facet sizes (number of triangles, quadrilaterals, ...) is determined by the $f$-vector in every case. The exact volume takes 155 distinct values, from 58 (square #427) to 292/3 (square #379), mean 84.05. The denominators are $1, 2, 3, 6$: all three coordinates of every point are half-odd integers when $n$ is even, so the translate of a gem by $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ is a lattice polytope of $\mathbb{Z}^3$ and its volume lies in $\frac{1}{6}\mathbb{Z}$. Surface areas range from 116.5 to 148.6.

**Combinatorial types.** Two gems have the same combinatorial type if their vertex–facet incidence structures (equivalently, their face lattices) are isomorphic. Grouping by a Weisfeiler–Lehman hash of the bipartite vertex–facet graph and confirming every group by an explicit graph isomorphism test gives

352 combinatorial types

among the 880 gems (the hash was already exact: no hash class had to be split). Their sizes are very uneven; the number of types with $1, 2, 3, \ldots$ members is

| squares per type | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 10 | 13 | 14 | 16 | 26 | 32 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| types | 93 | 201 | 8 | 20 | 4 | 7 | 4 | 5 | 5 | 1 | 1 | 1 | 1 | 1 |

(Figure 8c); the largest type, with $f = (10, 22, 14)$ (12 triangles, 2 quadrilaterals), is shared by 32 squares, and the second largest, $f = (12, 24, 14)$, by 26. The 7 gems with 8 vertices fall into 4 types, including a single $(8, 14, 8)$ gem with four triangular and four quadrilateral faces (square #54, Fig. 8d) and four $(8, 18, 12)$ gems sharing one simplicial type with 12 triangular faces. The 18 gems with all 16 points extreme belong to 15 types.

**Congruence classes and injectivity.** Two gems are *congruent* if an isometry of $\mathbb{R}^3$ (rotations, reflections, translations) maps one onto the other; since a distance-preserving bijection between finite point sets extends to an isometry, congruence is decided exactly by testing for a bijection of hull-vertex sets preserving all integer squared distances (Section A).

**Proposition 5.1** (class → gem)**.** *1. The* 880 *hull-vertex sets are pairwise distinct: the map from $D_4$-classes of order-*4 *magic squares to hull-vertex sets (as subsets of $\mathbb{R}^3$) is injective. (For the full* 16*-point sets this is trivial, since the point set determines the arrangement.)*

2. *Up to congruence there are exactly* 616 *gems. The congruence classes are precisely the orbits of the complement map* $S \mapsto 17-S$*, which acts on gems as the reflection* $z \mapsto -z$*:* 352 *squares are self-complementary (their gems have a non-trivial isometry, Theorem 4.3) and the other* 528 *form* 264 *mirror pairs. No two gems of non-complementary squares are congruent, not even as full* 16*-point sets.*

Part (2) is the only failure of injectivity, and it is the expected one: $17 - S$ is magic whenever $S$ is, and its gem is the mirror image of $\mathcal{G}(S)$ in the plane $z = 0$. The 352 self-complementary classes are exactly Dudeney's group III (the 48 associative squares) together with all 304 squares of group VI [7]. The coincidence $352 = \#\{\text{types}\} = \#\{\text{self-complementary squares}\}$ is numerical only: the two partitions of the 880 are different.

**Dudeney groups.** Crossing the hull-vertex count with Dudeney's twelve groups (table in the ancillary files) shows that geometry is only loosely tied to Dudeney type: every group with more than eight members spans at least three vertex counts and at least 20 combinatorial types; the 8-vertex gems occur only in groups VI (three) and VIII (four).

The groups III and VI, the self-complementary ones, have only even vertex counts; this is Proposition 4.4(3).





Figure 8: Geometry of the 880 order-4 Magic Gems. (a) Number of hull vertices (8 to 16; modal value 12). (b) Exact hull volume (155 distinct values between 58 and 292/3). (c) Sizes of the 352 combinatorial types (logarithmic scale): 93 types occur once, 201 twice, and one type is shared by 32 squares. (d)–(f) Three extreme gems, drawn with translucent faces after merging coplanar triangles: the unique $(8, 14, 8)$ gem (square #54; the eight interior points are shown in red), a gem with all 16 points extreme (square #802, Dürer's square with its first two rows exchanged and its last two rows exchanged), and the gem of largest volume (square #379).

**Named squares.** The gems of the Lo Shu, of Dürer's square and of the Siamese square of order 5 have hull fractions 8/9, 12/16 and 12/25 and exact volumes 18, 92 and 830/3. Dürer's own gem is centrally symmetric with twelve vertices, but exchanging its first two rows and its last two rows (which preserves the magic property) gives a square whose gem has all sixteen points extreme

(square #802, Figure 8e): the hull-vertex count is a property of the square, not of the Dudeney group or of the construction method.

### 5.2 Siamese gems have twelve vertices

For the Siamese (de la Loubère) squares of odd order the hull-vertex count does not grow with $n$. We computed the gems for $n = 3, 5, \ldots, 21$: the Lo Shu gem has $f = (8, 16, 10)$, and every Siamese gem with $n \geq 5$ has $f = (12, 26, 16)$ with 12 triangular and 4 quadrilateral facets, so the hull fraction $12/n^2$ tends to zero. This is a theorem, with a short computer-assisted proof.

**Proposition 5.2** (Siamese gems)**.** *Let $n = 2m + 1 \geq 5$ and let $S$ be the Siamese square of order $n$ (1 in the middle cell of the top row, successive entries one step up and to the right modulo $n$, one step down when blocked). Then $\mathcal{G}(S)$ has exactly 12 vertices, 26 edges and 16 facets (12 triangles, 4 quadrilaterals). With 0-indexed cells $(i, j)$ and $c = (n^2 + 1)/2$, the vertices are the points over the cells*

$$\begin{array}{lll} (0,0)\colon c+m+2, & (0,m-1)\colon n^2-1, & (0,m)\colon 1, \\ (0,2m)\colon c+m, & (m-1,0)\colon n^2-m, & (m,0)\colon m+2, \\ (m,2m)\colon n^2-m-1, & (m+1,2m)\colon m+1, & (2m,0)\colon c-m, \\ (2m,m)\colon n^2, & (2m,m+1)\colon 2, & (2m,2m)\colon c-m-2, \end{array}$$

*i.e. the four corners, and on each side of the square the two cells holding the side's extreme values. For $n = 3$ the same construction gives the Lo Shu gem, with 8 vertices (all cells except the centre).*

*Proof. Step 1: closed form.* Write $\langle t \rangle \in \{0, \ldots, n-1\}$ for $t$ mod $n$. We claim $s(i, j) = n\langle i + j - m \rangle + \langle i + 2j - 2m \rangle + 1$. Indeed, along a run of $n$ consecutive entries the move $(i, j) \to (i-1, j+1)$ fixes $i+j$ and increases $i+2j$ by 1 modulo $n$, while the "blocked" move $(i, j) \to (i+1, j)$ increases both $i + j$ and $i + 2j$ by 1; the entry 1 sits at $(0, m)$ where both expressions vanish. Hence by induction the entry $k$ lies in a cell with $\langle i+j-m \rangle = \lfloor (k-1)/n \rfloor$ and $\langle i+2j-2m \rangle = \langle k-1 \rangle$; since $(i, j) \mapsto (i + j,\, i + 2j)$ is a bijection of $\mathbb{Z}_n^2$ (determinant 1), this determines the cell, which proves the claim.

*Steps 2 and 3 (sketch; details in Section B).* With $m = (n-1)/2$, $x = j - m$, $y = m - i$, the closed form gives $z = L(x, y) + n\,\zeta(x, y)$ with $L$ affine and $\zeta = -n\lfloor (x-y+m)/n \rfloor - \lfloor (2x-y+m)/n \rfloor \in \{\pm(n+1), \pm 1, 0\}$. The affine bijection $\Phi(x, y, z) = (x, y, (z-L)/n)$ preserves the face lattice and the vertex cells, and $\Phi(\mathcal{G}(S))$ is the convex hull of the lattice points of five convex lattice polygons at the five heights $\zeta$: two triangles $T$, $B = -T$ at $\zeta = \pm(n+1)$, two triangles $U$, $D = -U$ at $\zeta = \pm 1$, and a parallelogram $M$ at $\zeta = 0$ (Figure 9a). Sixteen explicit planes, eight of them and their images under $(x, y, \zeta) \mapsto -(x, y, \zeta)$, are shown to support the convex hull of twelve named points $v_1, \ldots, v_{12}$—the three corners of $T$ and of $B$, one corner each of $U$ and $D$, and the four corners of $M$—in facets that close up into a surface with $(V, E, F) = (12, 26, 16)$ closed under edge-adjacency, hence all facets (so $P = Q$), and every vertex of every layer polygon satisfies all sixteen inequalities (Figure 9b). These are finitely many polynomial identities and inequalities in $m$, verified exactly for all integers $m \geq 2$ by the script `siamese_proof.py`. Hence $\Phi(\mathcal{G}(S)) = \mathrm{conv}\{v_1, \ldots, v_{12}\}$, and mapping back by $\Phi^{-1}$ gives the statement; for $m = 1$ the four triangles degenerate to points and the same argument gives the 8-vertex Lo Shu gem. □

## 6 Discussion and open questions

The Magic Gem is the convex hull of a magic square's height function over the centred grid, and the same polyhedron for all eight $D_4$-images of the square. The vector picture is exact but easily over-read: the centroid statement belongs to the full point set and holds for every arrangement; the magic conditions are the height-weighted ones; and the hull records only extent (Section 3). What the hull does see is symmetry: the complement map acts as a mirror reflection of the gem, and for order four this accounts for all the symmetry there is (Theorem 4.3) and for all coincidences between





Figure 9: The Siamese gem of order $n = 9$ after the affine reduction of Proposition 5.2. (a) The 81 cells in the $(x, y)$-plane coloured by the layer value $\zeta \in \{+10, +1, 0, -1, -10\}$, with the five lattice polygons $T, U, M, D, B$ outlined and the twelve vertex cells $v_1, \ldots, v_{12}$ circled. (b) The reduced polytope $\Phi(\mathcal{G}(S)) = \mathrm{conv}\{(x, y, \zeta)\}$ with its 12 vertices and 16 facets (12 triangles, 4 quadrilaterals); the shapes of the layers do not depend on $n$ except through scale, which is why the combinatorial type is constant for all odd $n \geq 5$.

gems (Proposition 5.1). The coarse invariants, the vertex count (nine values) and the volume (155 values), are not well aligned with Dudeney's groups, and the vertex count is a property of the square, not of its group or construction (Dürer's square versus square #802).

**Open questions.**

1. *Order five.* By Schroeppel's enumeration there are 275,305,224 essentially different squares of order 5 [9]; an exact hull computation for each is feasible. How many combinatorial types occur, what is the modal $f$-vector, and is the complement map again the only source of congruence? By Proposition 4.2 the self-complementary squares of odd order are exactly the associative ones, so every symmetric gem of order five is centrally symmetric.
2. *Symmetry in general.* Is $\mathrm{Iso}(\mathcal{G}(S))$ generated by the complement symmetry for every magic square of every order, as it is for order four?
3. *Bounded hull-vertex counts.* Siamese gems have twelve vertices for every odd $n \geq 5$ because, affinely, they are hulls of five lattice polygons whose shapes do not depend on $n$ (Proposition 5.2). Do the LUX, Strachey or diagonal-complement constructions have the same property, and is the fraction of vertices for a uniformly random magic square of large order bounded away from zero?
4. *Other square families.* The construction applies verbatim to semi-magic, pandiagonal and Franklin squares [12]; the statements of Proposition 3.2 already cover the semi-magic case. The gems of these families have not been examined.

# References


[1] Maya M. Ahmed, Jesús A. De Loera, and Raymond Hemmecke. Polyhedral cones of magic cubes and squares. In *Discrete and Computational Geometry*, volume 25 of *Algorithms and Combinatorics*, pages 25–41. Springer, 2003.

[2] W. S. Andrews. *Magic Squares and Cubes.* Dover Publications, 2nd edition, 1960.

[3] Matthias Beck and Andrew van Herick. Enumeration of $4 \times 4$ magic squares. *Mathematics of Computation*, 80(273):617–621, 2011.

[4] Matthias Beck and Thomas Zaslavsky. Inside-out polytopes. *Advances in Mathematics*, 205(1):134–162, 2006.

[5] Ian Cameron, Adam Rogers, and Peter D. Loly. Signatura of magic and Latin integer squares: isentropic clans and indexing. *Discussiones Mathematicae Probability and Statistics*, 33(1-2):121–149, 2013.

[6] Persi Diaconis and Anil Gangolli. Rectangular arrays with fixed margins. In *Discrete Probability and Algorithms (Minneapolis, MN, 1993)*, volume 72 of *The IMA Volumes in Mathematics and its Applications*, pages 15–41. Springer, New York, 1995.

[7] Henry E. Dudeney. *Amusements in Mathematics*. Thomas Nelson and Sons, London, 1917.

[8] Eugène Ehrhart. Sur les carrés magiques. *Comptes Rendus de l'Académie des Sciences, Série A*, 277:651–654, 1973.

[9] Martin Gardner. Mathematical games: A breakthrough in magic squares, and the first perfect magic cube. *Scientific American*, 234(1):118–123, 1976.

[10] Herbert Goldstein, Charles P. Poole, and John L. Safko. *Classical Mechanics*. Addison-Wesley, San Francisco, 3rd edition, 2002.

[11] Peter D. Loly. The invariance of the moment of inertia of magic squares. *The Mathematical Gazette*, 88(511):151–153, 2004. Note 88.30.

[12] Peter D. Loly. Franklin squares: A chapter in the scientific studies of magical squares. *Complex Systems*, 17(2):143–157, 2007.

[13] Peter D. Loly, Ian Cameron, Walter Trump, and Daniel Schindel. Magic square spectra. *Linear Algebra and its Applications*, 430(10):2659–2680, 2009.

[14] Kyle Elliott Mathewson. Magic squares as orthogonality conditions on a height function, 2026. In preparation.

[15] R. P. Nordgren. On properties of special magic square matrices. *Linear Algebra and its Applications*, 437(8):2009–2025, 2012.

[16] R. P. Nordgren. Spectra of order-4 special magic squares. *Linear Algebra and its Applications*, 455:320–331, 2014.

[17] Clifford A. Pickover. *The Zen of Magic Squares, Circles, and Stars*. Princeton University Press, 2002.

[18] Adam Rogers and Peter D. Loly. The inertia tensor of a magic cube. *American Journal of Physics*, 72(6):786–789, 2004.

[19] Adam Rogers and Peter D. Loly. The electric multipole expansion for a magic cube. *European Journal of Physics*, 26(5):809–813, 2005.

[20] Richard P. Stanley. Linear homogeneous Diophantine equations and magic labelings of graphs. *Duke Mathematical Journal*, 40:607–632, 1973.

[21] Walter Trump. Notes on magic squares and cubes. `https://www.trump.de/magic-squares/`.

[22] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer, New York, 1995.

# A Computational methods

**Hardware and software.** All computations were run on an Apple M4 Max (16 cores, 128 GB) under macOS, with Python 3.9.6, NumPy 2.0.2, SciPy 1.13.1 (Quickhull only seeds the exact hull computation) and NetworkX 3.2.1. The scalar identities of Section 3 (the value $n^2(n^4-1)/12$ for the semi-magic squares and the product-of-inertia statistics over the 880 classes) were checked in exact integer arithmetic.

**Exact hulls and their classification.** Doubling all coordinates makes every point of a gem an integer vector. Facet planes are recovered from an initial Quickhull triangulation as primitive integer equations, coplanar triangles are merged into faces, vertices are the points on at least three facet planes, edges are vertex pairs sharing two facets, and volumes are exact rationals; Euler's relation $V-E+F=2$ is checked for every gem. Combinatorial types are the isomorphism classes of the bipartite vertex–facet incidence graphs, grouped by a Weisfeiler–Lehman hash and confirmed by explicit isomorphism tests. Isometry groups (Theorem 4.3) are computed exactly as follows: the vertex set $V$ ($k$ points, doubled integer coordinates) is centred by $\mathbf{q}\mapsto k\mathbf{q}-\sum V$; an isometry of $V$ is an orthogonal map fixing the origin, determined by the images of three linearly independent vertices, which are searched among the vertices with the same squared norm and mutual inner products; each candidate is accepted only if the induced bijection of $V$ preserves all pairwise integer squared distances. Congruence is decided exactly by testing for a bijection of hull-vertex sets preserving all integer squared distances (grouping by the sorted distance multiset, then an edge-labelled graph isomorphism), with a floating-point Kabsch superposition as a cross-check. For the Siamese proof (Proposition 5.2, Section B) the finitely many polynomial identities and inequalities in $m$ are verified by exact integer polynomial arithmetic, an inequality $p(m)>0$ for all integers $m\geq 2$ being certified by positivity of the coefficients of $p(2+t)$ or, failing that, by exact evaluation up to a Cauchy root bound.

**Reproducibility.** The scripts for every table and figure and the JSON files containing all reported numbers are provided as ancillary files in the directory `anc/`, with a `README` listing the commands that regenerate them. Squares are numbered $0,\dots,879$ in the order of `anc/squares880.txt`. The web application at `https://kylemath.github.io/MagicGemWebpage/` displays the 880 essentially different order-four squares and renders the gem of a selected one.

# B Siamese gems: the remaining steps of the proof

We continue the proof of Proposition 5.2 from the closed form $s(i,j)=n\langle i+j-m\rangle+\langle i+2j-2m\rangle+1$.

*Step 2: affine reduction to five layers.* In gem coordinates $x=j-m$, $y=m-i$, with $\langle t\rangle=t-n\lfloor t/n\rfloor$,

$$z=\underbrace{n(x-y+m)+(2x-y+m)+1-\tfrac{n^2+1}{2}}_{L(x,y)\ \text{affine}}+n\,\zeta(x,y),\quad \zeta:=-n\left\lfloor\tfrac{x-y+m}{n}\right\rfloor-\left\lfloor\tfrac{2x-y+m}{n}\right\rfloor.$$

The map $\Phi(x,y,z)=(x,y,(z-L(x,y))/n)$ is an affine bijection, so $\Phi(\mathcal{G}(S))=\mathrm{conv}\{(x,y,\zeta)\}$ has the same face lattice and vertex cells as $\mathcal{G}(S)$. In the box $|x|,|y|\leq m$ both floors lie in $\{-1,0,1\}$, the pairs $(-1,1),(-1,0)$ and, by the central symmetry $\zeta(-x,-y)=-\zeta(x,y)$, $(1,-1),(1,0)$ are impossible, so $\zeta$ takes the five values $\pm(n+1),\pm1,0$ on the lattice points of five convex lattice polygons (Figure 9a): $T=\mathrm{conv}\{(-m,m),(-1,m),(-m,1)\}$ at $\zeta=n+1$ (cut out by $y\geq x+m+1$), $U=\mathrm{conv}\{(-m,1-m),(-m,0),(-1,m-1)\}$ at $\zeta=1$ (by $2x+m+1\leq y\leq x+m$), $M=\mathrm{conv}\{(-m,-m),(0,m),(m,m),(0,-m)\}$ at $\zeta=0$ (by $|y-2x|\leq m$), and $D=-U$, $B=-T$ at $\zeta=-1,-(n+1)$.

*Step 3: the twelve points and sixteen planes.* Let $v_1=(-m,m,n+1)$, $v_2=(-1,m,n+1)$, $v_3=(-m,1,n+1)$, $v_4=(-m,0,1)$, $v_5=(-m,-m,0)$, $v_6=(0,-m,0)$, $v_7=(0,m,0)$, $v_8=(m,m,0)$,

$v_9 = (m, 0, -1)$, $v_{10} = -v_1$, $v_{11} = -v_3$, $v_{12} = -v_2$ (the cells listed in the statement), and consider the inequalities

$$\begin{aligned}
&(1)\ \zeta \le 2m+2, \qquad (2)\ y \le m, \qquad (3)\ x \ge -m, \qquad (8)\ -2y+\zeta \le 2m,\\
&(4)\ -2(m{+}1)x + (2m{+}1)y - m\zeta \le m(2m{+}1),\\
&(5)\ 2(m{+}1)x - 2(m{+}1)y + (2m{+}1)\zeta \le 2m(m{+}1),\\
&(6)\ 2m(m{+}2)x - (2m^2{+}2m{-}1)y + m(2m{+}1)\zeta \le m(2m^2{+}2m{-}1),\\
&(7)\ 2mx - y + m\zeta \le m(2m{-}1),
\end{aligned}$$

together with their images $(1')$–$(8')$ under $(x, y, \zeta) \mapsto -(x, y, \zeta)$; let $Q$ be the polyhedron they define. The script `siamese_proof.py` verifies, as polynomial identities and inequalities in $m$ valid for all integers $m \ge 2$ (Section A): (a) every vertex of $T, U, M, D, B$ at its height satisfies all 16 inequalities, so $\Phi(\mathcal{G}(S)) \subseteq Q$; (b) each plane contains exactly the following $v$'s, all others lying strictly inside:

$$\begin{aligned}
&(1)\colon v_1v_2v_3, \quad (2)\colon v_1v_2v_8v_7, \quad (3)\colon v_1v_3v_5v_4, \quad (4)\colon v_1v_4v_7,\\
&(5)\colon v_2v_3v_6, \quad (6)\colon v_2v_6v_9, \quad (7)\colon v_2v_8v_9, \quad (8)\colon v_3v_5v_6,
\end{aligned}$$

with centrally symmetric lists for $(1')$–$(8')$; (c) the quadrilaterals are in convex position in the cyclic order given, so the 16 face polygons have 26 distinct edges, each shared by exactly two polygons; (d) each $v_k$ lies on three planes with independent normals. By (b) each plane supports $P := \mathrm{conv}\{v_1, \ldots, v_{12}\}$ in a facet with the stated vertices; by (c) this family of facets is closed under edge-adjacency, and since the facet-adjacency graph of a 3-polytope is connected these are all the facets, so $P = Q$. With (a), $P \subseteq \Phi(\mathcal{G}(S)) \subseteq Q = P$, whose vertex set is $\{v_1, \ldots, v_{12}\}$ by (d); mapping back by $\Phi^{-1}$ gives the statement, with $(V, E, F) = (12, 26, 16)$ read off from (b)–(c). For $m = 1$ the triangles degenerate to points and the same argument gives the 8-vertex Lo Shu gem.