

Lectures on Stabilized Ellipsoid Embeddings

KYLER SIEGEL*
LES MARÉCOTTES, SWITZERLAND
September 2025

Abstract

These notes are based on a five-part minicourse on stabilized symplectic embeddings given at a workshop in Les Marécottes, Switzerland during a September 2025 workshop.¹ Our main goal is to explain the recent resolution of the (restricted) stabilized ellipsoid embedding problem by D. McDuff and the author. Along the way we also introduce various other ideas which shed light on the context and hint at possible generalizations. Some of the concepts covered include sesquicuspidal curves, symplectic inflation, multidirectional tangency constraints, well-placed curves, cluster transformations, Looijenga pairs, toric models, scattering diagrams, and the tropical vertex theorem.

Contents

| | |
|---|-----------|
| Lecture 1: Overview | 2 |
| §1.1 Hamiltonian flows and ellipsoid embeddings | 2 |
| §1.2 The symplectic Fibonacci staircase | 4 |
| §1.3 Full fillings by long ellipsoids | 5 |
| §1.4 Stabilized ellipsoid embeddings | 7 |
| §1.5 Unicuspidal and sesquicuspidal curves | 8 |
| §1.6 Outline of the remaining lectures | 10 |
| Lecture 2: Enumerative geometry of sesquicuspidal curves | 10 |
| §2.1 The inner and outer corner curves | 10 |
| §2.2 Inflation along sesquicuspidal curves | 11 |
| §2.3 Multidirectional tangency constraints | 14 |
| Lecture 3: Well-placed curves and their symmetries | 18 |
| §3.1 Looijenga pairs | 18 |
| §3.2 Toric models | 18 |
| §3.3 Elementary transformations | 18 |
| §3.4 Well-placed curves | 18 |
| §3.5 The shift and reflection symmetries | 18 |
| Lecture 4: Scattering diagrams and the tropical vertex | 18 |
| §4.1 Scattering diagram basics | 18 |
| §4.2 Some toy examples | 18 |
| §4.3 The tropical vertex theorem | 18 |
| Lecture 5: Putting it all together | 18 |

*K.S. is partially supported by NSF grant DMS-2506639 and a Simon Foundation Travel Support for Mathematicians Award.

¹Special thanks to Felix Schlenk and Grisha Mikhalkin for organizing this workshop.

| | |
|--|----|
| §5.1 The change of lattice trick | 18 |
| §5.2 Proof of Theorem C | 18 |

Disclaimer. Many of the mathematical statements in these lectures are somewhat informal simplifications; we recommend consulting the original references for more precise formulations whenever possible. Also, we apologize in advance for any missing citations, and we welcome any comments, questions, or corrections.

Lecture 1: Overview

In this lecture, we first briefly introduce Hamiltonian flows, nonsqueezing phenomena, and the ellipsoid embedding problem in §1.1. We then recall the celebrated McDuff–Schlenk Fibonacci staircase in §1.2, and give a brief modern overview of ways to derive it. In §1.3 we recall some connections between four-dimensional ellipsoid embeddings, symplectic ball packings, and symplectic cones, and as an application we establish full symplectic fillings of the four-ball by sufficiently long ellipsoids. In §1.4, we introduce stabilized ellipsoid embeddings and formulate our main result, namely Theorem A. Lastly, in §1.5 we introduce sesquicuspidal curves as a key tool for obstructing stabilized ellipsoid embeddings. We end with a brief sketch in §1.6 of the remaining lectures.

§1.1 Hamiltonian flows and ellipsoid embeddings

Consider a Hamiltonian on \mathbb{R}^{2n} , i.e. a smooth function $H : \mathbb{R}^{2n} \rightarrow \mathbb{R}$. Its *symplectic gradient* (a.k.a. *Hamiltonian vector field*) is defined by $\nabla_\omega H := J\nabla H$, where ∇ is the usual gradient and J is the block matrix $J = \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}$. Integrating $\nabla_\omega H$ gives rise to the flow $\text{fl}_H^t : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ for $t \in \mathbb{R}$.

There are various interesting questions one can ask about Hamiltonian flows, for instance:

- periodic orbits (existence, lower bounds, density, . . .)
- integrability versus chaos (exact solvability, KAM tori)
- Poincaré recurrence (or “superrecurrence”)

and so on. These lectures are related to the so-called “symplectic nonsqueezing phenomena” and its generalizations. We first recall some basics.

Proposition 1.1 (“Liouville’s theorem”)

The symplectic gradient $\nabla_\omega H$ is divergence-free, i.e. fl_H^t preserves volume.

Notation 1.2. For $\vec{a} = (a_1, \dots, a_n) \in \mathbb{R}_{>0}^n$, put $E(\vec{a}) := \left\{ \pi \sum_{i=1}^n \frac{1}{a_i} (x_i^2 + y_i^2) \leq 1 \right\}$.

Remark 1.3. Note that $E(a, \dots, a)$ is the ball $B^{2n}(a)$ of radius $\sqrt{a/\pi}$. Also, if $\mu : \mathbb{C}^n \rightarrow \mathbb{R}_{\geq 0}^n$ denotes the moment map for the standard \mathbb{T}^n -action, i.e. $\mu(z_1, \dots, z_n) = (\pi|z_1|^2, \dots, \pi|z_n|^2)$, then $\mu(E(a_1, a_2))$ is the right triangle with vertices $(0, 0)$, $(a_1, 0)$, $(0, a_2)$ as in Figure 1.1.

Notation 1.4. Put $E(\vec{a}) \xrightarrow{s} E(\vec{b})$ if $\text{fl}_H^1(E(\vec{a})) \subset E(\vec{b})$ for some (time-dependent) $H : \mathbb{R}^{2n} \times [0, 1] \rightarrow \mathbb{R}$.

Remark 1.5. This turns out to be equivalent to abstract symplectic embedding, i.e. a smooth embedding




Figure 1: The image of $E(a_1, a_2)$ under the moment map $\mu : \mathbb{C}^2 \rightarrow \mathbb{R}_{\geq 0}^2$.

preserving the standard symplectic form, thanks to the “extension after restriction principle” – see e.g. [Sch18, §4.4]. The discrepancy between time-dependent and time-independent Hamiltonian flows is more subtle – see e.g. [PS16].

Theorem 1.6 (Gromov ‘85 [Gro85])

If $E(\vec{a}) \xrightarrow{s} E(\vec{b})$, then we must have $\min\{a_i\} \leq \min\{b_i\}$.

Definition 1.7. For a symplectic manifold M^{2n} , the **ellipsoid embedding function** is

$$\mathcal{E}_M : \mathbb{R}_{>0}^n \rightarrow \mathbb{R}_{>0}, \quad \mathcal{E}_M(\vec{a}) := \inf\{c \mid E(\tfrac{1}{c} \cdot \vec{a}) \xrightarrow{s} M\}.$$

Remark 1.8. Note that we have $\mathcal{E}_M(c \cdot \vec{a}) = c\mathcal{E}_M(\vec{a})$ for any scalar $c \in \mathbb{R}_{>0}$, i.e. $\mathcal{E}_M(\vec{a})$ is really a function of just $n - 1$ variables.

Problem 1.9 (“ellipsoid embedding problem” (EEP)). Compute $\mathcal{E}_M(\vec{a})$ for $M = E(\vec{b})$.

Problem 1.10 (“restricted ellipsoid embedding problem” (REEP)). Compute $\mathcal{E}_M(\vec{a})$ for $M = B^{2n}$.

Here are a few initial comments.

- The ellipsoid embedding problem is trivial for $n = 1$.
- The ellipsoid embedding problem is “solved” for $n = 2$ by the work of many people (McDuff, Hutchings, Schlenk, Hofer, Biran, Polterovich, Li, ...). Here “solved” means reduced to a deceptively simple combinatorial criterion.
- The solution to the restricted ellipsoid embedding problem for $n = 2$ was worked out in complete detail by McDuff–Schlenk [MS12].
- To our knowledge there is not even a conjecture for the restricted ellipsoid embedding problem for $n \geq 3$ (i.e. a formula for the function $\mathcal{E}_{B^6}(a_1, a_2, a_3)$).

§1.2 The symplectic Fibonacci staircase

Theorem 1.11 (McDuff–Schlenk [MS12])

The ellipsoid embedding function $\mathcal{E}_{B^4}(1, a)$ of the four-dimensional ball B^4 is as follows:




Figure 2: The ellipsoid embedding function $\mathcal{E}_{B^4}(1, a)$.

While Figure 2 is just a cartoon, the function $\mathcal{E}_{B^4}(1, a)$ is explicitly computed in [MS12] and it includes the following noteworthy features:

- it is an “infinite Fibonacci staircase” for $a \in [1, \tau^4]$, with infinitely many steps accumulating at $\tau^4 := \frac{1}{2}(7 + 3\sqrt{5}) \approx 6.85$, where the x and y coordinates of the outer and inner corners are ratios of Fibonacci numbers (or their squares)
- for $a \in [8\frac{1}{36}, \infty)$, we have $\mathcal{E}_{B^4}(1, a) = \sqrt{a}$, i.e. volume is the only embedding obstruction.
- it is the linear function $\frac{1}{3}(a + 1)$ on the segment $a \in [\tau^4, 7]$, and in the range $a \in [7, 8\frac{1}{36}]$ it exhibits transitional behavior.

Let Fib_i denote the i th Fibonacci number:

| | | | | | | | | | |
|----------------|---|---|---|---|---|---|---|----|-----|
| i | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
| Fib_i | 0 | 1 | 1 | 2 | 3 | 5 | 8 | 13 | ... |

Exercise 1.12. Show that $\mathcal{E}_{B^4}(1, a)$ is nondecreasing and $\frac{1}{a} \cdot \mathcal{E}_{B^4}(1, a)$ is nonincreasing. Conclude that $\mathcal{E}_{B^4}(1, a)$ for $a \in [1, \tau^4]$ is entirely characterized by:

- embedding obstructions:

$$\mathcal{E}_{B^4}\left(1, \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+1}}\right) \geq \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+3}} \quad (1.1)$$

- embedding constructions:

$$\mathcal{E}_{B^4} \left(1, \frac{\text{Fib}_{2k+3}^2}{\text{Fib}_{2k+1}^2} \right) \leq \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+1}} \quad (1.2)$$

There are multiple ways of establishing (1.1) and (1.2). For (1.1), all of the main approaches are based on **pseudoholomorphic curves**.² While the method of extracting symplectic embedding obstructions from pseudoholomorphic curves is by now fairly streamlined, there are many different approaches for *producing* such curves, for instance:

- applying Cremona transformations to basic exceptional curves
- embedded contact homology
- cluster symmetries.

In these lectures we will primarily explain the last bullet above, as it is the most directly connected to our primary goal.

For (1.2), there are several methods for constructing symplectic embeddings, most notably:

- inflation along pseudoholomorphic curves
- almost toric fibrations.

While almost toric fibrations (ATFs) provide a very visually and conceptually appealing approach to the embeddings for (1.2), we will primarily focus on the first bullet above in these lectures, partly because the ATF approach to embeddings is already discussed very thoroughly elsewhere (see e.g. [Sym, Eva23, CGHMP25, CV22]) and partly because it allows us to reduce both (1.1) and (1.2) to the existence of certain singular symplectic curves in the complex projective plane. There are then multiple methods for constructing the curves relevant for inflation approach to (1.2): Seiberg–Witten theory, tropical methods (see e.g. [MS25b, §5.3]), etc.

§1.3 Full fillings by long ellipsoids

We associate to each positive rational number $a \in \mathbb{Q}_{>1}$ its **weight sequence** $\text{wt}(a) = (w_1, \dots, w_k)$, whose definition is more or less explained by the “box diagram” in Figure 3, which divides a rectangle of height 1 and width a into a collection of squares.

Example 1.13. For $a = 17/5$, the corresponding weight sequence is $\text{wt}(17/5) = (1, 1, 1, 2/5, 2/5, 1/5, 1/5)$.


$\underbrace{\hspace{1.5cm}}_3 \quad \underbrace{\hspace{1.5cm}}_2 \quad \underbrace{\hspace{1.5cm}}_2$

Note that:

- $\frac{17}{5} = 3 + \frac{1}{2 + \frac{1}{2}}$, i.e. the multiplicities of the entries in the weight sequence give the continued fraction expansion
- $1 \geq w_1 \geq w_2 \geq \dots \geq w_k$
- $w_1^2 + \dots + w_k^2 = a$.

Each of these properties holds for general $a \in \mathbb{Q}_{>1}$.

²It is, however, interesting to ask how much of this story could be recovered using generating functions or microlocal sheaves.

Figure 3: The box diagram giving the weight sequence for $a = 17/5$.**Theorem 1.14 (McDuff [McD09])**

For each $a \in \mathbb{Q}_{>1}$, say with weight sequence $\text{wt}(a) = (w_1, \dots, w_k)$, we have

$$E(1, a) \xrightarrow{s} B^4(c) \iff \bigsqcup_{i=1}^k B^4(w_i) \xrightarrow{s} B^4(c).$$

Notation 1.15. Put:

- $\text{Bl}^k \mathbb{CP}^2 := \mathbb{CP}^2 \# (\underbrace{\mathbb{CP}^2 \# \dots \# \mathbb{CP}^2}_k)$, i.e. the k -fold blowup of \mathbb{CP}^2 as a smooth oriented manifold
- $H_2(\text{Bl}^k \mathbb{CP}^2) = \mathbb{Z}\langle \ell, e_1, \dots, e_k \rangle$, with ℓ the line class and e_1, \dots, e_k the obvious exceptional classes
- $C(\text{Bl}^k \mathbb{CP}^2) := \left\{ [\omega] \in H^2(\text{Bl}^k \mathbb{CP}^2) \mid \begin{array}{l} \omega \text{ symplectic form on } \text{Bl}^k \mathbb{CP}^2, \\ c_1(\omega) = \text{PD}(3\ell - e_1 - \dots - e_k) \end{array} \right\}$, the “symplectic cone with standard first Chern class”.

Theorem 1.16 (McDuff–Polterovich [MP94])

We have

$$\bigsqcup_{i=1}^k B^4(w_i) \xrightarrow{s} B^4(c) \iff \text{PD}(c\ell - w_1 e_1 - \dots - w_k e_k) \in C(\text{Bl}^k \mathbb{CP}^2).$$

Notation 1.17. Let $\text{Exc}(\text{Bl}^k \mathbb{CP}^2)$ denote the set of all homology classes $e \in H_2(\text{Bl}^k \mathbb{CP}^2)$ such that e is represented by a symplectically embedded sphere with self-intersection number -1 .

Remark 1.18. Note that a priori the definition of $\text{Exc}(\text{Bl}^k \mathbb{CP}^2)$ depends on the symplectic form on $\text{Bl}^k \mathbb{CP}^2$, but in fact one can show that it is independent of the choice of symplectic form in $C(\text{Bl}^k \mathbb{CP}^2)$ (this uses a Gromov compactness argument and the fact that any two $\omega, \omega' \in C(\text{Bl}^k \mathbb{CP}^2)$ are homotopic through symplectic forms).

Remark 1.19. Note that for $e = d\ell - m_1 e_1 - \dots - m_k e_k \in \text{Exc}(\text{Bl}^k \mathbb{CP}^2)$, we have

$$e \cdot e = d^2 - m_1^2 - \dots - m_k^2 = -1 \quad \text{and} \quad c_1(e) = 3d - m_1 - \dots - m_k = 1. \quad (1.3)$$

Theorem 1.20 ([McD98, Bir97, LL01, LL02])

We have

$$C(\mathrm{Bl}^k \mathbb{C}\mathbb{P}^2) = \left\{ \alpha \in H^2(\mathrm{Bl}^k \mathbb{C}\mathbb{P}^2) \mid \begin{array}{l} \alpha \cdot \alpha > 0, \\ \langle \alpha, e \rangle > 0 \ \forall e \in \mathrm{Exc}(\mathrm{Bl}^k \mathbb{C}\mathbb{P}^2) \end{array} \right\}.$$

Corollary 1.21. For $a \in \mathbb{Q}_{>1}$, say with $\mathrm{wt}(a) = (w_1, \dots, w_k)$, we have

$$\mathcal{E}_{B^4}(1, a) = \max \left(\sqrt{a}, \sup \left\{ \frac{m_1 w_1 + \dots + m_k w_k}{d} \mid d\ell - m_1 e_1 - \dots - m_k e_k \in \mathrm{Exc}(\mathrm{Bl}^k \mathbb{C}\mathbb{P}^2) \right\} \right)$$

Lemma 1.22. We have $\mathcal{E}_{B^4}(1, a) = \sqrt{a}$ for all $a \geq 9$.

Proof. It suffices to establish $\frac{m_1 w_1 + \dots + m_k w_k}{d} \leq \sqrt{a}$ for all $d\ell - m_1 e_1 - \dots - m_k e_k \in \mathrm{Exc}(\mathrm{Bl}^k \mathbb{C}\mathbb{P}^2)$. Observe that we have

$$\frac{m_1 w_1 + \dots + m_k w_k}{d} \leq \frac{m_1 + \dots + m_k}{d} = \frac{3d - 1}{d} < 3 < \sqrt{a}.$$

□

§1.4 Stabilized ellipsoid embeddings

Definition 1.23. For a symplectic manifold M^{2n} and $N \in \mathbb{Z}_{\geq 1}$, the **stabilized ellipsoid embedding function** is

$$\mathcal{E}_M^N : \mathbb{R}_{>0}^n \rightarrow \mathbb{R}_{>0}, \quad \mathcal{E}_M^N(\vec{a}) := \inf \{ c \mid E(\tfrac{1}{c} \cdot \vec{a}) \times \mathbb{R}^{2N} \xhookrightarrow{s} M \times \mathbb{R}^{2N} \}. \quad (1.4)$$

Remark 1.24. Note that by taking the identity map in the extra dimensions, we always have

$$\mathcal{E}_M^N(\vec{a}) \leq \mathcal{E}_M(\vec{a}).$$

Problem 1.25 (“stabilized ellipsoid embedding problem” (SEEP)). Compute $\mathcal{E}_M^N(\vec{a})$ for $M = E(\vec{b})$ and $N \in \mathbb{Z}_{\geq 1}$.

Problem 1.26 (“restricted stabilized ellipsoid embedding problem” (RSEEP)). Compute $\mathcal{E}_M^N(\vec{a})$ for $M = B^{2n}$ and $N \in \mathbb{Z}_{\geq 1}$.

Remark 1.27. Note that we can think of $E(a_1, \dots, a_n) \times \mathbb{R}^{2N}$ as $E(a_1, \dots, a_n, \underbrace{\infty, \dots, \infty}_N)$, and hence

the stabilized ellipsoid embedding problem is essentially a special case of the corresponding unstabilized ellipsoid embedding problem (but now in higher dimensions).

Our primary goal in these lectures is to explain the proof of the following theorem.

Theorem A ([MS24a])

For all $N \in \mathbb{Z}_{\geq 1}$, the stabilized ellipsoid embedding of the four-dimensional ball is given by

$$\mathcal{E}_{B^4}^N(1, a) = \begin{cases} \mathcal{E}_{B^4}(1, a) & a \in [1, \tau^4] \\ \frac{3a}{a+1} & a > \tau^4 \end{cases}$$

See Figure 4 for an illustration.




Figure 4: The stabilized ellipsoid embedding function $\mathcal{E}_{B^4}^N(1, a)$.

In brief, the function $\mathcal{E}_{B^4}^N(1, a)$ has a “subcritical regime” which is an infinite staircase and a “supercritical regime” which is a smooth rational function, with the phase transition at $a = \tau^4$. One can check that the rational function $\frac{3a}{a+1}$ meets the infinite Fibonacci staircase precisely at its outer corners and lies otherwise above it for $a \in [1, \tau^4]$, whereas it lies below \sqrt{a} for $a > \tau^4$. Note that the volume bound \sqrt{a} evidently no longer holds for the stabilized problem, i.e. the supercritical behavior is a truly higher dimensional phenomenon.

Remark 1.28. There were important earlier partial results by subsets of Hind, Kerman, Cristofaro-Gardiner, McDuff, and the author (see e.g. [HK14, CGH18, CGHM18, McD18, Sie22]). In particular, the function for $a \in [1, \tau^4]$ was worked out in [CGH18], which by Exercise 1.12 and (1.4) boils down to showing that the inequality (1.1) persists after stabilization.

The upper bounds for Theorem A are given by an explicit “symplectic folding” construction due to Hind, which adapts Guth’s polydisk embedding construction in [Gut08].

Theorem 1.29 (Hind [Hin15])

For each $a \in \mathbb{R}_{\geq 1}$ and $N \in \mathbb{Z}_{\geq 1}$ we have

$$E(1, a) \times \mathbb{R}^{2N} \xhookrightarrow{s} B^4(c) \times \mathbb{R}^{2N} \quad \text{for} \quad c > \frac{3a}{a+1}. \quad (1.5)$$

Thus, our main goal is to find optimal lower bounds for $\mathcal{E}_{B^4}^N(1, a)$, especially for $a > \tau^4$.

§1.5 Unicuspidal and sesquicuspidal curves

Let p, q be positive integers satisfying $\gcd(p, q) = 1$ and $p > q \geq 2$.

Definition 1.30. A (p, q) -sesquicuspidal symplectic curve in a symplectic four-manifold M^4 is

a subset $C \subset M$ such that

- C has one (p, q) cusp singularity (i.e. modeled locally on $\{x^p + y^q = 0\} \subset \mathbb{C}^2$)
- C has finitely many additional positive ordinary double points (i.e. modeled locally on $\{xy = 0\} \subset \mathbb{C}^2$)
- C is otherwise symplectically embedded.

Remark 1.31. The second condition in Definition 1.30 holds after a generic perturbation, and the number of double points $\underline{\delta}$ away from the distinguished cusp is fixed by adjunction, namely

$$\underline{\delta} = \frac{1}{2}(C \cdot C + \chi(C) - c_1(C) - (p-1)(q-1)). \quad (1.6)$$

In the case $\underline{\delta} = 0$, the curve is called **unicuspidal**.³ Note that we can similarly define (p, q) -sesquicuspidal algebraic curves, which in particular give examples of (p, q) -sesquicuspidal symplectic curves.

Definition 1.32. We will define the (real) **index** of a (p, q) -sesquicuspidal curve $C \subset M$ to be $\text{ind}(C) := 2(c_1(C) - p - q)$.

We will see later that this is the expected (real) dimension of the moduli space of (p, q) -sesquicuspidal curves in a fixed homology class and with fixed “position” of the distinguished cusp.

Proposition B ([MS23, MS25a])

Given a rational index zero (p, q) -sesquicuspidal symplectic curve C in a closed symplectic four-manifold M^4 , we have

$$\mathcal{E}_M^N(p, q) \geq \frac{pq}{\text{area}(C)} \quad (1.7)$$

for all $N \in \mathbb{Z}_{\geq 0}$.

Here $\text{area}(C)$ denotes the symplectic area of C , which depends only on its homology class $[C] \in H_2(M)$. Note that, after rescaling, (1.7) can be written equivalently as $\mathcal{E}_M^N(1, p/q) \geq \frac{p}{\text{area}(C)}$.

The above proposition allows us to reduce Theorem A to the following key construction of singular algebraic curves. As above we assume $\gcd(p, q) = 1$ and $p > q \geq 2$.

Theorem C ([MS24a])

There exists a rational index zero (p, q) -sesquicuspidal symplectic curve $C \subset \mathbb{CP}^2$ if and only if

- (a) $p/q = \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+1}}$ for $k \in \mathbb{Z}_{\geq 1}$, or
- (b) $p/q > \tau^4$.

Exercise 1.33. Combining Proposition B with Theorem C gives precisely the lower bounds on $\mathcal{E}_{B^4}^N$ needed for Theorem A.

Thus in the remaining lectures we will focus on the proofs of Proposition B and Theorem C, albeit with various detours.

³The term sesquicuspidal comes from Latin sesqui meaning “one and a half”, reflecting the fact that sesquicuspidal curves are only mildly more singular than unicuspidal curves.

§1.6 Outline of the remaining lectures

Here is the plan for the remaining lectures:

- In Lecture 2, we discuss the enumerative geometry of sesquicuspidal curves, and we use this formalism to sketch a proof of Proposition B.
- In Lecture 3, we define well-placed curves and discuss their cluster symmetries, and we give one construction of the “outer corner curves” alluded to in §1.2.
- In Lecture 4, we give a brief crash course in scattering diagrams and formulate the tropical vertex theorem proved in [GPS10].
- Finally, in Lecture 5 we put all of this together to give the proof of Theorem C.

Remark 1.34. While much of the theory developed in this lectures can be extended in various directions, for instance by replacing \mathbb{CP}^2 with other del Pezzo surfaces, for simplicity we will mostly focus our exposition on the case of the complex projective plane.

Lecture 2: Enumerative geometry of sesquicuspidal curves

§2.1 The inner and outer corner curves

Let us begin by recalling the following result as motivation.

Theorem 2.1 ([FdBLMHN06])

There exists a (p, q) -unicuspidal rational algebraic curve in \mathbb{CP}^2 of degree d with $p > q \geq 2$ and $\gcd(p, q) = 1$ if and only if (d, p, q) is one of the following:

- (a) $(p, q) = (d, d - 1)$ for $d \geq 3$
- (b) $(p, q) = (2d - 1, d/2)$ for $d \geq 4$ even
- (c) $(p, q) = (\text{Fib}_{2k+3}^2, \text{Fib}_{2k+1}^2)$ and $d = \text{Fib}_{2k+3}\text{Fib}_{2k+1}$ for $k \geq 1$
- (d) $(p, q) = (\text{Fib}_{2k+5}, \text{Fib}_{2k+1})$ and $d = \text{Fib}_{2k+3}$ for $k \geq 1$
- (e) $(p, q) = (22, 3)$ and $d = 8$
- (f) $(p, q) = (43, 6)$ and $d = 16$.

Recall that the index of such a curve C is given by

$$\text{ind}(C) = 2(c_1(C) - p - q) = 2(3d - p - q).$$

Exercise 2.2. The indices of the curves in Theorem 2.1 are as follows:

| | (a) | (b) | (c) | (d) | (e) | (f) |
|-------|----------|---------|-----|-----|-----|-----|
| index | $2d + 2$ | $d + 2$ | 2 | 0 | -2 | -2 |

Observe that, in (c), $\frac{\text{Fib}_{2k+3}^2}{\text{Fib}_{2k+1}^2}$ is precisely the x -value of the k th inner corner point in the Fibonacci staircase from Theorem 1.11, so we will refer to these as **inner corner curves**. Similarly, in (d), $\frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+1}}$ is precisely the x -value of the k th outer corner point, and we will refer to these as **outer**

corner curves. Note also that the outer corner curves have index zero, so by Proposition B their existence gives

$$\mathcal{E}_{B^4}^N \left(1, \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+1}} \right) \geq \frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+3}},$$

where $\frac{\text{Fib}_{2k+5}}{\text{Fib}_{2k+3}}$ is precisely the y -value of the k th outer corner point.

Remark 2.3. Here are a few more remarks about the curve families (a),(b),(c),(d),(e),(f) in Theorem 2.1:

- family (a) is given by $\{ZY^{d-1} = X^d\}$
- family (b) is given by $\{(ZY - X^2)^{d/2} = XY^{d-1}\}$
- Miyanishi–Sugie [MS81] or Kashiwara [Kas87] construct a pencil with general member (c) and special member (d)
- Orevkov [Ore02] constructs families (d),(e),(f) (we will explain a related construction in §3.5 below)
- families (e),(f) are a bit “surprising” to a symplectic geometer in the sense that they should not persist under a generic perturbation of the almost complex structure.

§2.2 Inflation along sesquicuspidal curves

The following proposition, which is essentially an application of the technique of **symplectic inflation**, constructs four-dimensional symplectic ellipsoid embeddings from singular curves. While it is not directly related to the proof of Theorem C, it does help illustrate the role connection between sesquicuspidal curves and (stabilized) symplectic embeddings.

Proposition D

Let M be a closed symplectic four-manifold and $C \subset M$ a (p, q) -sesquicuspidal symplectic curve such that $[C] = c\text{PD}([\omega_M])$ for some $c \in \mathbb{R}_{>0}$ and $[C] \cdot [C] \geq pq$. Then $\mathcal{E}_M(p, q) \leq c$ (i.e. $\mathcal{E}_M(1, p/q) \leq c/q$).

Remark 2.4. Note that we have

$$\text{vol}(M) = \frac{1}{2} \int \omega_M \wedge \omega_M = \frac{1}{2} \text{PD}([\omega_M]) \cdot \text{PD}([\omega_M]) = \frac{1}{2c^2} [C] \cdot [C].$$

In the case $[C] \cdot [C] = pq$, Proposition D corresponds to an embedding $E\left(\frac{p}{c}, \frac{q}{c}\right) \xrightarrow{s} M$ (up to small error), which is a full filling since $\text{vol}(E\left(\frac{p}{c}, \frac{q}{c}\right)) = \frac{pq}{2c^2} = \text{vol}(M)$.

Exercise 2.5. Show that the inner corner curves (i.e. family (c) from Theorem 2.1) satisfy $[C] \cdot [C] = d^2 = pq$, and these translate into full fillings via Proposition D.

Combining Exercise 1.33 and Exercise 2.5, we see that the entire (stabilized) Fibonacci staircase $\mathcal{E}_{B^4}^N(1, a)$ for $a \in [1, \tau^4]$ and $N \in \mathbb{Z}_{\geq 0}$ follows from the existence of the outer and inner corner unicuspidal curves (i.e. families (c) and (d) in Theorem 2.1).

Although Proposition D will suffice for our purposes here, let us also mention that symplectic inflation can also be performed along curves with more complicated singularities, for example a **k -fold (p, q) -multicusp**, i.e. the singularity locally modeled on

$$\left\{ \prod_{j=1}^k (x^p - C_j y^p) = 0 \right\} \subset \mathbb{C}^2$$

for some (pairwise distinct) constants $C_j \in \mathbb{C}$. In fact, the following theorem shows that the existence of such a curve is *equivalent* to the existence of a corresponding ellipsoid embedding.

Theorem 2.6 (Ophstein [Ops15])

Let M^4 be a symplectic four-manifold with rational symplectic form class $[\omega_M] \in H^2(M; \mathbb{Q})$, and fix $\tau \in \mathbb{Q}_{>0}$ and relatively prime $p, q \in \mathbb{Z}_{\geq 1}$. Then we have $E(\tau p, \tau q) \xrightarrow{s} M$ if and only if there exists an irreducible symplectic curve $C \subset M$ which has a $k\tau$ -fold (p, q) -multicusp and satisfies $[C] = k\text{PD}([\omega_M])$ and $[C] \cdot [C] \geq k\tau pq$, for some $k \in \mathbb{Z}_{\geq 1}$.

We end this subsection with a proof sketch of Proposition D, which introduces the method of symplectic inflation and also some useful pictures around resolutions of singularities.

Proof sketch of Proposition D.

Step 1: We first resolve the cusp singularity of C by a (p, q) -weighted symplectic blowup as in Figure 5, giving a new curve $\tilde{C} \subset \tilde{M}$. In more detail, this means that we start with a neighborhood of the cusp point p which is symplectomorphic to a neighborhood of the form $\mu^{-1}(U) \subset \mathbb{R}^4$, where U is a neighborhood of the origin on $\mathbb{R}_{\geq 0}^2$ and $\mu : \mathbb{R}^4 \rightarrow \mathbb{R}_{\geq 0}^2$ is the standard moment map. We may assume that this symplectomorphism identifies C locally near p with the pre-image under μ of a ray in $\mathbb{R}_{\geq 0}^2$ through the origin with direction (p, q) . We can assume that $\mu^{-1}(U)$ contains the ellipsoid $E(\varepsilon p, \varepsilon q)$ for some $\varepsilon > 0$ sufficiently small.

The weighted blowup $\tilde{M} := \text{Bl}_{p,q} M$ is then obtained by symplectically cutting out the ellipsoid $E(\varepsilon p, \varepsilon q)$, i.e. replacing the local toric picture with 2 edges with one with 3 edges as in the right side of Figure 5. Here $\tilde{C} \subset \tilde{M}$ is the strict transform of C , given by suitably truncating the line segment $\mu(C)$. Note that \tilde{C} no longer has a (p, q) cusp but may still have some additional double points, while \tilde{M} has (at most) two orbifold points, due to the two non-Delzant corners in the local toric picture.




Figure 5: The local toric picture for the (p, q) -weighted symplectic blowup $M \rightsquigarrow \tilde{M}$.

Step 2 (optional): Although the orbifold points are not necessarily problematic, it is sometimes preferable to avoid them, which we can do by further resolving \tilde{M} . In terms of the local toric picture, this corresponds to adding many small edges so that all of the vertices become Delzant – see Figure 6. We denote the resulting nonsingular symplectic manifold by $\tilde{\tilde{M}}$ and the new (essentially unaffected) curve by $\tilde{\tilde{C}} \subset \tilde{\tilde{M}}$.




Figure 6: The local toric picture for the resolution of orbifold singularities $\tilde{M} \rightsquigarrow \tilde{\tilde{M}}$.

Example 2.7. Figure 7 shows a concrete example of the process of going from \tilde{M} to $\tilde{\tilde{M}}$ by resolving the two orbifold points. In the local toric pictures, \tilde{M} has edges with outward normals $(-1, 0)$, $(-2, -3)$, $(0, -1)$, while $\tilde{\tilde{M}}$ has two new edges with outward normals $(-1, -1)$ and $(-1, -2)$. One can easily check that all the corners for $\tilde{\tilde{M}}$ are Delzant.




Figure 7: The orbifold resolution process $\tilde{M} \rightsquigarrow \tilde{\tilde{M}}$ in the case $(p, q) = (3, 2)$.

Remark 2.8. Note that the process of going from $C \subset M$ to $\tilde{C} \subset \tilde{\tilde{M}}$ is a symplectic version of embedded resolution of singularities for a (p, q) cusp. The combinatorics of this resolution process are neatly encoded in the box diagram for (p, q) . For example, the box diagram for $(p, q) = (3, 2)$ is shown in Figure 8, with the 3 boxes corresponding to the 2 small added edges in Figure 7 – see [MS25a, §4.1] for more details.




Figure 8: The box diagram for $(p, q) = (3, 2)$.

□

Step 3: We now smooth the remaining double points of $\tilde{\tilde{C}}$. This is straightforward in the

symplectic category, and locally modeled on trading $\{xy = 0\}$ for $\{xy = \delta\}$ in \mathbb{C}^2 for $\delta > 0$ small. We denote the resulting curve by $\tilde{\tilde{C}}_{\text{sm}} \subset \tilde{\tilde{M}}$.

Step 4: We can find a closed two-form Ω on $\tilde{\tilde{M}}$ which is supported in a small neighborhood of $\tilde{\tilde{C}}_{\text{sm}}$ and satisfies $[\Omega] = \text{PD}(\tilde{\tilde{C}}_{\text{sm}})$. Here it is necessary that $\tilde{\tilde{C}}_{\text{sm}}$ has nonnegative self-intersection number, which comes from our assumption $[C] \cdot [C] \geq pq$. Note that finding Ω is straightforward in the special case that $\tilde{\tilde{C}}_{\text{sm}}$ has zero self-intersection number, since then it has a trivial symplectic normal bundle and we can take Ω to be simply a two-form pulled back from the normal direction.

Step 5: We now “inflate” $\tilde{\tilde{M}}$ by replacing its symplectic form $\omega_{\tilde{\tilde{M}}}$ with the new one $\omega_{\tilde{\tilde{M}}}^s := \omega_{\tilde{\tilde{M}}} + s\Omega$ for $s \in \mathbb{R}_{>0}$. Let E denote the symplectic divisor in $\tilde{\tilde{M}}$ corresponding to the edge in the local toric picture with outward normal direction $(-p, -q)$ as in Figure 9. Note that E has symplectic area approximately ε with respect to $\omega_{\tilde{\tilde{M}}}$, given by the affine length of the corresponding edge in the local toric picture. Then E has area approximately $\varepsilon + s$ with respect to the symplectic form $\omega_{\tilde{\tilde{M}}}^s$.




Figure 9: The edge $E \subset \tilde{\tilde{M}}$ along which we blow down.

Step 6: We now reverse the blowup process $M \rightsquigarrow \tilde{\tilde{M}}$ by blowing down E and the small edges introduced in Step 2, which gives back M but now with a symplectic form ω_M^s . This can be again understand as a straightforward local toric substitution, but observe that the symplectic manifold (M, ω_M^s) now naturally contains the ellipsoid $E([\varepsilon + s]p, [\varepsilon + s]q)$, and we have $[\omega_M^s] = [\omega_M] + s\text{PD}(C) = (1 + sc)[\omega_M^c]$ in $H^2(M; \mathbb{R})$. After rescaling by the factor $1 + sc$ and applying Moser’s stability theorem, we thus have a symplectic embedding

$$E\left(\frac{\varepsilon+s}{1+sc}p, \frac{\varepsilon+s}{1+sc}q\right) \xhookrightarrow{s} (M, \omega_M).$$

Taking $s \rightarrow \infty$, we get an embedding $E(\frac{p}{c'}, \frac{q}{c'}) \xhookrightarrow{s} (M, \omega_M)$ for c' arbitrarily close to c .

§2.3 Multidirectional tangency constraints

Let M^{2n} be a closed $2n$ -dimensional symplectic manifold, and $\vec{D} = (D_1, \dots, D_n)$ be a collection of nonsingular local symplectic divisors which have a symplectic normal crossing intersection at a point $p \in M$ (that is, they look locally like $\{z_1 \cdots z_n = 0\} \subset \mathbb{C}^n$ in symplectic Darboux coordinates).

Notation 2.9. Let $\mathcal{J}(M, \vec{D})$ denote the set of tame almost complex structures on M which are integrable near p and preserve J_1, \dots, J_n .

Put $z_0 := [0 : 1] \in \mathbb{CP}^1$.

Definition 2.10. Given an almost complex structure $J \in \mathcal{J}(M, \vec{D})$, a homology class $A \in H_2(M)$, and a tuple $\vec{m} = (m_1, \dots, m_n) \in \mathbb{Z}_{\geq 1}^n$, let $\mathcal{M}_{M,A}^J \llbracket C_{\vec{D}}^{\vec{m}} \mathfrak{p} \rrbracket$ denote the moduli space of J -holomorphic maps $u : \mathbb{CP}^1 \rightarrow M$ which satisfy $[u] = A$, $u(z_0) = \mathfrak{p}$, and such that u has contact order at least m_i with D_i at z_0 for $i = 1, \dots, n$, modulo biholomorphic reparametrizations of \mathbb{CP}^1 fixing z_0 .

Here the contact order constraints $\llbracket C_{\vec{D}}^{\vec{m}} \mathfrak{p} \rrbracket$ with D_1, \dots, D_n is called a **multidirectional tangency constraint**.

Remark 2.11.

- When $\vec{m} = (m, \underbrace{1, \dots, 1}_{n-1})$, this reduces to the m -fold **local tangency constraint** (with respect to $D := D_1$) studied by Cieliebak–Mohnke and others (see e.g. [CM05, CM18, Ton18, MS21]).
- If $u : \mathbb{CP}^1 \rightarrow M^4$ has contact order exactly p with D_1 and exactly q with D_2 , then it necessarily has a (p, q) cusp singularity at \mathfrak{p} (assuming no other branches of u meet \mathfrak{p}).

Theorem E

Fix a semipositive symplectic manifold M^{2n} and homology class $A \in H_2(M)$, and put $\vec{m} = (p, q, \underbrace{1, \dots, 1}_{n-2})$ for some coprime $p, q \in \mathbb{Z}_{\geq 1}$ satisfying $p + q = c_1(A)$.

- For generic $J \in \mathcal{J}(M, \vec{D})$, the moduli space $\mathcal{M}_{M,A}^J \llbracket C_{\vec{D}}^{\vec{m}} \mathfrak{p} \rrbracket$ is finite and regular, and the signed count $\#_{M,A} \llbracket C^{\vec{m}} \rrbracket := \# \mathcal{M}_{M,A}^J \llbracket C_{\vec{D}}^{\vec{m}} \mathfrak{p} \rrbracket$ is independent of \mathfrak{p}, \vec{D}, J .
- For any other closed symplectic manifold Q^{2N} such that $M^{2n} \times Q^{2N}$ is semipositive, we have

$$\#_{M \times Q, A \times [\text{pt}]} \llbracket C^{(\vec{m}, \vec{1})} \rrbracket = \#_{M,A} \llbracket C^{\vec{m}} \rrbracket,$$

where $(\vec{m}, \vec{1}) = (m_1, \dots, m_n, \underbrace{1, \dots, 1}_N)$.

- If $\#_{M,A} \llbracket C^{\vec{m}} \rrbracket \neq 0$, then we have

$$\mathcal{E}_M^N(p, q, a_3, \dots, a_n) \geq \frac{pq}{\text{area}(A)}$$

whenever $a_3, \dots, a_n > pq$.

- If $2n = 4$, then $\#_{M,A} \llbracket C^{\vec{m}} \rrbracket \in \mathbb{Z}_{\geq 0}$, and if there exists a rational (p, q) -sesquicuspidal symplectic curve $C \subset M$ with $[C] = A$, then $\#_{M,A} \llbracket C^{\vec{m}} \rrbracket > 0$.

Here semipositivity is a technical topological assumption which allows us to achieve generic transversality without virtual techniques.

Remark 2.12. There is also a version of Theorem E for more general $\vec{m} \in \mathbb{Z}_{\geq 1}^n$, but the combinatorial assumptions on \vec{m} and $c_1(A)$ are more complicated to state – see [MS25a].

Example 2.13. According to Remark 2.11(i), $\#_{\mathbb{CP}^2, d\ell}^{(3d-1, 1)}$ is the count of degree d rational plane curves satisfying an order $3d - 1$ local tangency constraint. These were computed in [MS21] as

follows:

| | | | | | |
|--|---|---|---|----|-----|
| d | 1 | 2 | 3 | 4 | 5 |
| $\#_{\mathbb{CP}^2, d\ell}^{(3d-1,1)}$ | 1 | 1 | 4 | 26 | 217 |

Note that this sequence appears as A364973 in the On-line Encyclopedia of Integer Sequences, where it is also seen to make important appearances in the mirror symmetry literature.

Example 2.14. The counts corresponding to the outer corner curves in \mathbb{CP}^2 are all equal to 1:

| | | | | | |
|--|---|---|---|---|---|
| d | 1 | 2 | 3 | 4 | 5 |
| $\#_{\mathbb{CP}^2, \text{Fib}_{2k+3}\ell}^{(\text{Fib}_{2k+5}, \text{Fib}_{2k+1})}$ | 1 | 1 | 1 | 1 | 1 |

The general philosophy with Examples 2.13 and 2.14 is that unicuspidal curves tend to be unique (after imposing a suitable multidirectional tangency constraint), whereas sesquicuspidal curve counts tend to grow rapidly.

We end this lecture by briefing giving some of the ideas underlying Theorem E.

Proof ideas for Theorem E.

- (i) The standard approach is to use Gromov's compactness theorem, but here we need extra care to rule out additional "bad degenerations". For instance, a curve with a $(11, 5)$ cusp could degenerate into two components, one with an $(8, 3)$ cusp and one with a $(3, 2)$ cusp, which meet at the singular point as in Figure 10.




Figure 10: Degeneration of a curve with a $(11, 5)$ cusp

The following is our key tool for ruling out such degenerations.

Lemma F ("hidden constraints" [MS25a, §3.2])

If $\langle C^{\vec{m}} \rangle$ degenerates into $\langle C^{\vec{m}_1} \rangle, \dots, \langle C^{\vec{m}_k} \rangle$ then we must have

$$\sum_{i=1}^k \min\{a_1 m_1^i, \dots, a_n m_n^i\} \geq \min\{a_1 m_1, \dots, a_n m_n\} \quad (2.1)$$

for any $\vec{a} = (a_1, \dots, a_n) \in \mathbb{R}_{>0}^n$. Here we put $\vec{m} = (m_1, \dots, m_n)$ and $\vec{m}_i = (m_1^i, \dots, m_n^i)$ for $i = 1, \dots, k$,

For example, taking $\vec{a} = (5, 11)$ rules out the hypothetical degeneration depicted in Figure 10.

The rough heuristic idea behind Lemma F is as follows. After surrounding the cusp by an ellipsoid $E := E(\varepsilon \cdot \vec{a})$ for $\varepsilon > 0$ sufficiently small and stretching the neck along ∂E , the resulting degeneration picture involves a rational punctured curve C in the symplectization

$\mathbb{R} \times \partial E$ with one negative asymptotic Reeb orbit γ and many positive asymptotic Reeb orbits $\gamma_1, \dots, \gamma_k$. An explicit dictionary between cusp singularities and Reeb orbits in ∂E is given in [MS23], and in particular nonnegativity of C gives the action inequality $\sum_{i=1}^k \mathcal{A}_{\partial E}(\gamma_i) \geq \mathcal{A}_{\partial E}(\gamma)$, which translates directly into (2.1).

Remark 2.15. *The proof of Lemma F relies on neck stretching and the SFT compactness theorem. It would be interesting to see if there is a more algebraic or classical argument.*

(ii) We argue using a product almost complex structure on $M \times Q$:

- using some elementary Fredholm theory, one argues that regular curves $C \subset M$ remain regular as curves in $M \times Q$
- using some basic intersection theory, one shows that all relevant curves in $M \times Q$ must be entirely contained in $M \times \{\text{pt}\}$.

Together, these essentially show that our curve counts are unaffected by the stabilization process.

(iii) We will assume $2n = 4$ for simplicity. Suppose that we have an embedding $E(\frac{p}{c}, \frac{q}{c}) \xrightarrow{s} M$. By assumption we have a curve C with a (p, q) cusp and $[C] = A$, which looks locally near the cusp like the left hand side of Figure 5 with respect to an ellipsoid embedding $E(\varepsilon p, \varepsilon q) \xrightarrow{s} M$. By Exercise 2.16, the portion of the curve lying over the triangle with vertices $(0, 0), (\varepsilon p, 0), (0, \varepsilon q)$ has symplectic area $\varepsilon p q$. It is not difficult to show that we can find a family of ellipsoid embeddings $E(sp, sq) \xrightarrow{s} M, s \in [\varepsilon, 1/c]$, which interpolates between these two embeddings (using e.g. the “extension after restriction principle” – see [Sch18, §4.4]). By a compactness argument which extends (i) above, we can then find another curve C' with the same local picture, but where the triangle now has vertices $(0, 0), (p/c, 0), (0, q/c)$. Since the portion of the curve lying over this triangle has symplectic area $\frac{pq}{c}$ (again using Exercise 2.16), we must have $\text{area}(C) > \frac{pq}{c}$.

Exercise 2.16. Let T be the triangle in $\mathbb{R}_{\geq 0}^2$ with vertices $(0, 0), (rp, 0), (0, rq)$ for some $r \in \mathbb{R}_{> 0}$ and coprime $p, q \in \mathbb{Z}_{\geq 1}$, and let $\gamma \subset \mathbb{R}_{\geq 0}^2$ be the shortest line segment from the origin to the hypotenuse of T . Let $C \subset \mathbb{R}^4$ be the visible symplectic curve whose moment map image is γ , such that each fiber is a geodesic in the direction (p, q) in the corresponding torus. Show that $\text{area}(C) = rpq$. *Hint: see [MS25b, §5.1a] for more on visible symplectic curves and their areas.*

(iv) This follows by a version of automatic transversality in dimension four, in the sense of [HLS97, Wen10]. More precisely, a well-known corollary of automatic transversality is that sufficiently nice curves count positively (c.f. [MS24b, §5.2]).

□

Lecture 3: Well-placed curves and their symmetries

§3.1 Looijenga pairs

§3.2 Toric models

§3.3 Elementary transformations

§3.4 Well-placed curves

§3.5 The shift and reflection symmetries

Lecture 4: Scattering diagrams and the tropical vertex

§4.1 Scattering diagram basics

§4.2 Some toy examples

§4.3 The tropical vertex theorem

Lecture 5: Putting it all together

§5.1 The change of lattice trick

§5.2 Proof of Theorem C

References

- [Bir97] Paul Biran. Symplectic packing in dimension 4. *Geometric & Functional Analysis GAFA*, 7(3):420–437, 1997. [7](#)
- [CGH18] Daniel Cristofaro-Gardiner and Richard Hind. Symplectic embeddings of products. *Comment. Math. Helv.*, 93(1):1–32, 2018. [8](#)
- [CGHM18] Daniel Cristofaro-Gardiner, Richard Hind, and Dusa McDuff. The ghost stairs stabilize to sharp symplectic embedding obstructions. *J. Topol.*, 11(2):309–378, 2018. [8](#)
- [CGHMP25] Dan Cristofaro-Gardiner, Tara S Holm, Alessia Mandini, and Ana Rita Pires. On infinite staircases in toric symplectic four-manifolds. *Journal of Differential Geometry*, 129(2):335–413, 2025. [5](#)
- [CM05] K. Cieliebak and K. Mohnke. Compactness for punctured holomorphic curves. *Journal of Symplectic Geometry*, 3(4):589 – 654, 2005. [15](#)
- [CM18] Kai Cieliebak and Klaus Mohnke. Punctured holomorphic curves and Lagrangian embeddings. *Invent. Math.*, 212(1):213–295, 2018. [15](#)
- [CV22] Roger Casals and Renato Vianna. Full ellipsoid embeddings and toric mutations. *Selecta Mathematica*, 28(3):61, 2022. [5](#)
- [Eva23] Jonny Evans. *Lectures on Lagrangian torus fibrations*, volume 105. Cambridge University Press, 2023. [5](#)

- [FdBLMHN06] Javier Fernández de Bobadilla, Ignacio Luengo, Alejandro Melle Hernández, and Andras Némethi. Classification of rational unicuspidal projective curves whose singularities have one Puiseux pair. In *Real and complex singularities*, pages 31–45. Springer, 2006. [10](#)
- [GPS10] Mark Gross, Rahul Pandharipande, and Bernd Siebert. The tropical vertex. *Duke Mathematical Journal*, 153(2):297–362, 2010. [10](#)
- [Gro85] Mikhael Gromov. Pseudo holomorphic curves in symplectic manifolds. *Inventiones Mathematicae*, 82(2):307–347, 1985. [3](#)
- [Gut08] Larry Guth. Symplectic embeddings of polydisks. *Invent. Math.*, 172(3):477–489, 2008. [8](#)
- [Hin15] Richard Hind. Some optimal embeddings of symplectic ellipsoids. *Journal of Topology*, 8(3):871–883, 2015. [8](#)
- [HK14] Richard Hind and Ely Kerman. New obstructions to symplectic embeddings. *Invent. Math.*, 196(2):383–452, 2014. [8](#)
- [HLS97] Helmut Hofer, Véronique Lizan, and Jean-Claude Sikorav. On genericity for holomorphic curves in four-dimensional almost-complex manifolds. *The Journal of Geometric Analysis*, 7:149–159, 1997. [17](#)
- [Kas87] Hiroko Kashiwara. Fonctions rationnelles de type $(0, 1)$ sur le plan projectif complexe. *Osaka J. Math.*, 24(3):521–577, 1987. [11](#)
- [LL01] Tian-Jun Li and Ai-Ko Liu. Uniqueness of symplectic canonical class, surface cone and symplectic cone of 4-manifolds with $b^+ = 1$. *Journal of Differential Geometry*, 58(2):331–370, 2001. [7](#)
- [LL02] Bang-He Li and Tian-Jun Li. Symplectic genus, minimal genus and diffeomorphisms. *Asian Journal of Mathematics*, 6(1):123–144, 2002. [7](#)
- [McD98] Dusa McDuff. From symplectic deformation to isotopy. In *Topics in symplectic 4-manifolds (Irvine, CA, 1996)*, volume I of *First Int. Press Lect. Ser.*, pages 85–99. Int. Press, Cambridge, MA, 1998. [7](#)
- [McD09] Dusa McDuff. Symplectic embeddings of 4-dimensional ellipsoids. *Journal of Topology*, 2(1):1–22, 2009. [6](#)
- [McD18] Dusa McDuff. A remark on the stabilized symplectic embedding problem for ellipsoids. *Eur. J. Math.*, 4(1):356–371, 2018. [8](#)
- [MP94] Dusa McDuff and Leonid Polterovich. Symplectic packings and algebraic geometry. *Inventiones mathematicae*, 115:405–429, 1994. [6](#)
- [MS81] Masayoshi Miyanishi and Tohru Sugie. On a projective plane curve whose complement has logarithmic Kodaira dimension $-\infty$. *Osaka Journal of Mathematics*, 18(1):1 – 11, 1981. [11](#)
- [MS12] Dusa McDuff and Felix Schlenk. The embedding capacity of 4-dimensional symplectic ellipsoids. *Ann. of Math. (2)*, 175(3):1191–1282, 2012. [3](#), [4](#)
- [MS21] Dusa McDuff and Kyler Siegel. Counting curves with local tangency constraints. *Journal of Topology*, 14(4):1176–1242, 2021. [15](#)

- [MS23] Grigory Mikhalkin and Kyler Siegel. Ellipsoidal superpotentials and stationary descendants. *arXiv:2307.13252*, 2023. [9](#), [17](#)
- [MS24a] Dusa McDuff and Kyler Siegel. Sesquicuspidal curves, scattering diagrams, and symplectic nonsqueezing. *arXiv:2412.00561*, 2024. [7](#), [9](#)
- [MS24b] Dusa McDuff and Kyler Siegel. Symplectic capacities, unperturbed curves and convex toric domains. *Geometry & Topology*, 28(3):1213–1285, 2024. [17](#)
- [MS25a] Dusa McDuff and Kyler Siegel. Ellipsoidal superpotentials and singular curve counts. *International mathematics research notices*, 2025. To appear. [9](#), [13](#), [15](#), [16](#)
- [MS25b] Dusa McDuff and Kyler Siegel. Singular algebraic curves and infinite symplectic staircases. *Inventiones Mathematicae*, 2025. To appear. [5](#), [17](#)
- [Ops15] Emmanuel Opshtein. Symplectic packings in dimension 4 and singular curves. *J. Symplectic Geom.*, 13(2):305–342, 2015. [12](#)
- [Ore02] S Yu Orevkov. On rational cuspidal curves: I. Sharp estimate for degree via multiplicities. *Mathematische Annalen*, 324(4):657–673, 2002. [11](#)
- [PS16] Leonid Polterovich and Egor Shelukhin. Autonomous hamiltonian flows, hofer’s geometry and persistence modules. *Selecta Mathematica*, 22(1):227–296, 2016. [3](#)
- [Sch18] Felix Schlenk. Symplectic embedding problems, old and new. *Bull. Amer. Math. Soc. (N.S.)*, 55(2):139–182, 2018. [3](#), [17](#)
- [Sie22] Kyler Siegel. Computing higher symplectic capacities I. *International Mathematics Research Notices*, 2022(16):12402–12461, 2022. [8](#)
- [Sym] Margaret Symington. Four dimensions from two in symplectic topology. topology and geometry of manifolds (athens, ga, 2001), 153–208. In *Proc. Sympos. Pure Math*, volume 71. [5](#)
- [Ton18] Dmitry Tonkonog. String topology with gravitational descendants, and periods of Landau-Ginzburg potentials. *arXiv:1801.06921*, 2018. [15](#)
- [Wen10] Chris Wendl. Automatic transversality and orbifolds of punctured holomorphic curves in dimension four. *Comment. Math. Helv.*, 85(2):347–407, 2010. [17](#)