

# Edit Distance

---

## 1 The Edit Distance Problem

### Definition of a ref-word

Given a finite set of variables  $V \subseteq \text{SVars}$  we define the alphabet of ref words as:  $\Gamma_V := \{x \vdash, \dashv x\}$ . And given an alphabet  $\Sigma$  such that  $\Sigma \cap \Gamma_V = \emptyset$  we can define the set of ref words over  $\Sigma$  and  $V$  as:  $\mathbf{r} \in (\Sigma \cup \Gamma_V)^*$ . Next, a ref-word is valid if and only if, every occurrence of a variable in the ref-word is opened exactly once and closed afterwards, exactly once.

### Functions on ref-words

We can define the projection of a ref word over a set  $S$ ,  $r \uparrow S$ , recursively as:<sup>1</sup>

1.  $r \in S \rightarrow r \uparrow S = r$
2.  $r \notin S \rightarrow r \uparrow S = \epsilon$
3.  $(r_1 \cdot r_2) \uparrow S = (r_1 \uparrow S) \cdot (r_2 \uparrow S)$

$\text{Vars}(r)$  is the set of variables  $x \in V$  that occurs in the ref-word:

$$\text{Vars}(r) := \{x \in V \mid \exists r_x^{pre}, r_x, r_x^{post} \in (\Sigma \cup \Gamma_V)^* \text{ such that } r = r_x^{pre} \cdot x \vdash \cdot r_x \cdot \dashv x \cdot r_x^{post}\} \quad (1)$$

$\text{tup}(r)$  are the positions each ref-word is referencing, and is defined as:

$$\text{tup}(r) := \{x \mapsto [i_x, j_x] \mid x \in \text{Vars}(R), i_x = |r_x^{pre} \uparrow \Sigma|, j_x = i_x + |r_x \uparrow \Sigma|\} \quad (2)$$

**Postulate:**

$$\text{valid}(r) \rightarrow |\text{tup}(r)| = |\text{Vars}(r)| \quad (3)$$

### Definition of ref-word tuple

### Distance between two ref words

Next, given two ref words  $r_1, r_2 \in (\Sigma \cup \Gamma_V)^*$  and a distance function  $\mathbf{d} : \Sigma^* \times \Sigma^* \rightarrow \mathbb{R}$  the distance  $\mathbf{d}_\Gamma$  between  $r_1$  and  $r_2$  is defined as:  $\mathbf{d}_\Gamma(r_1, r_2) = \mathbf{d}(r_1 \uparrow \Sigma, r_2 \uparrow \Sigma)$

### Ref-word distance languages

Given a ref-word language  $R \subseteq (\Sigma \cup \Gamma_V)^*$  and a distance  $k \in \mathbb{R}$ , the k-distance ref-word language is defined as:

$$R \pm k = \{r \in (\Sigma \cup \Gamma_V)^* \mid \text{valid}(r), \exists r' (r' \in R \wedge \mathbf{d}_\Gamma(r, r') \leq k)\} \quad (4)$$

Given a document  $d$ , the spanner over a ref-word language  $R$  is:

$$\llbracket R \rrbracket_d = \{\text{tup}(r) \mid r \uparrow \Sigma = d, \exists r \in R\} \quad (5)$$

---

<sup>1</sup>In the paper (Doleschal, 2021) this operation is defined for  $\Sigma$  as  $\text{doc}(\sigma)$

**Theorem 1.** *If the distance function  $\mathbf{d}$  is a metric, then:*

$$(R \pm n) \pm k = R \pm (n + k) \quad (6)$$

*Proof.* Given  $n, k \in \mathbb{R}$ , we want to prove that:  $(R \pm n) \pm k = R \pm (n + k)$ . By definition, we have that:

$$(R \pm n) \pm k = \{r \in (\Sigma \cup \Gamma_V)^* \mid \text{valid}(r), \exists r' (r' \in R \pm n \wedge \mathbf{d}_{\mathbf{r}}(r, r') \leq k)\}$$

First we prove that  $(R \pm n) \pm k \subseteq R \pm (n + k)$ . By contradiction let's assume there exists an element  $r_1 \in (R \pm n) \pm k$  such that  $r_1 \notin R \pm (n + k)$ . By the previous definition, we have that  $\exists r' \in R \pm n$  such that  $\mathbf{d}_{\mathbf{r}}(r_1, r') \leq k$ . By definition of  $R \pm (n)$  we have that for any  $r \in R$ ,  $\mathbf{d}_{\mathbf{r}}(r', r) \leq n$ . Next, by the definition of  $R \pm (n + k)$ , and our supposition we have that  $\mathbf{d}_{\mathbf{r}}(r_1, r) > n + k$  which contradicts the triangle inequality. The proof that  $R \pm (n + k) \subseteq (R \pm n) \pm k$  uses this same argument.  $\square$

## Variable-set automaton over ref words (VSet-automaton)<sup>2</sup>

**Definition 1.** A VSet-automaton is a sextuple  $A := (\Sigma, V, Q, q_0, Q_F, \delta)$

- $\Sigma$ : Alphabet symbols
- $V$ : Finite set of variables
- $Q$ : Finite set of states
- $q_0 \in Q$ : Initial state
- $Q_F \subseteq Q$ : Set of final states
- $\delta : Q \times (\Sigma \cup \{\epsilon\} \cup \Gamma_V) \rightarrow 2^Q$ : Transition function
  - $\Gamma_V := \{x \vdash, \dashv x \mid x \in V\}$
  - $2^Q$ : power set of  $Q$

### Ref-word language

The ref-word language of  $A$  is:  $\mathcal{R}(A) = \mathcal{R}^0(A) = \{r \in \mathcal{L}(A) \subseteq (\Sigma \cup \Gamma_V)^* \mid r \text{ is accepted by the } \epsilon - \text{NFA } A\}$ . This is direct from interpreting  $A$  as an  $\epsilon - \text{NFA}$ .

### Run of a VSet-automaton over a ref-word

Given a ref-word  $r = \sigma_1 \cdots \sigma_n$ , the run  $\rho$  of  $A$  is the sequence:

$$\rho := q_0 \xrightarrow{\sigma_1} q_1 \cdots q_{n-1} \xrightarrow{\sigma_n} q_n \quad (7)$$

Where  $\forall i \in [0, n) (q_{i+1} \in \delta(q_i, \sigma_{i+1}))$  and  $q_n \in Q_F$

From previous publications we know that  $r \in \mathcal{R}(A)$  if and only if there is a run  $\rho$  of  $A$  on  $r$ .

---

<sup>2</sup>(Doleschal, 2021)

## Distance automaton

Given a VSet-automaton  $A$  we can define, under Levenshtein distance, the automaton  $A \pm 1 := (\Sigma, V, Q', q'_0, Q'_F, \delta')$  Where

- $Q' = \{q_1, \dots, q_{|Q|}\} \cup \{q_1^1, \dots, q_{|Q|}^1\}$  Where there exists two bijective functions:

1.  $f : Q \rightarrow \{q_1, \dots, q_{|Q|}\}$
2.  $f' : Q \rightarrow \{q_1^1, \dots, q_{|Q|}^1\}$

And two bijective functions  $F$  and  $F'$  that map  $f$  and  $f'$  respectively to sets.

- $q'_0 = f(q_0)$
- $Q'_F = Q_F \cup \{q_i^1 \mid q_j \in Q_F \wedge f'(q_j) = q_i^1\}$
- The function  $\delta'$  is defined by:

$$\delta'(q_i, e) = \begin{cases} F(\delta(f^{-1}(q_i), e)) & e \in \Gamma_V \\ F(\delta(f^{-1}(q_i), e)) \cup q_i^1 \cup \bigcup_{a \in \Sigma} F'(\delta(f^{-1}(q_i), a)) & e.o.c \end{cases}$$

$$\delta'(q_i^1, e) = F'(\delta(f'^{-1}(q_i^1), e))$$

**Definition 2.** A ref-word language  $R$  is sequential if every ref-word  $r \in R$  is valid.

**Lemma 1.** All runs of  $\mathcal{R}(A)$  have one of the following structures:

1.

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \dots \xrightarrow{c_{a-1}} f(\phi_a) \xrightarrow{c_a} f'(\phi_a) \xrightarrow{c_{a+1}} f'(\phi_{a+1}) \dots \xrightarrow{c_n} f'(\phi_{n-1})$$

2.

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \dots \xrightarrow{c_{a-1}} f(\phi_a) \xrightarrow{c_a} f'(\phi_{a+1}) \xrightarrow{c_{a+1}} f'(\phi_{a+2}) \dots \xrightarrow{c_n} f'(\phi_n)$$

3.

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \dots \xrightarrow{c_{a-1}} f(\phi_a) \xrightarrow{c_a} f'(\phi_{a+2}) \dots \xrightarrow{c_n} f'(\phi_{n+1})$$

4.

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \xrightarrow{c_2} \dots \xrightarrow{c_n} f(\phi_n)$$

*Proof.* This is direct from the form of the transitions under the definition of  $A \pm 1$ , and noticing that there is no transition from  $f'(q)$  to  $f(q)$ .  $\square$

**Theorem 2.** if  $\mathcal{R}(A)$  is sequential, then  $\mathcal{R}(A \pm 1)$  is sequential.

*Proof.* For purposes of contradiction let's assume that there exists a ref-word  $r = c_1 \dots c_n \in A \pm 1$  that is not valid. Then there must exist a run  $\rho_{A \pm 1}$  on  $r$ . Furthermore, If  $r = c_1 \dots c_n$  is not valid, then there are two cases:

1.  $\exists c_i, c_j$  such that  $i < j$  and  $c_i = x \dashv, c_j = x \vdash$

Under the first case of lemma ??,  $c_a \notin \Gamma_V$  from the definition of  $A \pm 1$ . Therefore,

2.  $\exists c_i, c_j$  such that  $c_i = c_j \in \Gamma_V$

$\square$

**Theorem 3.** Given a VSet-automaton with a sequential ref-word language  $\mathcal{R}(A)$ , using Levenshtein distance we obtain that  $\mathcal{R}(A \pm 1) = \mathcal{R}(A) \pm 1$ .

*Proof.* This is equivalent to proving that, given a ref-word  $r$ ,  $r \in \mathcal{R}(A \pm 1) \leftrightarrow r \in \mathcal{R}(A) \pm 1$ . Therefore this is a two part proof.

First let's assume that  $r \in \mathcal{R}(A \pm 1)$ . In that case we know that there must exist a run  $\rho$  of  $A \pm 1$  on  $r$ .  $r$  is valid due to the theorem ???. By definition of  $\delta'$  there is no transition from states of the form  $q_i^1$  to  $q_i$ . Therefore, there are two cases:

1. There are only states of the form  $q_i$ . In this case, the subset of nodes and transitions used have an isomorphism to  $A$  using the function  $f$ , therefore,  $r \in \mathcal{R}(A)$ , using this and that  $r$  is valid we conclude that  $r \in \mathcal{R}(A) \pm 1$
2. There is one transition from a state of the form  $q_i$  to  $q_i^1$ . The respective run of  $A \pm 1$  on  $r = c_1 \cdots c_n$  could have one of three forms:

**Insertion**

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \cdots \xrightarrow{c_{i-1}} f(\phi_i) \xrightarrow{c_i} f'(\phi_i) \xrightarrow{c_{i+1}} f'(\phi_{i+1}) \cdots \xrightarrow{c_n} f'(\phi_n) \quad (8)$$

In this case, there exists a run:

$$\rho_A = q_0 \xrightarrow{c_1} \phi_1 \cdots \xrightarrow{c_{i-1}} \phi_i \xrightarrow{c_{i+1}} \phi_{i+1} \cdots \xrightarrow{c_n} \phi_n \quad (9)$$

Such that every transition is in  $A$ , therefore the word  $r' = c_1 \cdots c_{i-1} \cdot c_{i+1} \cdots c_n$  is in  $\mathcal{R}(A)$ . And therefore by Levenshtein distance we obtain that  $\mathbf{d}_\Gamma(r', r) = 1$ . And because of this and because  $r$  is valid:  $r \in \mathcal{R}(A) \pm 1$

**Substitution**

TODO

**Elimination**

TODO

Next, let's assume that  $r \in \mathcal{R}(A) \pm 1$ . In that case,  $r$  is valid, and there exists  $r' \in R$  such that using Levenshtein distance:  $\mathbf{d}_\Gamma(r, r') \leq 1$ . Since Levenshtein distance is discrete, there are two cases:

1.  $\mathbf{d}_\Gamma(r, r') = 0$ . Then it is clear that  $r \in \mathcal{R}(A \pm 1)$  since a subset of  $A \pm 1$  forms an isomorphism with  $A$ .
2.  $\mathbf{d}_\Gamma(r, r') = 1$ . In this case, because of the structure of Levenshtein's distance, there are three possible cases:

- (a) **Insertion**  $r = c_1 \cdots c_i \cdot c_{inserted} \cdot c_{i+1} \cdots c_n$  And  $r' = c_1 \cdots c_n$

In this case, because there exists a run  $\rho_A$  on  $r'$  of the form:

$$\rho_A = q_0 \xrightarrow{c_1} \phi_1 \cdots \phi_i \xrightarrow{c_i} \phi_{i+1} \cdots \phi_n$$

Therefore there exists a run  $\rho_{A \pm 1}$  on  $r$ , and it is:

$$\rho_{A \pm 1} = f(q_0) \xrightarrow{c_1} f(\phi_1) \cdots f(\phi_i) \xrightarrow{c_{ins}} f'(\phi_i) \xrightarrow{c_i} f'(\phi_{i+1}) \cdots \xrightarrow{c_n} f'(\phi_n)$$

- (b) **Elimination** TODO, Same as above

- (c) **Substitution** TODO, Same as above

□

**Theorem 4.** For any sequential automaton  $A$  there exists an automaton  $B$  such that  $\mathcal{R}(B) = \mathcal{R}(A) \pm k$  for all  $k \in \mathbb{N}$  under Levenshtein distance.

*Proof.* Trivial proof using theorems 1 and 3, and induction over  $k$ . □