

**Search for high-mass leptoquark with
tau-lepton and missing transverse
momentum using 140 fb of pp collision
data at $\sqrt{s} = 13$ TeV in the LHC-ATLAS
experiment**

(LHC-ATLAS実験における重心系エネルギー13 TeVの140 fbの陽子陽子衝突データを用いたタウレプトンおよび消失横運動量を使った高質量レプトクォーク探索)

Jiaqi Zang

藏 嘉琦

Department of Physics, Graduate School of Science,
The University of Tokyo

This dissertation is submitted for the degree of
Doctor of Philosophy

August 2026

Acknowledgements

¹⁰ To be filled

11

Abstract

12 To be filled

13 Table of contents

14	List of figures	xi
15	List of tables	xiii
16	1 Introduction	1
17	1.1 Introduction	1
18	2 Theoretical Background	5
19	2.1 The Standard Model	5
20	2.1.1 Particle content and gauge group of the SM	6
21	2.1.2 Local gauge invariance	7
22	2.1.3 The $SU(3)_C$ gauge sector	8
23	2.1.4 The electroweak gauge sector	9
24	2.1.5 Covariant derivative and gauge field strengths	11
25	2.1.6 The Standard Model Lagrangian	12
26	2.1.7 Electroweak symmetry breaking and the Higgs mechanism	14
27	2.1.8 Fermion masses and flavour mixing	16
28	2.2 Beyond the Standard Model	17
29	2.2.1 Gravity and the Planck scale	17
30	2.2.2 The hierarchy problem	17
31	2.2.3 Neutrino masses	18
32	2.2.4 Dark matter	19
33	2.2.5 Matter–antimatter asymmetry	19
34	2.2.6 The strong CP problem	19
35	2.2.7 Flavour anomalies and lepton flavour universality	20
36	2.3 Possible Solutions to the Remaining Problems of the Standard Model	21
37	2.3.1 Supersymmetry	21
38	2.3.2 Axions and the strong CP problem	22

39	2.3.3	Leptoquarks	22
40	3	Leptoquark Search at the LHC	25
41	3.1	Leptoquark Models and Phenomenology	25
42	3.1.1	General properties of leptoquarks	25
43	3.1.2	The simplified U_1 leptoquark model	26
44	3.1.3	Low-energy fit and benchmark scenarios	29
45	3.1.4	Leptoquark searches at the LHC	32
46	3.1.5	Signal model and target signatures	34
47	4	LHC-ATLAS Experiment	35
48	4.1	Large Hadron Collider	35
49	4.2	ATLAS experiment	38
50	4.2.1	Overview	38
51	4.2.2	Coordinate system	39
52	4.2.3	Magnet system	39
53	4.2.4	Inner Detector	40
54	4.2.5	Calorimeter	42
55	4.2.6	Muon Spectrometer	45
56	4.2.7	Data Acquisition and Trigger System	47
57	5	Data and Monte Carlo Simulations	51
58	5.1	Dataset selection	51
59	5.2	Signal samples	52
60	5.3	Background samples	54
61	5.4	Triggers	55
62	6	Object Reconstruction and Identification	59
63	6.1	Track	59
64	6.2	Primary vertex	59
65	6.3	Electron	59
66	6.4	Muon	59
67	6.5	Hadronic tau	59
68	6.6	Jet	59
69	6.7	Jet flavor tagging	59
70	6.8	Overlap removal	59
71	6.9	Missing transverse momentum	59

72	7 Event Selection	61
73	7.1 Preselection and variables definition	61
74	7.1.1 Analysis variables	61
75	7.1.2 Multijet suppression	63
76	7.1.3 Common preselection	63
77	7.2 One b-jet signal region (SR1b) optimisation and definition	64
78	7.2.1 Scan variables and optimisation criterion	65
79	7.2.2 Signal efficiency and selected regions	66
80	7.3 b-veto signal region definition (SR0b)	68
81	8 Background Estimation Strategy	71
82	8.1 Strategy overview	71
83	8.2 W +jets estimation	71
84	8.3 $t\bar{t}$ estimation	71
85	8.4 Fake- τ estimation	71
86	8.5 Multi-jet cleaning	71
87	8.6 VRs	71
88	9 Systematics Uncertainties	73
89	9.1 Experimental uncertainties	73
90	9.2 Theoretical uncertainties	73
91	10 Statistical Analysis and Results	75
92	10.1 Statistical analysis	75
93	10.1.1 Hypothesis test	75
94	10.1.2 Likelihood function	75
95	10.2 Results	75
96	10.2.1 Background-only fit	75
97	10.2.2 Unblinded S+B fit in CR+SRs	75
98	10.2.3 Exclusion limits with toy models	75
99	11 Discussion and Conclusion	77
100	References	79
101	Appendix A Correction of the signal internal event weights	87

102	Appendix B Multijet background study	89
103	B.1 Origin of the discriminating variables	89
104	B.2 Study selection and distributions	89
105	B.3 Threshold scan and signal retention	91
106	B.4 Residual multijet contribution	92
107	Appendix C Zero-b-jet signal-region optimisation	93

List of figures

109	2.1	Average of $R(D^*)$ and $R(D)$	21
110	3.1	Fit results in (C_{LL}^c, C_{LR}^c) plane using low-energy observables	30
111	3.2	Example Feynman diagrams of LQ production and exchange process for	
112		$\tau\tau(+b)$ searches.	32
113	3.3	ATLAS constraints on the U_1 leptoquark	33
114	3.4	Signal topologies in the $\tau\nu(+b)$ final state	34
115	4.1	Schematic overview of the CERN accelerator complex.	36
116	4.2	The schematic view of the ATLAS detector and the subdetector systems . .	38
117	4.3	The coordinate system used in ATLAS	40
118	4.4	ATLAS magnet system	41
119	4.5	ATLAS Inner Detector	42
120	4.6	ATLAS calorimeter system	43
121	4.7	Photograph of EM barrel and end-cap LAr modules	44
122	4.8	A cut-away view of an LAr electromagnetic calorimeter module	44
123	4.9	ATLAS Muon Spectrometer	46
124	4.10	ATLAS Run-2 trigger and data acquisition system	47
125	5.1	Run 2 single-lepton trigger efficiencies	57
126	5.2	Run 2 missing-transverse-momentum trigger efficiencies	58
127	7.1	Discriminating-variable distributions in SR1b	67
128	7.2	Discriminating-variable distributions in SR0b	69
129	B.1	Variables used for multijet suppression	90
130	B.2	Scan of the multijet rejection threshold	91

List of tables

132	2.1	Standard Model particle content	7
133	5.1	Integrated luminosity of the Run 2 proton–proton collision data used in this	
134		analysis.	52
135	5.2	Summary of the principal MC generator configurations used for the signal	
136		and SM background processes. The quoted perturbative order refers to the	
137		hard-scattering matrix-element calculation. For the $W/Z + \text{jets}$ samples,	
138		matrix elements with up to two additional partons are calculated at NLO	
139		and those with up to five additional partons at LO. The symbol ℓ denotes an	
140		electron or muon.	55
141	5.3	Run 2 single-lepton and missing-transverse-momentum trigger chains . . .	56
142	7.1	Common preselection for the signal-region optimisation. Object definitions	
143		and overlap removal are given in Chapter 6.	64
144	7.2	Thresholds tested in the SR1b optimisation. Each set lists the trial values for	
145		the indicated inequality. A zero mass threshold or an angular upper threshold	
146		of π represents no effective requirement on that variable.	65
147	7.3	Final signal-region selections, applied in addition to the common preselection	
148		in Table 7.1. A dash denotes no additional requirement. The jet multiplicity	
149		includes b -tagged jets.	69
150	B.1	Loose selection used to study the multijet suppression. This selection defines	
151		the denominator of the rejection fractions shown below.	90
152	C.1	Signal benchmarks and preferred thresholds in the supplementary SR0b	
153		E_T^{miss} scan. The last column gives the thresholds used in the final analysis. .	93

Chapter 1

Introduction

1.1 Introduction

Modern high-energy physics seeks to understand the elementary constituents of matter and the fundamental interactions that govern their behaviour. At present, the most successful theoretical description of particle physics is the Standard Model (SM), a quantum field theory based on the gauge symmetry

$$SU(3)_C \times SU(2)_L \times U(1)_Y. \quad (1.1)$$

Within this framework, matter is described by three generations of quarks and leptons, while the electromagnetic, weak, and strong interactions are mediated by gauge bosons. The electroweak symmetry is spontaneously broken through the Higgs mechanism, giving masses to the weak gauge bosons and to the fermions through their interactions with the Higgs field. The discovery of the Higgs boson by the ATLAS and CMS experiments in 2012 completed the particle content predicted by the SM and marked a major milestone in modern particle physics.

The SM has been tested with remarkable precision over a wide range of energy scales and has successfully described a vast number of experimental observations. Nevertheless, it is widely regarded as an incomplete theory. It does not provide a particle candidate for dark matter, does not explain the observed matter–antimatter asymmetry of the universe, and does not account for the origin of non-zero neutrino masses. In addition, the pattern of fermion masses and flavour mixing remains unexplained, and the large separation between the electroweak and Planck scales suggests the possible existence of new physics beyond the SM. These open questions motivate direct searches for new particles and interactions at the highest experimentally accessible energies.

The Large Hadron Collider (LHC) at CERN provides a unique environment for probing such new phenomena. By colliding protons at tera-electronvolt centre-of-mass energies, the LHC allows the production of particles that may be associated with physics beyond the SM. During Run 2 of the LHC from year 2015 to 2018, proton–proton (pp) collisions were delivered at a centre-of-mass energy of $\sqrt{s} = 13$ TeV.

ATLAS is a detector located at one of the interaction points of the LHC. It is designed to observe and study a wide range of physics processes, mainly from the pp collisions, from precision measurements of SM interactions to searches for new particles. The detector consists of several sub-detectors surrounding the collision interaction point, including an inner tracking detector, electromagnetic and hadronic calorimeters, and a muon spectrometer. Together with a powerful magnet system, these components allow the reconstruction and identification of electrons, photons, muons, hadronically decaying τ -leptons, jets, and missing transverse momentum, denoted by E_T^{miss} .

Among the many possible extensions of the SM, leptoquarks (LQs) provide a well-motivated class of new particles connecting the quark and lepton sectors. They are hypothetical bosons that couple directly to a quark and a lepton, and they appear naturally in theories such as grand unified theories and models of quark–lepton unification. Leptoquarks are also of particular phenomenological interest because they can modify semileptonic processes and may therefore be connected to the observed tensions in flavour observables.

A search for the leptoquarks is presented in the thesis using 140 fb^{-1} of pp collision data recorded by the ATLAS detector at a centre-of-mass energy of $\sqrt{s} = 13$ TeV. This thesis focuses on the vector singlet leptoquark U_1 as the benchmark model. This model can contribute to processes involving b quarks, τ leptons, and neutrinos. Therefore, the analysis targets events with a hadronically decaying τ -lepton, large missing transverse momentum, and jets. The observed data are compared with SM background predictions in signal regions designed to enhance sensitivity to possible leptoquark contributions. In the absence of a significant excess over the SM expectation, constraints are placed on the allowed parameter space of the U_1 model, expressed in terms of the leptoquark mass and its couplings.

This dissertation is organised as follows. Chapter 2 provides an overview of the SM’s theoretical framework and the physics beyond it. Chapter 3 describes the leptoquark models and the searches performed at LHC. Chapter 4 introduces the LHC and the ATLAS detector. Chapter 5 provides the dataset and simulated samples. Chapter 6 details the reconstruction and the identification of the physics objects that are used for event selection. Chapter 7 discusses the optimisation for the event selection used for signal signature extraction. Chapter 8 presents the background estimation strategy. Chapter 9 discusses the systematic

213 uncertainties for the background and signals. Chapter 10 presents the statistical interpretation
214 of the results. Chapter 11 summarises this analysis.

Chapter 2

Theoretical Background

This chapter introduces the theoretical background which motivates this search performed in the thesis. The discussion begins with the Standard Model (SM), which provides the current theoretical framework for describing elementary particles and their interactions. The limitations of the SM are then summarised, followed by a discussion of several possible extensions beyond the SM. Particular emphasis is placed on leptoquark models, which provide a direct connection between the quark and lepton sectors and motivate the experimental search presented in this thesis.

2.1 The Standard Model

The Standard Model (SM) is a relativistic quantum field theory describing the known elementary particles and their strong, weak, and electromagnetic interactions. Its structure is determined by local gauge invariance under

$$SU(3)_C \times SU(2)_L \times U(1)_Y. \quad (2.1)$$

The group $SU(3)_C$ describes the colour symmetry of the strong interaction, while $SU(2)_L \times U(1)_Y$ describes the electroweak interaction [1–3]. In quantum field theory, particles are classified not only by their mass and spin, but also by their transformation properties under these gauge symmetries. Particles with half-integer spin are fermions and obey Fermi–Dirac statistics, while particles with integer spin are bosons and obey Bose–Einstein statistics [4]. The matter particles of the SM are spin-1/2 fermions, the interaction carriers are spin-1 gauge bosons, and the Higgs boson is a spin-0 scalar boson.

The minimal SM Lagrangian used here is constructed from the SM field content and is invariant under the local gauge group. Therefore, instead of introducing the Lagrangian as

an isolated expression, this section first summarises the field content and gauge quantum numbers of the SM. The principle of local gauge invariance is then introduced, followed by the gauge transformations of the strong sector, based on $SU(3)_C$, and of the electroweak sector, based on $SU(2)_L \times U(1)_Y$. The covariant derivative and field-strength tensors are then constructed from these symmetries, leading to the full SM Lagrangian. Finally, electroweak symmetry breaking and the generation of fermion masses are discussed.

2.1.1 Particle content and gauge group of the SM

The gauge quantum numbers of a field specify how it transforms under the SM gauge group. In this thesis, the electric charge is written as

$$Q = T^3 + Y, \quad (2.2)$$

where T^3 is the third component of weak isospin and Y is the weak hypercharge. This convention is used throughout the thesis.

For each generation, the SM fermions are arranged as

$$Q_L^i = \begin{pmatrix} u_L^i \\ d_L^i \end{pmatrix} \sim (3, 2, 1/6), \quad u_R^i \sim (3, 1, 2/3), \quad d_R^i \sim (3, 1, -1/3), \quad (2.3)$$

$$L_L^i = \begin{pmatrix} \nu_L^i \\ e_L^i \end{pmatrix} \sim (1, 2, -1/2), \quad e_R^i \sim (1, 1, -1), \quad (2.4)$$

where $i = 1, 2, 3$ is the generation index, and the notation denotes the representation under $SU(3)_C \times SU(2)_L \times U(1)_Y$, which are the QCD representation, the weak isospin representation and the weak hypercharge, respectively. In the minimal SM, no right-handed neutrino field is included. The scalar sector contains one complex Higgs doublet,

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \sim (1, 2, 1/2). \quad (2.5)$$

The gauge bosons are associated with the generators of the three gauge groups. The eight gluon fields G_μ^a , with $a = 1, \dots, 8$, are associated with $SU(3)_C$. The three weak gauge fields W_μ^I , with $I = 1, 2, 3$, are associated with $SU(2)_L$. The hypercharge gauge field B_μ is associated with $U(1)_Y$. Before electroweak symmetry breaking, W_μ^1 , W_μ^2 , W_μ^3 , and B_μ are gauge eigenstates. After electroweak symmetry breaking, they are reorganised into the physical electroweak bosons $W^\pm$, Z , and γ .

The main particle content of the SM is summarised in Table 2.1. For fermions and the Higgs boson, the table lists the corresponding electroweak quantum numbers using the convention in Eq. (2.2). For gauge bosons, the entries describe their gauge fields and the electric charges of the physical states where appropriate.

Table 2.1 Particle content of the SM and the corresponding spin and electroweak quantum numbers. For fermions and the Higgs doublet, T^3 and Y denote the third component of weak isospin and the weak hypercharge, respectively, with the electric charge defined as $Q = T^3 + Y$. For gauge bosons, only the electric charges of the physical states after electroweak symmetry breaking are shown.

Category	Type	Generation			Spin	Quantum numbers		
		1st	2nd	3rd		T^3	Y	Q
Fermion	Left-handed quark	u_L	c_L	t_L	1/2	+1/2	1/6	+2/3
		d_L	s_L	b_L		-1/2		-1/3
	Right-handed up-type quark	u_R	c_R	t_R	1/2	0	2/3	+2/3
	Right-handed down-type quark	d_R	s_R	b_R	1/2	0	-1/3	-1/3
	Left-handed lepton	ν_{eL}	$\nu_{\mu L}$	$\nu_{\tau L}$	1/2	+1/2	-1/2	0
		e_L	μ_L	τ_L		-1/2		-1
	Right-handed charged lepton	e_R	μ_R	τ_R	1/2	0	-1	-1
Gauge boson	Gluon	$g^a, a = 1, \dots, 8$			1	-	-	0
	Photon	γ			1	-	-	0
	Charged weak boson	$W^\pm$			1	-	-	± 1
	Neutral weak boson	Z			1	-	-	0
Scalar field	Higgs doublet	ϕ^+			0	+1/2	1/2	+1
		ϕ^0				-1/2		0

The quantum numbers listed above determine which interactions each field can have. They also determine which terms can be written in the SM Lagrangian. For example, explicit Dirac mass terms for charged fermions are not gauge invariant before electroweak symmetry breaking, because the left- and right-handed fermions transform differently under $SU(2)_L \times U(1)_Y$. Fermion masses instead arise from gauge-invariant Yukawa interactions with the Higgs field.

2.1.2 Local gauge invariance

The guiding principle in constructing the SM is local gauge invariance. A global transformation acts in the same way at every point in spacetime, while a local gauge transformation depends on the spacetime coordinate x . For a field $\psi(x)$ transforming under a gauge group with generators T^A , a generic local transformation is

$$\psi(x) \rightarrow \psi'(x) = U(x)\psi(x), \quad U(x) = \exp \left[i\alpha^A(x)T^A \right]. \quad (2.6)$$

Here $\alpha^A(x)$ are arbitrary real, spacetime-dependent gauge parameters. They specify how the field is transformed locally along each generator T^A of the gauge group, and the repeated index A is summed over all generators.

The ordinary derivative of the transformed field is

$$\partial_\mu \psi'(x) = U(x) \partial_\mu \psi(x) + (\partial_\mu U(x)) \psi(x). \quad (2.7)$$

The second term in Eq. (2.7) shows that $\partial_\mu \psi$ does not transform in the same way as ψ . Therefore, the kinetic term constructed using the ordinary derivative is not invariant under a local gauge transformation.

To restore local gauge invariance, the ordinary derivative is promoted to a covariant derivative D_μ satisfying

$$D_\mu \psi(x) \rightarrow D'_\mu \psi'(x) = U(x) D_\mu \psi(x). \quad (2.8)$$

This requirement introduces gauge fields. The gauge fields are therefore not added by hand as arbitrary new particles; they are required in order to compare fields at neighbouring spacetime points in a way that is independent of the local gauge convention. The following subsections describe how this principle is realised for the strong and electroweak gauge sectors of the SM.

2.1.3 The $SU(3)_C$ gauge sector

The strong interaction is described by quantum chromodynamics (QCD), a gauge theory based on the colour group $SU(3)_C$. Quark fields transform as colour triplets, while leptons and the Higgs field are colour singlets. For a colour triplet quark field $q(x)$, the local $SU(3)_C$ transformation is

$$q(x) \rightarrow q'(x) = U_C(x) q(x), \quad U_C(x) = \exp \left[i \alpha_C^a(x) \frac{\lambda^a}{2} \right], \quad (2.9)$$

where λ^a are the Gell-Mann matrices and $a = 1, \dots, 8$. The corresponding gauge fields are the eight gluon fields G_μ^a .

It is useful to introduce the matrix-valued gluon field

$$G_\mu(x) = G_\mu^a(x) \frac{\lambda^a}{2}. \quad (2.10)$$

With the convention

$$D_\mu = \partial_\mu - ig_s G_\mu^a \frac{\lambda^a}{2}, \quad (2.11)$$

where g_s is the strong coupling constant, the gluon field transforms as

$$G_\mu(x) \rightarrow G'_\mu(x) = U_C(x) G_\mu(x) U_C^\dagger(x) - \frac{i}{g_s} (\partial_\mu U_C(x)) U_C^\dagger(x). \quad (2.12)$$

This transformation law ensures that $D_\mu q$ transforms in the same way as q under local $SU(3)_C$ transformations.

The $SU(3)_C$ field-strength tensor is

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f^{abc} G_\mu^b G_\nu^c, \quad (2.13)$$

where f^{abc} are the structure constants of $SU(3)$, defined by the commutation relation

$$\left[\frac{\lambda^a}{2}, \frac{\lambda^b}{2} \right] = i f^{abc} \frac{\lambda^c}{2}. \quad (2.14)$$

The last term in Eq. (2.13) arises from the non-Abelian nature of $SU(3)_C$. It gives rise to gluon self-interactions, which are a defining feature of QCD and are essential for asymptotic freedom at high energies and confinement at low energies.

2.1.4 The electroweak gauge sector

Before electroweak symmetry breaking, the weak and electromagnetic interactions are described together by the gauge group $SU(2)_L \times U(1)_Y$ [1–3]. The $SU(2)_L$ factor is the non-Abelian weak-isospin gauge symmetry, while the $U(1)_Y$ factor is the Abelian hypercharge gauge symmetry. They are introduced separately below, but the physical photon and Z boson arise only after electroweak symmetry breaking, when the neutral fields W_μ^3 and B_μ mix.

The $SU(2)_L$ gauge transformation

The $SU(2)_L$ gauge symmetry acts on left-handed fermion doublets and on the Higgs doublet:

$$Q_L^i, \quad L_L^i, \quad \Phi. \quad (2.15)$$

For an $SU(2)_L$ doublet $\chi(x)$, the local transformation is

$$\chi(x) \rightarrow \chi'(x) = U_L(x)\chi(x), \quad U_L(x) = \exp \left[i\alpha^I(x) \frac{\tau^I}{2} \right], \quad (2.16)$$

where τ^I are the Pauli matrices and $I = 1, 2, 3$. The right-handed charged fermions u_R^i, d_R^i , and e_R^i are singlets under $SU(2)_L$ and therefore have $T_L^I = 0$.

The three gauge fields associated with $SU(2)_L$ are denoted by W_μ^I . They can be written in matrix form as

$$W_\mu(x) = W_\mu^I(x) \frac{\tau^I}{2}. \quad (2.17)$$

With the convention

$$D_\mu = \partial_\mu - igW_\mu^I \frac{\tau^I}{2}, \quad (2.18)$$

where g is the $SU(2)_L$ coupling constant, the weak gauge field transforms as

$$W_\mu(x) \rightarrow W'_\mu(x) = U_L(x)W_\mu(x)U_L^\dagger(x) - \frac{i}{g}(\partial_\mu U_L(x))U_L^\dagger(x). \quad (2.19)$$

This transformation law is the non-Abelian generalisation of the Abelian gauge-field transformation. The matrix structure reflects the fact that the $SU(2)_L$ generators do not commute.

The $SU(2)_L$ field-strength tensor is

$$W_{\mu\nu}^I = \partial_\mu W_\nu^I - \partial_\nu W_\mu^I + g\epsilon^{IJK}W_\mu^J W_\nu^K. \quad (2.20)$$

The last term arises from the non-Abelian nature of $SU(2)_L$ and leads to self-interactions among weak gauge bosons.

The $U(1)_Y$ gauge transformation

The $U(1)_Y$ gauge symmetry is the Abelian part of the electroweak gauge group. Its corresponding gauge charge is the weak hypercharge Y , which determines how each field transforms under a local $U(1)_Y$ phase rotation. For a field $\psi(x)$ with hypercharge Y , the local $U(1)_Y$ transformation is

$$\psi(x) \rightarrow \psi'(x) = \exp[iY\beta(x)]\psi(x), \quad (2.21)$$

where $\beta(x)$ is a spacetime-dependent transformation parameter. The corresponding gauge field is the hypercharge field B_μ . With the convention

$$D_\mu = \partial_\mu - ig'YB_\mu, \quad (2.22)$$

the gauge field transforms as

$$B_\mu(x) \rightarrow B'_\mu(x) = B_\mu(x) + \frac{1}{g'}\partial_\mu\beta(x). \quad (2.23)$$

This transformation ensures that $D_\mu\psi$ transforms in the same way as ψ .

The hypercharge assignments in Table 2.1 are chosen so that the electric charge is obtained from Eq. (2.2). For the left-handed lepton doublet, $Y = -1/2$. The upper component has $T^3 = +1/2$ and therefore $Q = 0$, corresponding to the neutrino. The lower component has $T^3 = -1/2$ and therefore $Q = -1$, corresponding to the charged lepton. Similarly, the left-handed quark doublet has $Y = 1/6$, giving charges $+2/3$ and $-1/3$ for the up- and down-type components.

Because $U(1)_Y$ is Abelian, the corresponding field-strength tensor is

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (2.24)$$

There is no self-interaction term involving two B_μ fields in $B_{\mu\nu}$. Although $U(1)_Y$ is Abelian, it should not be identified with the electromagnetic gauge symmetry $U(1)_{\text{em}}$. Before electroweak symmetry breaking, the corresponding gauge field is the hypercharge field B_μ rather than the photon field. The electromagnetic gauge symmetry appears only after electroweak symmetry breaking, when B_μ mixes with the neutral weak gauge field W_μ^3 to form the photon and the Z boson.

2.1.5 Covariant derivative and gauge field strengths

Combining the three gauge symmetries, the covariant derivative acting on a generic SM field can be written as

$$D_\mu = \partial_\mu - ig_s G_\mu^a T_C^a - ig W_\mu^I T_L^I - ig' Y B_\mu. \quad (2.25)$$

Here g_s , g , and g' are the gauge couplings of $SU(3)_C$, $SU(2)_L$, and $U(1)_Y$, respectively. The matrices T_C^a and T_L^I are the generators acting on the particular field. For a colour triplet, $T_C^a = \lambda^a/2$, while for a colour singlet $T_C^a = 0$. For an $SU(2)_L$ doublet, $T_L^I = \tau^I/2$, while

for an $SU(2)_L$ singlet $T_L^I = 0$. The value of Y is the hypercharge of the field, as listed in Table 2.1.

The gauge kinetic terms are constructed from the field-strength tensors:

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f^{abc} G_\mu^b G_\nu^c, \quad (2.26)$$

$$W_{\mu\nu}^I = \partial_\mu W_\nu^I - \partial_\nu W_\mu^I + g \epsilon^{IJK} W_\mu^J W_\nu^K, \quad (2.27)$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (2.28)$$

The tensors $G_{\mu\nu}^a$ and $W_{\mu\nu}^I$ contain non-linear terms because $SU(3)_C$ and $SU(2)_L$ are non-Abelian groups. By contrast, $B_{\mu\nu}$ has the Abelian form because $U(1)_Y$ has only one generator.

2.1.6 The Standard Model Lagrangian

The SM Lagrangian is constructed from the field content described above and is invariant under the local gauge transformations of $SU(3)_C \times SU(2)_L \times U(1)_Y$. Using the covariant derivative and field-strength tensors introduced in the previous subsections, the minimal SM Lagrangian can be written as the sum of gauge-boson kinetic terms, fermion kinetic terms, the Higgs-sector terms, and Yukawa interactions. Ignoring gauge-fixing terms, ghost terms, and the QCD θ -term, it is written schematically as

$$\mathcal{L}_{\text{SM}} = \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{fermion}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{Yukawa}}. \quad (2.29)$$

The four terms in Eq. (2.29) are fixed by Lorentz invariance, local gauge invariance, and renormalisability, once the SM field content is specified [5, 6].

The gauge-boson kinetic terms are

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4} W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu}. \quad (2.30)$$

Here $G_{\mu\nu}^a$, $W_{\mu\nu}^I$, and $B_{\mu\nu}$ are the field-strength tensors of $SU(3)_C$, $SU(2)_L$, and $U(1)_Y$, respectively. The indices $a = 1, \dots, 8$ and $I = 1, 2, 3$ label the generators of $SU(3)_C$ and $SU(2)_L$, and repeated Lorentz and gauge indices are summed over. The common factor $-1/4$ gives the conventional canonical normalisation of the gauge-boson kinetic terms. With this normalisation, the quadratic parts of Eq. (2.30) describe the free propagation of the gluon, weak, and hypercharge gauge fields, while the non-Abelian parts of $G_{\mu\nu}^a$ and $W_{\mu\nu}^I$ generate self-interactions among the corresponding gauge bosons.

The fermion kinetic terms are

$$\mathcal{L}_{\text{fermion}} = \sum_{i=1}^3 \left[\bar{Q}_L^i i\gamma^\mu D_\mu Q_L^i + \bar{u}_R^i i\gamma^\mu D_\mu u_R^i + \bar{d}_R^i i\gamma^\mu D_\mu d_R^i + \bar{L}_L^i i\gamma^\mu D_\mu L_L^i + \bar{e}_R^i i\gamma^\mu D_\mu e_R^i \right]. \quad (2.31)$$

The sum runs over the three fermion generations. The fields Q_L^i and L_L^i are the left-handed quark and lepton doublets, while u_R^i , d_R^i , and e_R^i are right-handed singlets under $SU(2)_L$. The Dirac adjoint is defined as $\bar{\psi} = \psi^\dagger \gamma^0$, and γ^μ are the Dirac gamma matrices. The covariant derivative D_μ contains the gauge fields and couplings appropriate to the gauge quantum numbers of each fermion field. Therefore, Eq. (2.31) describes both the free propagation of fermions and their gauge interactions.

The Higgs-sector terms are

$$\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi), \quad V(\Phi) = -\mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2. \quad (2.32)$$

The first term is the kinetic term of the Higgs doublet and also contains the gauge interactions of the Higgs field through the covariant derivative. The function $V(\Phi)$ is the Higgs potential. With the convention used here, $\mu^2 > 0$ and $\lambda > 0$. The condition $\lambda > 0$ ensures that the potential is bounded from below, while the negative quadratic term causes the minimum of the potential to occur at a non-zero value of $\Phi^\dagger \Phi$. This leads to electroweak symmetry breaking through the Brout–Englert–Higgs mechanism [7–9].

The Yukawa interactions are

$$\mathcal{L}_{\text{Yukawa}} = -y_d^{ij} \bar{Q}_L^i \Phi d_R^j - y_u^{ij} \bar{Q}_L^i \tilde{\Phi} u_R^j - y_e^{ij} \bar{L}_L^i \Phi e_R^j + \text{h.c.}, \quad (2.33)$$

where $i, j = 1, 2, 3$ are generation indices, and y_u^{ij} , y_d^{ij} , and y_e^{ij} are complex Yukawa coupling matrices in flavour space. h.c. denotes the Hermitian conjugate. The conjugate Higgs doublet is defined as

$$\tilde{\Phi} = i\tau^2 \Phi^*, \quad (2.34)$$

where τ^2 is the second Pauli matrix. The field $\tilde{\Phi}$ transforms as an $SU(2)_L$ doublet with hypercharge opposite to that of Φ , allowing a gauge-invariant Yukawa interaction for up-type quarks. After electroweak symmetry breaking, the Yukawa interactions generate masses for the charged fermions. In the minimal SM, no right-handed neutrino field is included, and therefore no renormalisable neutrino Yukawa mass term appears in Eq. (2.33).

2.1.7 Electroweak symmetry breaking and the Higgs mechanism

The Higgs potential in Eq. (2.32) has a minimum at a non-zero value of $\Phi^\dagger \Phi$. For the convention used in Eq. (2.32), the minimum satisfies $\Phi^\dagger \Phi = v^2/2$, where $v = \sqrt{\mu^2/\lambda}$ is the Higgs vacuum expectation value and sets the electroweak scale. A convenient choice of vacuum expectation value is

$$\langle \Phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}. \quad (2.35)$$

The value of v is related to the Fermi constant by $v = (\sqrt{2}G_F)^{-1/2} \simeq 246 \text{ GeV}$ [10].

This vacuum breaks the electroweak symmetry according to

$$SU(2)_L \times U(1)_Y \rightarrow U(1)_{\text{em}}. \quad (2.36)$$

The unbroken symmetry is electromagnetism. This mass-generation mechanism for gauge bosons is the Brout–Englert–Higgs mechanism [7–9].

In unitary gauge, the Higgs doublet can be written as

$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}, \quad (2.37)$$

where $h(x)$ is the physical Higgs boson field. Expanding the Higgs potential around the vacuum gives

$$V(\Phi) = V(\langle \Phi \rangle) + \frac{1}{2}m_h^2 h^2 + \lambda v h^3 + \frac{\lambda}{4}h^4, \quad (2.38)$$

where the physical Higgs-boson mass is

$$m_h^2 = 2\lambda v^2 = 2\mu^2. \quad (2.39)$$

Unlike the W and Z boson masses, which are fixed at tree level by the electroweak gauge couplings and the vacuum expectation value, the Higgs-boson mass depends on the quartic coupling λ . Therefore, the SM does not predict the numerical value of m_h before λ is fixed experimentally. The Higgs boson was observed by the ATLAS and CMS experiments in 2012 [11, 12]. The observed Higgs-boson mass is approximately 125 GeV, corresponding to $\lambda \simeq m_h^2/(2v^2) \simeq 0.13$ [13].

The masses of the electroweak gauge bosons arise from the Higgs kinetic term,

$$(D_\mu \Phi)^\dagger (D^\mu \Phi), \quad (2.40)$$

after substituting the Higgs vacuum expectation value. It is useful to define the charged weak fields as

$$W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \mp iW_\mu^2). \quad (2.41)$$

The terms quadratic in the electroweak gauge fields are then

$$(D_\mu \langle \Phi \rangle)^\dagger (D^\mu \langle \Phi \rangle) = \frac{g^2 v^2}{4} W_\mu^+ W^{-\mu} + \frac{v^2}{8} (gW_\mu^3 - g'B_\mu)^2. \quad (2.42)$$

The first term is the mass term of the charged weak bosons. The second term shows that the neutral gauge fields W_μ^3 and B_μ mix and must be diagonalised.

The weak mixing angle is defined by

$$\cos \theta_W = \frac{g}{\sqrt{g^2 + g'^2}}, \quad \sin \theta_W = \frac{g'}{\sqrt{g^2 + g'^2}}. \quad (2.43)$$

The corresponding neutral mass eigenstates are

$$A_\mu = B_\mu \cos \theta_W + W_\mu^3 \sin \theta_W, \quad (2.44)$$

$$Z_\mu = -B_\mu \sin \theta_W + W_\mu^3 \cos \theta_W. \quad (2.45)$$

The field A_μ is identified with the photon, while Z_μ is the neutral weak boson. The photon remains massless because it is associated with the unbroken gauge symmetry $U(1)_{\text{em}}$.

The resulting gauge-boson masses are

$$m_W = \frac{1}{2} g v, \quad m_Z = \frac{1}{2} \sqrt{g^2 + g'^2} v = \frac{m_W}{\cos \theta_W}, \quad m_\gamma = 0. \quad (2.46)$$

Thus, the SM predicts the relation $m_W = m_Z \cos \theta_W$. Inserting $v \simeq 246$ GeV and the electroweak couplings gives $m_W \simeq 80$ GeV and $m_Z \simeq 91$ GeV. Therefore, the Higgs mechanism gives masses to the W and Z bosons while leaving the photon massless.

After electroweak symmetry breaking, the unbroken gauge symmetry is $U(1)_{\text{em}}$, whose gauge field is the photon field A_μ . At energies much below the electroweak scale, the electromagnetic interactions of charged fermions are described by quantum electrodynamics (QED).

480 For a charged fermion field ψ with electric charge Q , the QED Lagrangian can be written as

$$481 \quad \mathcal{L}_{\text{QED}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}(i\gamma^\mu D_\mu - m)\psi, \quad (2.47)$$

482 where

$$483 \quad D_\mu = \partial_\mu - ieQA_\mu, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (2.48)$$

484 Here e is the electromagnetic coupling constant, Q is the electric charge of the fermion in
485 units of e , and $F_{\mu\nu}$ is the electromagnetic field-strength tensor. Thus, QED appears as the
486 low-energy gauge theory associated with the unbroken $U(1)_{\text{em}}$ symmetry of the SM.

487 2.1.8 Fermion masses and flavour mixing

488 The Yukawa interactions generate fermion masses after electroweak symmetry breaking.
489 When the Higgs field acquires its vacuum expectation value v , the Yukawa coupling matrices
490 are converted into fermion mass matrices:

$$491 \quad M_u^{ij} = \frac{v}{\sqrt{2}}y_u^{ij}, \quad M_d^{ij} = \frac{v}{\sqrt{2}}y_d^{ij}, \quad M_e^{ij} = \frac{v}{\sqrt{2}}y_e^{ij}. \quad (2.49)$$

492 Here M_u , M_d , and M_e are the mass matrices of the up-type quarks, down-type quarks, and
493 charged leptons, respectively. The matrices y_u , y_d , and y_e are the corresponding dimensionless
494 Yukawa coupling matrices, and the indices $i, j = 1, 2, 3$ label the fermion generations.

495 The physical fermion masses are obtained by diagonalising the mass matrices. In the
496 quark sector, the up-type and down-type mass matrices are diagonalised by different rotations
497 in flavour space. As a result, flavour mixing appears in the charged-current weak interaction.
498 This mixing is described by the Cabibbo–Kobayashi–Maskawa (CKM) matrix [14, 15]. In
499 the mass eigenstate basis, the charged-current interaction of quarks is

$$500 \quad \mathcal{L}_{\text{CC}} = \frac{g}{\sqrt{2}}\bar{u}_L^i \gamma^\mu V_{ij}^{\text{CKM}} d_L^j W_\mu^+ + \text{h.c.} \quad (2.50)$$

501 Here g is the $SU(2)_L$ gauge coupling, u_L^i and d_L^j are the left-handed up-type and down-type
502 quark mass eigenstates, and W_μ^+ is the charged weak gauge boson. The matrix element
503 V_{ij}^{CKM} describes the strength of the charged-current transition from the down-type quark d^j
504 to the up-type quark u^i . The matrices γ^μ are the Dirac gamma matrices, and the Hermitian-
505 conjugate term describes the corresponding interaction involving W_μ^- . The CKM matrix is

essential for describing flavour-changing charged-current processes such as $b \rightarrow c\tau\nu$. Its complex phase also provides a source of CP violation in the quark sector of the SM.

In the minimal SM, neutrino masses are absent because no right-handed neutrino fields are included and no renormalisable neutrino Yukawa interaction can be written with the minimal field content. This provides one of the motivations for considering extensions of the SM.

2.2 Beyond the Standard Model

Although the SM has been tested with high precision and has successfully described a wide range of experimental observations, it is not a complete theory of nature. Several experimental and theoretical facts point to physics beyond the SM.

2.2.1 Gravity and the Planck scale

The SM describes the electromagnetic, weak, and strong interactions as quantum gauge field theories, but it does not include gravity. Gravity is instead described at macroscopic scales by general relativity. A complete theory of nature is therefore expected to provide a consistent quantum description of gravity and to explain its connection to the SM interactions.

The difficulty of connecting gravity with the SM appears in two related ways. First, quantum gravitational effects are expected to become important at the Planck scale, $M_{\text{Pl}} \sim 10^{19}$ GeV. This is much higher than the electroweak scale, $v \simeq 246$ GeV. This large separation of scales is closely related to the hierarchy problem discussed below.

Second, constructing a quantum theory of gravity is theoretically challenging. If general relativity is quantised perturbatively in the same way as ordinary quantum field theories, it is not renormalisable as a fundamental theory. This makes it difficult to combine gravity with the quantum-field-theoretic framework of the SM in a straightforward way. Several possible approaches to quantum gravity have been proposed, including string theory, loop quantum gravity, and asymptotic safety, although no complete theory has been experimentally confirmed.

2.2.2 The hierarchy problem

The electroweak scale is set by the Higgs vacuum expectation value, $v \simeq 246$ GeV, and is closely related to the masses of the weak gauge bosons and the Higgs boson. In the SM, the

Higgs-boson mass is determined at tree level by

$$m_h^2 = 2\lambda v^2, \quad (2.51)$$

where λ is the Higgs quartic coupling. The observed value $m_h \simeq 125$ GeV therefore corresponds to $\lambda \simeq 0.13$.

The hierarchy problem arises because the Higgs mass parameter is highly sensitive to physics at very high energy scales. If the SM is treated as an effective theory valid up to an ultraviolet cutoff scale Λ , loop corrections to the Higgs mass can be written schematically as

$$m_{h,\text{phys}}^2 = m_{h,0}^2 + \delta m_h^2, \quad (2.52)$$

where $m_{h,0}^2$ is the bare Higgs mass parameter and δm_h^2 denotes the quantum correction. At one-loop order, the correction contains terms quadratically sensitive to the cutoff scale,

$$\delta m_h^2 \sim \frac{\Lambda^2}{16\pi^2}, \quad (2.53)$$

If the cutoff scale is close to the Planck scale, $\Lambda \sim M_{\text{Pl}} \sim 10^{19}$ GeV, the magnitude of δm_h^2 is many orders of magnitude larger than the observed value $m_h^2 \sim (125 \text{ GeV})^2$. Therefore, the bare parameter $m_{h,0}^2$ and the quantum correction δm_h^2 must cancel each other with extremely high precision in order to leave the observed electroweak-scale Higgs mass. This required cancellation is known as fine-tuning.

The hierarchy problem is the question of why the electroweak scale remains so much smaller than a possible ultraviolet scale such as the Planck scale. It motivates many extensions of the SM, such as supersymmetry, composite Higgs models, and other scenarios in which new physics appears near the TeV scale and stabilises the Higgs mass against large ultraviolet corrections. More generally, the hierarchy problem provides one of the motivations for experimental searches for new particles at the TeV scale.

2.2.3 Neutrino masses

In the minimal SM, neutrinos are massless because no right-handed neutrino fields are included and no renormalisable neutrino mass term can be written using only the SM field content. However, neutrino oscillation experiments have shown that neutrinos change flavour as they propagate, which requires non-zero neutrino mass differences. Evidence for atmospheric neutrino oscillations was reported by the Super-Kamiokande experiment [16], and solar neutrino flavour transformation was established by the Sudbury Neutrino Obser-

vatory [17]. Therefore, neutrino masses provide direct experimental evidence for physics beyond the minimal SM.

2.2.4 Dark matter

Astrophysical and cosmological observations indicate the existence of non-luminous matter that cannot be explained by the known particles of the SM. Measurements of the cosmic microwave background, for example by the Planck satellite, are well described by a cosmological model containing a significant non-baryonic dark matter component [18]. The SM does not contain a suitable cold dark matter candidate. Although neutrinos are electrically neutral and weakly interacting, their small masses and relativistic behaviour in the early universe do not provide the required cold dark matter component.

2.2.5 Matter–antimatter asymmetry

The observable universe contains much more matter than antimatter. Sakharov identified three necessary conditions for dynamically generating such a baryon asymmetry: baryon-number violation, C and CP violation, and departure from thermal equilibrium [19]. The SM contains CP violation through the CKM matrix and baryon-plus-lepton number violation through non-perturbative electroweak effects. However, the amount of CP violation in the SM and the nature of the electroweak phase transition are not sufficient to explain the observed baryon asymmetry of the universe [20, 21]. This motivates new sources of CP violation and new dynamics in the early universe.

2.2.6 The strong CP problem

The most general QCD Lagrangian allows a CP-violating term of the form

$$\mathcal{L}_\theta = \theta \frac{g_s^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}, \quad \tilde{G}^{a\mu\nu} = \frac{1}{2} \varepsilon^{\mu\nu\rho\sigma} G_{\rho\sigma}^a. \quad (2.54)$$

Here g_s is the strong coupling constant, and $G_{\mu\nu}^a$ is the gluon field-strength tensor of the $SU(3)_C$ gauge group, with the colour index $a = 1, \dots, 8$ labelling the eight gluon fields. The tensor $\tilde{G}^{a\mu\nu}$ is the dual field-strength tensor, defined using the totally antisymmetric Levi-Civita tensor $\varepsilon^{\mu\nu\rho\sigma}$. The parameter θ is a dimensionless QCD vacuum angle that controls the size of this CP-violating interaction. The factor $g_s^2/(32\pi^2)$ is a conventional normalisation.

The operator $G_{\mu\nu}^a \tilde{G}^{a\mu\nu}$ is gauge invariant and has mass dimension four, so it is allowed by the gauge symmetry and renormalisability of QCD. However, it violates CP symmetry. Therefore, a non-zero value of the effective QCD vacuum angle would lead to CP violation in the strong interaction.

This term would contribute to the electric dipole moment of the neutron. Experimental limits on the neutron electric dipole moment strongly constrain the effective QCD vacuum angle $\bar{\theta}$ [22]. The effective parameter $\bar{\theta}$ includes both the QCD vacuum angle θ and phases from the quark mass matrix. The absence of observable strong CP violation, despite the fact that such a term is allowed by the SM gauge symmetries, is known as the strong CP problem.

2.2.7 Flavour anomalies and lepton flavour universality

The flavour structure of the SM is mainly determined by the Yukawa matrices. After electroweak symmetry breaking, these matrices generate the fermion mass matrices; their eigenvalues determine the fermion masses, while the mismatch between the diagonalisation of the up- and down-type quark mass matrices gives rise to quark flavour mixing through the CKM matrix.

Although the SM accommodates the observed masses and mixing angles, it does not explain their hierarchical pattern or their origin. Precision measurements in flavour physics therefore provide sensitive probes of possible physics beyond the SM.

A key property of the SM is lepton flavour universality (LFU), which means that the electroweak gauge interactions couple equally to the three charged-lepton generations, apart from effects due to their different masses. Semileptonic B -meson decays provide precise tests of LFU by comparing decay rates involving different charged leptons. In charged-current transitions of the form $b \rightarrow c\ell\nu$, the commonly studied ratios are

$$R(D^{(*)}) = \frac{\mathcal{B}(\bar{B} \rightarrow D^{(*)} \tau \bar{\nu}_\tau)}{\mathcal{B}(\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}_\ell)}, \quad \ell = e, \mu. \quad (2.55)$$

These observables compare decays involving a τ lepton with the corresponding decays involving light charged leptons.

Neutral-current transitions of the form $b \rightarrow s\ell^+\ell^-$ can be tested through ratios such as

$$R(K^{(*)}) = \frac{\mathcal{B}(B \rightarrow K^{(*)} \mu^+ \mu^-)}{\mathcal{B}(B \rightarrow K^{(*)} e^+ e^-)}. \quad (2.56)$$

These ratios compare final states with muon pairs and electron pairs. Recent measurements of $R(K)$ and $R(K^*)$ are compatible with SM predictions [23, 24]. However, measurements of

$R(D)$ and $R(D^*)$ continue to show tensions with SM predictions. The HFLAV CKM 2025 average gives a combined discrepancy of about 3.8σ with respect to the SM prediction [25] as shown in Figure 2.1. These observables remain important motivations for models that couple differently to different lepton generations.




Fig. 2.1 Average of $R(D^*)$ and $R(D)$ [25] measured by the BaBar [26, 27], Belle [28–31], Belle-II [32, 33], and LHCb [34–36] experiments.

2.3 Possible Solutions to the Remaining Problems of the Standard Model

Many extensions of the SM have been proposed to address one or more of the problems described above. This section briefly introduces several representative ideas: supersymmetry, axions, and leptiquarks. These models are not mutually exclusive, and in many cases they address different aspects of the SM's limitations.

2.3.1 Supersymmetry

Supersymmetry (SUSY) is a spacetime symmetry relating bosons and fermions [37, 38]. In supersymmetric theories, each SM particle is accompanied by a superpartner whose spin differs by one half unit. The fermionic superpartners of bosons are conventionally named by adding the suffix '-ino', such as the gluino, wino, bino, and higgsino, while the bosonic superpartners of fermions are named by adding the prefix 's-', such as squarks and sleptons.

If supersymmetry were exact, particles and their superpartners would have equal masses. Since no such superpartners have been observed, SUSY must be broken if it is realised in nature.

One of the main motivations for SUSY is the hierarchy problem. Loop corrections to the Higgs mass from SM particles can be cancelled by corresponding contributions from their superpartners, reducing the sensitivity of the electroweak scale to very high energy scales [39]. In models with conserved R -parity, supersymmetric particles are produced in pairs and the lightest supersymmetric particle (LSP) is stable [40]. If the LSP is electrically neutral and weakly interacting, such as a neutralino in many SUSY scenarios, it can provide a dark matter candidate [41, 42]. SUSY can also improve the unification of gauge couplings at high energy, since the additional superpartners modify the renormalisation-group evolution of the gauge couplings [43, 44]. However, no direct evidence for supersymmetric particles has yet been observed, and the simplest low-scale SUSY scenarios are increasingly constrained by collider searches [45].

2.3.2 Axions and the strong CP problem

The Peccei–Quinn mechanism provides a dynamical solution to the strong CP problem [46, 47]. It introduces a new global chiral symmetry, usually denoted as the Peccei–Quinn symmetry $U(1)_{\text{PQ}}$, which is spontaneously broken. The associated pseudo-Nambu–Goldstone boson is the axion [48, 49]. The axion field dynamically relaxes the effective value of $\bar{\theta}$ to zero, thereby explaining the absence of observable strong CP violation. In many models, axions or axion-like particles can also contribute to the dark matter density of the universe [50].

2.3.3 Leptoquarks

Leptoquarks provide another well-motivated class of particles beyond the Standard Model. They are hypothetical bosons that couple directly to a quark and a lepton, thereby connecting the quark and lepton sectors. Such particles appear naturally in several extensions of the SM, including models based on quark–lepton unification, such as the Pati–Salam framework [51].

Leptoquarks are particularly interesting because they can affect semileptonic processes. This makes them relevant to the flavour anomalies observed in several B -meson decay measurements, where the experimental results have shown tensions with the SM predictions in observables testing lepton-flavour universality.

Among the possible leptoquark scenarios, the vector singlet leptoquark U_1 is especially well motivated for charged-current anomalies such as $b \rightarrow c \tau \nu$. This motivates collider

670 searches for final states containing hadronically decaying τ leptons. The following chapter
671 introduces the leptoquark models in more detail and the leptoquark searches at the LHC.

Chapter 3

Leptoquark Search at the LHC

3.1 Leptoquark Models and Phenomenology

3.1.1 General properties of leptoquarks

Leptoquarks (LQs) are hypothetical colour-charged bosons that couple directly to a quark and a lepton and therefore mediate interactions connecting the quark and lepton sectors. Depending on their spin, leptoquarks can be classified as scalar leptoquarks with spin zero or vector leptoquarks with spin one. They appear in several theoretical frameworks, including grand unified theories such as $SU(5)$ and $SO(10)$, quark–lepton unification models such as the Pati–Salam model, composite models, and models motivated by flavour anomalies [52, 51, 53].

Leptoquarks are particularly interesting in the context of semileptonic flavour physics. Their tree-level exchange can generate effective four-fermion interactions involving two quark fields and two lepton fields. Such interactions can modify flavour observables in processes such as $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$. In addition, the flavour structure of LQ couplings is model dependent and can be arranged to favour interactions with third-generation fermions. This feature is motivated by the distinctive role of the top quark, bottom quark, and τ -lepton in many flavour-motivated scenarios.

At the LHC, leptoquarks can be searched for directly. Since they carry colour charge, they can be pair-produced through QCD interactions. Additional production modes, such as single production or non-resonant exchange, are also possible, depending on the LQ coupling to SM quarks and leptons. Therefore, the LQ searches in ATLAS provide information complementary to low-energy flavour measurements: flavour observables probe virtual effects through precision measurements, while the search at pp collider probes direct LQ production or high-energy virtual exchange.

3.1.2 The simplified U_1 leptoquark model

Several leptoquark representations have been studied in connection with semileptonic B -meson anomalies, including the scalar singlet S_1 and the scalar doublet R_2 . In this thesis, however, the benchmark interpretation is based on the vector singlet leptoquark U_1 . The U_1 benchmark is well motivated because its left-handed coupling to quark and lepton doublets can generate the SM-like vector operator for $b \rightarrow c\tau\nu$. In addition, flavour structures with enhanced couplings to third-generation fermions naturally lead to final states involving τ -leptons, neutrinos, and jets.

The U_1 leptoquark transforms as $U_1^\mu \sim (3, 1, 2/3)$ under the SM gauge group. It is therefore a colour-triplet, weak-singlet vector boson with hypercharge $Y = 2/3$. Since it is an $SU(2)$ singlet, its electric charge is $Q = T^3 + Y = 2/3$.

A commonly used simplified-model Lagrangian [54] for U_1 is

$$\begin{aligned} \mathcal{L}_U = & -\frac{1}{2}U_{1\mu\nu}^\dagger U_1^{\mu\nu} + M_U^2 U_{1\mu}^\dagger U_1^\mu \\ & -ig_c(1 - \kappa_c)U_{1\mu}^\dagger T^a U_{1\nu} G^{a\mu\nu} - i\frac{2}{3}g_Y(1 - \kappa_Y)U_{1\mu}^\dagger U_{1\nu} B^{\mu\nu} \\ & + (U_1^\mu J_\mu + \text{h.c.}). \end{aligned} \quad (3.1)$$

Here U_1^μ is the vector leptoquark field and M_U is the leptoquark mass. The first term is the kinetic term, while the second term is the mass term. The field-strength tensor of U_1 is defined as

$$U_{1\mu\nu} = D_\mu U_{1\nu} - D_\nu U_{1\mu}, \quad (3.2)$$

with the covariant derivative

$$D_\mu = \partial_\mu - ig_c G_\mu^a T^a - i\frac{2}{3}g_Y B_\mu. \quad (3.3)$$

Here g_c and g_Y are the colour and hypercharge gauge couplings. The fields G_μ^a and B_μ are the gluon and hypercharge gauge fields, and $G^{a\mu\nu}$ and $B^{\mu\nu}$ are their corresponding field-strength tensors. The matrices T^a are the generators of $SU(3)$. For a colour-triplet field, they are given by $T^a = \lambda^a/2$, where λ^a are the Gell-Mann matrices and $a = 1, \dots, 8$.

The parameters κ_c and κ_Y control the interactions of the vector leptoquark with the SM gauge fields. $\kappa_c = \kappa_Y = 0$ corresponds to the Yang–Mills scenario, whereas $\kappa_c = \kappa_Y = 1$ corresponds to the minimal-coupling scenario [55]. The Yang–Mills scenario is adopted throughout this thesis.

The last term in Eq. (3.1) describes the interaction of U_1 with SM fermions through the current J_μ . It is given by

$$J_\mu = \frac{g_U}{\sqrt{2}} [\beta_L^{i\alpha} (\bar{q}_L^i \gamma_\mu \ell_L^\alpha) + \beta_R^{i\alpha} (\bar{d}_R^i \gamma_\mu e_R^\alpha)]. \quad (3.4)$$

Here $i = 1, 2, 3$ labels the quark generation and $\alpha = 1, 2, 3$ labels the lepton generation. q_L^i and ℓ_L^α are left-handed quark and lepton doublets, while d_R^i and e_R^α are right-handed down-type quark and charged-lepton singlets. The parameter g_U denotes the overall strength of the U_1 coupling to fermions. The matrices $\beta_L^{i\alpha}$ and $\beta_R^{i\alpha}$ are complex 3×3 flavour-coupling matrices for the left- and right-handed interactions, respectively. The strength of a given flavour coupling is therefore proportional to $g_U \beta_{L,R}^{i\alpha}$.

The down-quark and charged-lepton mass eigenstate basis is adopted for the $SU(2)$ multiplets:

$$q_L^i = \begin{pmatrix} V_{ji}^* u_L^j \\ d_L^i \end{pmatrix}, \quad \ell_L^\alpha = \begin{pmatrix} \nu_L^\alpha \\ e_L^\alpha \end{pmatrix}, \quad (3.5)$$

where V is the CKM matrix.

Expanding the left-handed current from Eq. 3.4 gives

$$\bar{q}_L^i \gamma_\mu \ell_L^\alpha = V_{ji} \bar{u}_L^j \gamma_\mu \nu_L^\alpha + \bar{d}_L^i \gamma_\mu e_L^\alpha. \quad (3.6)$$

This expression shows that a single left-handed flavour coupling $\beta_L^{i\alpha}$ gives two related interactions. One involves an up-type quark together with a neutrino, while the other involves a down-type quark together with a charged lepton.

The simplified U_1 model allows general flavour couplings to different quark and lepton generations. Meanwhile, we focus on the charged-current $b \rightarrow c \tau \nu$ transition and the signatures containing a τ -lepton and a neutrino. Therefore, the couplings to second-generation leptons in our benchmark are set to zero, which is $\beta_L^{22} = \beta_L^{32} = 0$, and only the couplings involving third-generation leptons are retained. Finally, the flavour-coupling matrices are taken to be

$$\beta_L = \begin{pmatrix} 0 & 0 & \beta_L^{13} \\ 0 & 0 & \beta_L^{23} \\ 0 & 0 & \beta_L^{33} \end{pmatrix}, \quad \beta_R = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \beta_R^{33} \end{pmatrix}. \quad (3.7)$$

The third-generation coupling is normalised to $\beta_L^{33} = 1$, while g_U remains a free parameter controlling the overall fermionic coupling strength.

Within this benchmark, β_L^{23} controls the $s\tau$ and $c\nu_\tau$ coupling through the CKM factor V_{cs} , while β_L^{33} controls the $b\tau$ coupling and contributes $c\nu_\tau$ coupling as $V_{cb}\beta_L^{33}$.

The heavy U_1 field can be integrated out at the matching scale $\mu \sim M_U$. At tree level, its exchange generates four-fermion operators involving left- and right-handed quark and lepton fields [56]. The corresponding effective Lagrangian can be written as

$$\mathcal{L}_{\text{eff}}^U = -\frac{2}{v^2} \left[C_{LL}^{ij\alpha\beta} \mathcal{O}_{LL}^{ij\alpha\beta} + C_{RR}^{ij\alpha\beta} \mathcal{O}_{RR}^{ij\alpha\beta} + \left(C_{LR}^{ij\alpha\beta} \mathcal{O}_{LR}^{ij\alpha\beta} + \text{h.c.} \right) \right]. \quad (3.8)$$

The four-fermion operators are defined as

$$\mathcal{O}_{LL}^{ij\alpha\beta} = (\bar{q}_L^i \gamma_\mu \ell_L^\alpha) (\bar{\ell}_L^\beta \gamma^\mu q_L^j), \quad (3.9)$$

$$\mathcal{O}_{RR}^{ij\alpha\beta} = (\bar{d}_R^i \gamma_\mu e_R^\alpha) (\bar{e}_R^\beta \gamma^\mu d_R^j), \quad (3.10)$$

$$\mathcal{O}_{LR}^{ij\alpha\beta} = (\bar{q}_L^i \gamma_\mu \ell_L^\alpha) (\bar{e}_R^\beta \gamma^\mu d_R^j). \quad (3.11)$$

The Wilson coefficients are

$$C_{LL}^{ij\alpha\beta} = C_U \beta_L^{i\alpha} (\beta_L^{j\beta})^*, \quad (3.12)$$

$$C_{RR}^{ij\alpha\beta} = C_U \beta_R^{i\alpha} (\beta_R^{j\beta})^*, \quad (3.13)$$

$$C_{LR}^{ij\alpha\beta} = C_U \beta_L^{i\alpha} (\beta_R^{j\beta})^*. \quad (3.14)$$

The overall coefficient is defined as

$$C_U = \frac{g_U^2 v^2}{4M_U^2}, \quad (3.15)$$

where $v \simeq 246$ GeV is the Higgs vacuum expectation value. The coefficient C_U sets the overall strength of the effective interaction and scales as g_U^2/M_U^2 , while the flavour dependence is encoded in the coupling matrices β_L and β_R .

The effective Lagrangian for the $b \rightarrow c\tau\nu$ transition can be written as:

$$\begin{aligned} \mathcal{L}_{\text{eff}}^{b \rightarrow c\tau\nu} = & -\frac{4G_F}{\sqrt{2}} V_{cb} \left[(1 + C_{LL}^c) (\bar{c}_L \gamma_\mu b_L) (\bar{\tau}_L \gamma^\mu \nu_{\tau L}) \right. \\ & \left. - 2C_{LR}^c (\bar{c}_L b_R) (\bar{\tau}_R \nu_{\tau L}) \right], \end{aligned} \quad (3.16)$$

where G_F is the Fermi constant. The coefficient C_{LL}^c modifies the SM-like left-handed vector interaction, while C_{LR}^c is associated with the scalar operator arising from the combination of left- and right-handed fermion couplings of the U_1 leptoquark.

For the flavour structure introduced above, the corresponding coefficients are

$$C_{LL}^c = C_U \frac{(\beta_L^{33})^*}{V_{cb}} (V_{cd}\beta_L^{13} + V_{cs}\beta_L^{23} + V_{cb}\beta_L^{33}), \quad (3.17)$$

$$C_{LR}^c = C_U \frac{(\beta_R^{33})^*}{V_{cb}} (V_{cd}\beta_L^{13} + V_{cs}\beta_L^{23} + V_{cb}\beta_L^{33}), \quad (3.18)$$

Using the normalisation $\beta_L^{33} = 1$ and neglecting the small first-generation coupling β_L^{13} , the coefficients reduce to¹

$$C_{LL}^c \simeq C_U \left(1 + \frac{V_{cs}}{V_{cb}} \beta_L^{23} \right), \quad (3.19)$$

$$C_{LR}^c \simeq C_U (\beta_R^{33})^* \left(1 + \frac{V_{cs}}{V_{cb}} \beta_L^{23} \right). \quad (3.20)$$

3.1.3 Low-energy fit and benchmark scenarios

The Wilson coefficients C_{LL}^c and C_{LR}^c introduced above can be constrained using measurements of semileptonic charged-current B -meson decays. Ref. [56] performed a combined fit to low-energy $b \rightarrow c \tau \nu$ observables and determines the preferred region in the (C_{LL}^c, C_{LR}^c) plane. The fit result is shown in Fig. 3.1. The Wilson coefficients in this figure are evolved to a common reference scale of 1 TeV to facilitate comparison with high-energy observables. The best-fit point is $(C_{LL}^c, C_{LR}^c) = (0.05, -0.02)$, and the preferred region is compatible with both pure-left-handed ($\beta_R^{33} = 0$) and sizeable right-handed ($\beta_R^{33} = -1$) scenarios. Although the sign of β_R^{33} affects the low-energy relation between C_{LL}^c and C_{LR}^c , the collider predictions used in this thesis are identical for $\beta_R^{33} = +1$ and -1 . The new-physics amplitudes with different fermion chiralities do not interfere, and the amplitude proportional to a product of left- and right-handed couplings does not interfere with the SM amplitude. The collider interpretation therefore depends only on $|\beta_R^{33}|$.

Pure-left-handed scenario. For $\beta_R^{33} = 0$, Eq. 3.20 gives $C_{LR}^c = 0$. Therefore, the U_1 leptoquark contribution to $b \rightarrow c \tau \nu$ is described only by the left-handed vector operator, as given by Eq. 3.19. This is the primary benchmark scenario considered in this thesis.

¹The small first-generation coupling $\beta_L^{13} = (V_{td}^*/V_{ts}^*)\beta_L^{23} \sim 0.2\beta_L^{23}$.




Fig. 3.1 The result in (C_{LL}^c, C_{LR}^c) plane obtained from the fit to low-energy $b \rightarrow c \tau \nu$ observables. The blue solid ellipses correspond to the 1, 2 and 3 σ contours. The blue dotted lines are obtained including also the constraint from $\mathcal{B}(B_u^- \rightarrow \tau \bar{\nu})$. The black point indicates the best-fit point of the fit, $(C_{LL}^c, C_{LR}^c) = (0.05, -0.02)$. The coloured bands show the $\Delta\chi^2 = 1$ regions preferred by the individual observables. The pure-left-handed scenario ($\beta_R^{33} = 0$) and the equal magnitude right-handed scenario ($\beta_R^{33} = -1$) are given by the black solid lines [56].

The corresponding B -anomaly-favoured region can be obtained from the intersection of the low-energy fit with the line $C_{LR}^c = 0$ in Fig. 3.1.

Right-handed scenario. The second benchmark is obtained by allowing a sizeable right-handed third-generation coupling. For $\beta_R^{33} = -1$, the relation between the Wilson coefficients becomes

$$C_{LR}^c = -C_{LL}^c. \quad (3.21)$$

This benchmark therefore corresponds to the diagonal line in Fig. 3.1, where the right-handed third-generation coupling has the same magnitude as the left-handed coupling.

The two scenarios are described by the parameter space (β_L^{23}, M_U, g_U) with different fixed values of β_R^{33} . Particular emphasis is placed on β_L^{23} , since charged-current low-energy data alone do not provide a stringent constraint on this coupling. For fixed M_U and g_U , β_L^{23} has a strong impact on C_{LL}^c because its contribution is enhanced by the CKM ratio V_{cs}/V_{cb} .

Indeed, one can constrain the size of β_L^{23} by considering possible ultraviolet (UV) completions of the simplified U_1 model. While the simplified model parametrises the relevant interactions directly in terms of the couplings β_L^{ij} , the UV completion provides a more fundamental description of their origin. One example is the “4321” model, which is based on the extended gauge group $\mathcal{G}_{4321} = SU(4) \times SU(3)' \times SU(2)_L \times U(1)'$. In the flavour structure considered in this model, the U_1 couplings are non-universal, with $|\beta_L^{23}| < |\beta_L^{33}|$ [54].

Additional constraints can be introduced when considering $\Delta F = 2$ observables. These are processes in which a quark-flavour quantum number changes by two units, including neutral-meson mixing such as $B_s - \bar{B}_s$, $K - \bar{K}$, and $D - \bar{D}$ mixing. In a given UV completion, the flavour-symmetry breaking that generates non-universal U_1 couplings may also be transmitted to the neutral-current sector, inducing flavour-changing neutral currents (FCNC) and consequently additional contributions to $\Delta F = 2$ processes. The strong experimental constraints on neutral-meson mixing therefore restrict the allowed flavour-violating structure of the UV model and indirectly constrain β_L^{23} .

Combining such considerations of the UV flavour structure with constraints from $\Delta F = 2$ observables, Ref. [56] identifies

$$\beta_L^{23} \in [0.06, 0.16] \quad (3.22)$$

as a theoretically motivated range for experimental searches. **However, this constraint should not be interpreted as a model-independent experimental bound on β_L^{23} .** Additional particles, flavour symmetries, or cancellation mechanisms may suppress flavour-

changing transitions and thereby allow larger values of β_L^{23} . Therefore, large values of β_L^{23} remain a possibility within the U_1 model, and it is important to test this complementary region directly.

3.1.4 Leptoquark searches at the LHC

Leptoquarks can be searched for at the LHC through several production modes, including pair production, resonant single production, and non-resonant t -channel exchange. Pair production is dominated by QCD interactions. For a fixed κ choice, its cross section depends strongly on M_U but weakly on the couplings g_U and $\beta^{i\alpha}$. Pair-production searches therefore provide relatively robust constraints on the leptoquark mass.

ATLAS and CMS have performed searches for pair-produced leptoquarks coupling to third-generation fermions in final states containing τ -leptons, b -jets, and, depending on the decay mode, missing transverse momentum [57, 58]. For the vector U_1 , the mass limit extends to approximately 1.5–1.8 TeV as shown in Fig. 3.3. The limit also depends on the choice of the U_1 -gluon coupling, such as the Yang–Mills or minimal-coupling scenario, as well as on the decay pattern determined by the assumed fermionic couplings.

For single production, the cross section also depends strongly on the leptoquark–fermion couplings. Searches for singly produced leptoquarks decaying into $b\tau$ have been performed in the $b\tau\tau$ final state, where the leptoquark is produced in association with a τ -lepton as shown in Fig. 3.2a [57, 59]. In addition, virtual t -channel U_1 exchange can contribute to final states such as $\tau\tau$ or $\tau\nu$, enhancing the event yields in the high-mass region. ATLAS and CMS have therefore searched for non-resonant leptoquark contributions in high-mass $\tau\tau$ production [60, 61]. These searches can remain sensitive to masses beyond the reach of the direct pair-production searches, particularly for large g_U , since the sensitivity is approximately governed by g_U/M_U .




Fig. 3.2 Example Feynman diagrams of LQ production and exchange process for $\tau\tau(+b)$ searches.




Fig. 3.3 Summary of the observed and expected 95% CL constraints on the U_1 vector-leptoquark coupling $\lambda_{b\tau}$ as a function of the leptoquark mass in the Yang–Mills coupling scenario, with $\beta_R^{33} = 0$. Pair-production searches dominate the direct mass constraint at lower masses, while single-production and $\tau\tau$ searches provide coupling-dependent sensitivity at larger masses. Here $\lambda_{b\tau} = g_U/\sqrt{2}$ for $\beta_L^{33} = 1$ [62].

On the other hand, the existing $\tau\tau(+b)$ searches generally assume β_L^{23} to be small or neglect its contribution, motivated by the flavour assumptions discussed in Sec. 3.1.3. Consequently, scenarios with a sizeable $c\nu_\tau$ coupling, and in particular the large- β_L^{23} region, have not been systematically explored at the LHC.

The $\tau\nu$ final state provides a complementary probe of the $b\tau$ and $c\nu_\tau$ coupling. CMS has performed an inclusive search for events with a hadronically decaying τ -lepton and large missing transverse momentum, including an interpretation in terms of leptoquark exchange [63], but without an explicit b -jet requirement. This thesis presents a dedicated search in the $\tau\nu(+b)$ final state. Prior to the analysis, no dedicated ATLAS or CMS leptoquark search had targeted this topology. The $\tau\nu(+b)$ search presented here is therefore complementary to the existing $\tau\tau(+b)$ searches: it probes both the $b\tau$ and $c\nu_\tau$ couplings and extends the collider interpretation to large values of β_L^{23} . The corresponding benchmark and target signatures are introduced in the following subsection.

3.1.5 Signal model and target signatures

In this thesis, we focus on the simplified U_1 model introduced in Sec. 3.1.2 in the Yang–Mills scenario, corresponding to $\kappa_c = \kappa_Y = 0$ in Eq. 3.1. The third-generation coupling is normalised to $\beta_L^{33} = 1$, while the primary interpretation assumes a pure-left-handed scenario ($\beta_R^{33} = 0$). An additional interpretation including a right-handed coupling is presented with the final results. Therefore, the signal is characterised by (M_U, g_U, β_L^{23}) .

Figure 3.4 shows representative U_1 signal diagrams targeted by the analysis. Both resonant single production and non-resonant t -channel exchange are considered, and each is divided according to the presence or absence of a b -tagged jet. This results in four target signatures: resonant $\tau\nu + b$, non-resonant $\tau\nu + b$, resonant $\tau\nu + 0b$, and non-resonant $\tau\nu + 0b$.




Fig. 3.4 Signal diagrams targeted in this analysis: non-resonant t -channel exchange (left) and resonant single production (right). The final-state jet can originate from a b - or s -quark.

The b -jet requirement provides strong discrimination against the dominant SM $\tau\nu + \text{jets}$ background, since genuine SM production of a $\tau\nu$ system in association with a b -quark is CKM suppressed [64]. The $\tau\nu + b$ categories are therefore particularly sensitive to the combination of the $b\tau$ and $c\nu_\tau$ interactions.

The zero- b categories extend the sensitivity to the large β_L^{23} region explored in this thesis. As β_L^{23} increases, the $s\tau$ interaction also becomes sizeable, producing signal events without a final-state b -quark. This contribution is particularly relevant for resonant production, where an on-shell U_1 can decay into $s\tau$. The four categories therefore provide complementary sensitivity to the different flavour and production configurations of the U_1 signal.

Chapter 4

LHC-ATLAS Experiment

4.1 Large Hadron Collider

The European Organisation for Nuclear Research (CERN) is an international particle-physics laboratory located near Geneva on the border between France and Switzerland. Since its foundation in 1954, CERN has developed and operated a chain of accelerators and experimental facilities to study the fundamental constituents of matter and their interactions [65]. The Large Hadron Collider (LHC) is the largest and highest-energy particle collider in the CERN accelerator complex. It is installed in the former Large Electron–Positron Collider (LEP) tunnel, which has a circumference of about 27 km and lies between approximately 45 m and 170 m below the surface [66, 67]. A schematic overview of the LHC ring and its interaction points is shown in Fig. 4.1.

The LHC was proposed as the next major collider after LEP. The first circulating beams were achieved in 2008, and the first physics run began in 2010. The discovery of a new boson compatible with the Standard Model (SM) Higgs boson by the ATLAS and CMS experiments in 2012 marked one of the central achievements of the LHC physics programme [11, 12]. Beyond Higgs-boson measurements and precision studies of SM processes, a major purpose of the LHC is to search for new particles and interactions beyond the SM.

The LHC collides protons at high centre-of-mass energies in order to extend the kinematic reach for the production of heavy particles. For two proton beams with equal beam energy E_{beam} , the proton–proton centre-of-mass energy is approximately $\sqrt{s} = 2E_{\text{beam}}$. However, the hard-scattering process in a proton–proton (pp) collision occurs between partons, namely quarks and gluons, which carry only fractions of the proton momenta. The partonic centre-of-mass energy therefore varies from event to event and is generally lower than the nominal pp collision energy. Increasing $\sqrt{s}$ consequently extends the accessible mass range for new heavy particles.




Fig. 4.1 Schematic layout of the CERN accelerator complex, including the LHC ring and its main interaction regions [68].

Another key machine parameter is the luminosity, which quantifies the collision rate per unit cross section. For a process with cross section σ , the production rate is

$$R = \mathcal{L}\sigma, \quad (4.1)$$

where $\mathcal{L}$ is the instantaneous luminosity. For two Gaussian bunches colliding with frequency f_{coll} , the luminosity can be expressed as [69]

$$\mathcal{L} = f_{\text{coll}} \frac{n_1 n_2}{4\pi\sigma_x^* \sigma_y^*} \mathcal{F}, \quad (4.2)$$

where n_1 and n_2 are the numbers of particles in the two colliding bunches, σ_x^* and σ_y^* are the RMS transverse beam sizes in the horizontal and vertical directions at the interaction point, and $\mathcal{F}$ is a geometrical reduction factor accounting for effects such as the crossing angle. The integrated luminosity,

$$\mathcal{L}_{\text{int}} = \int \mathcal{L} dt, \quad (4.3)$$

determines the total expected number of produced events by $N = \sigma \mathcal{L}_{\text{int}}$. High instantaneous luminosity increases the event rate, while high integrated luminosity improves the sensitivity to rare processes with small production cross sections.

Four major experiments are installed around the LHC ring. ATLAS, located at Point 1, and CMS, located at Point 5, are general-purpose detectors designed for a broad physics programme including Higgs-boson measurements, precision SM studies, and searches for new particles. ALICE, located at Point 2, is optimised for heavy-ion physics and the study of strongly interacting matter at high energy density. LHCb, located at Point 8, is designed for precision studies of heavy-flavour hadrons and flavour-changing processes [66]. The analysis presented in this thesis uses data recorded by the ATLAS detector.

The LHC physics programme is organised into run periods. Run 1 took place from 2010 to 2012, with pp collision data recorded at $\sqrt{s} = 7$ TeV and $\sqrt{s} = 8$ TeV. Run 2 took place from 2015 to 2018 at $\sqrt{s} = 13$ TeV. After standard data-quality selections, the full ATLAS Run 2 pp dataset corresponds to an integrated luminosity of $140.1 \pm 1.2 \text{ fb}^{-1}$, with a relative uncertainty of 0.83% [70]. Run 3 began in 2022 after the second long shutdown and operated at the higher centre-of-mass energy of $\sqrt{s} = 13.6$ TeV [71]. The Run 2 dataset is used for the leptoquark search presented in this thesis.

An important consequence of high-luminosity operation at the LHC is pile-up. The proton beams are divided into bunches, and a bunch crossing occurs when bunches from the two beams pass through the interaction point. Several pp interactions can occur in a single bunch crossing. The interaction corresponding to the hard-scattering process of interest is referred to as the hard-scatter interaction, while additional inelastic pp interactions in the same bunch crossing constitute in-time pile-up. Detector signals may also originate from interactions in neighbouring bunch crossings, depending on the detector response time. This contribution is referred to as out-of-time pile-up.

The mean number of inelastic pp collisions per bunch crossing is commonly denoted by μ . During Run 2, the nominal bunch spacing was 25 ns, corresponding to a maximum bunch-crossing frequency of 40 MHz. The luminosity-weighted mean value over the full Run 2 dataset was $\langle \mu \rangle = 33.7$, corresponding to approximately 34 interactions per bunch crossing [72].

Pile-up has an important impact on event reconstruction. It produces extra charged-particle tracks, calorimeter energy deposits, and reconstructed jets. These contributions can affect the reconstruction of the primary vertex, jet energies, hadronically decaying τ -leptons, flavour-tagged jets, and missing transverse momentum. Pile-up effects must therefore be modelled in simulated samples and mitigated through dedicated reconstruction and calibration procedures.

4.2 ATLAS experiment

4.2.1 Overview

ATLAS, which stands for A Toroidal LHC ApparatuS, is a detector located at Point 1 of the Large Hadron Collider. It was designed to exploit the full physics potential of the LHC, including precision measurements of SM processes, studies of the Higgs boson, and searches for new physics beyond the SM. The detector surrounds the nominal interaction point and provides nearly full solid-angle coverage, allowing the reconstruction of a wide range of final-state particles produced in proton–proton collisions.

The ATLAS detector is approximately 44 m long, 25 m in diameter, and has a total weight of about 7000 tonnes [73]. It has a cylindrical, forward–backward symmetric layout centred on the interaction point. A schematic view of the detector is shown in Figure 4.2.




Fig. 4.2 Schematic view of the ATLAS detector and its subdetector systems [73].

Starting from the beam line and moving outwards, the main detector systems are the Inner Detector (ID), the calorimeters, and the Muon Spectrometer (MS):

- **Inner Detector (ID)** is immersed in a 2 T axial magnetic field provided by a thin superconducting solenoid and is used to reconstruct charged-particle trajectories close to the interaction point.
- **Calorimeters** surround the ID and measure the energies of electrons, photons, hadrons, hadronically decaying τ -leptons, and jets.

- ***Muon Spectrometer (MS)*** forms the outermost major detector system and measures muons in the magnetic field generated by large superconducting toroids.

The magnetic fields provided by the solenoid and toroids allow the momentum and charge of charged particles to be determined from the curvature of their trajectories. The trigger and data acquisition system selects potentially interesting events from the high LHC bunch-crossing rate for permanent storage and offline reconstruction.

The detector configuration described in this chapter corresponds primarily to Run 2. Several detector and trigger upgrades were installed for Run 3 to cope with higher luminosity conditions, but these upgrades are not used in the analysis presented in this thesis [74].

4.2.2 Coordinate system

ATLAS uses a right-handed coordinate system centred at the nominal interaction point. The z -axis is defined along the beam direction, the x -axis points from the interaction point towards the centre of the LHC ring, and the y -axis points upwards [73]. Cylindrical coordinates (r, ϕ) are used in the plane transverse to the beam, where r is the distance from the beam axis and ϕ is the azimuthal angle around the z -axis.

The polar angle θ is measured with respect to the positive z -axis. The pseudorapidity η is defined as

$$\eta = -\ln \tan \left(\frac{\theta}{2} \right). \quad (4.4)$$

The transverse momentum is defined as $p_T = p \sin \theta$, and the transverse energy is defined similarly. The angular separation between two reconstructed objects is commonly expressed as

$$\Delta R = \sqrt{(\Delta \eta)^2 + (\Delta \phi)^2}. \quad (4.5)$$

These quantities are used throughout this thesis to describe object kinematics, angular separations, overlap removal, and event selections. Figure 4.3 illustrates the coordinate system used in ATLAS.

4.2.3 Magnet system

The ATLAS magnet system consists of four large superconducting magnets: a central solenoid, one barrel toroid, and two end-cap toroids, as shown in Figure 4.4. The central solenoid surrounds the Inner Detector with a length of approximately 5.3 m and provides an




Fig. 4.3 The coordinate system used in ATLAS [75].

axial magnetic field of 2 T for charged-particle momentum measurements in the tracking system. The solenoid is designed to minimise the amount of inactive material in front of the electromagnetic calorimeter.

The toroid system provides magnetic fields of approximately 0.5 T and 1 T in the barrel and end-cap regions of the Muon Spectrometer, respectively. The barrel toroid surrounds the calorimeter system in the central region, while the two end-cap toroids provide bending power in the forward regions. The toroidal field bends muon trajectories, allowing their momenta to be measured using the precision tracking chambers of the MS. The air-core design of the toroids reduces multiple scattering and helps preserve the momentum resolution for high- p_T muons [73]. The combination of the solenoid and toroid systems allows charged-particle momenta to be measured independently in the ID and, for muons, in the MS.

4.2.4 Inner Detector

The Inner Detector (ID) is the innermost tracking system of ATLAS and provides tracking for charged particles within $|\eta| < 2.5$ [73]. It is immersed in the 2 T magnetic field of the central solenoid and records local position measurements, called hits, along their trajectories. During Run 2, it consisted of the pixel detector, the Semiconductor Tracker (SCT), and the Transition Radiation Tracker (TRT). The Insertable B-Layer (IBL) is installed before Run 2 as the innermost pixel layer [76]. Figure 4.5 shows a transverse view of the Inner Detector.

- *Insertable B-Layer (IBL) and Pixel detector*

The pixel detector provides high-granularity measurements close to the beam line. Its precise spatial resolution is essential for reconstructing tracks near the interaction point and for identifying displaced vertices from the decays of long-lived particles such




Fig. 4.4 Layout of the ATLAS superconducting magnet system, including the central solenoid, barrel toroid, and end-cap toroids [73].

1032 as b - and c -hadrons. The IBL further improves the impact-parameter resolution and
 1033 enhances the robustness of tracking and flavour tagging under high pile-up conditions.

1034 • ***Semiconductor Tracker (SCT)***

1035 The SCT surrounds the pixel detector and forms the intermediate precision-tracking
 1036 system of the ID. It is a silicon microstrip detector covering $|\eta| < 2.5$, consisting of four
 1037 cylindrical barrel layers and two end-caps with nine disks on each side. Each module
 1038 contains silicon strip sensors arranged back-to-back with a small stereo angle, allowing
 1039 two-dimensional position information to be reconstructed. The SCT typically provides
 1040 four space-point measurements per track, improving pattern recognition, momentum
 1041 reconstruction, and vertex reconstruction when combined with the high-granularity
 1042 pixel measurements.

1043 • ***Transition Radiation Tracker (TRT)***

1044 The TRT is the outermost tracking subsystem of the ID and provides tracking informa-
 1045 tion at larger radii, covering approximately $|\eta| < 2.0$. It is based on thin straw-tube
 1046 drift chambers, which provide a large number of measurements along the trajectory
 1047 of each charged particle. Although the spatial precision of each TRT hit is lower than
 1048 that of the silicon detectors, the large number of straw hits and the longer lever arm
 1049 improve the momentum measurement and help stabilise track reconstruction.

1050 Together, these subsystems provide the tracking information needed in ATLAS.




Fig. 4.5 Transverse view of the ATLAS Inner Detector, showing the IBL, pixel detector, SCT, and TRT subsystems [76].

4.2.5 Calorimeter

The ATLAS calorimeter system surrounds the Inner Detector and measures the energies and directions of particles [73]. It consists of electromagnetic (EM) and hadronic calorimeters, as shown in Figure 4.6. Specifically, it provides measurements of charged particles, photons and jets, and covers the region up to $|\eta| < 4.9$. It also provides the shower information to the Level-1 trigger system.

- **Electromagnetic (EM) calorimeter**

The EM calorimeter uses liquid argon (LAr) as the active medium and lead absorbers arranged in an accordion geometry, as shown in Figure 4.7. The accordion geometry provides continuous azimuthal coverage and fine segmentation of electromagnetic showers. The EM calorimeter is divided into a barrel section covering $|\eta| < 1.475$ and two end-cap sections covering $1.375 < |\eta| < 3.2$. It is used to reconstruct and identify electrons and photons and also contributes to the reconstruction of hadronically decaying τ -leptons, particularly through the measurement of electromagnetic energy deposits from neutral-pion decays, which don't leave the charged-particle tracks in the ID.

The EM calorimeter consists of a presampler followed by three longitudinal sampling layers, each segmented into readout cells, as illustrated in Figure 4.8. The presampler




Fig. 4.6 Schematic view of the ATLAS calorimeter system [73].

is installed for $|\eta| < 1.8$ and is used to correct for energy lost in material upstream of the calorimeter. The first sampling layer has fine segmentation in η , which is useful for resolving the transverse shape of electromagnetic showers, while the middle layer collects the largest fraction of the shower energy. The back layer provides an additional measurement of the longitudinal shower development.

The energy resolution of the electromagnetic calorimeter can be parameterised as

$$\frac{\sigma_E}{E} = \frac{a}{\sqrt{E}} \oplus b \oplus \frac{c}{E}, \quad (4.6)$$

where a is the intrinsic sampling term, b is the constant term, and c represents the contribution from electronic and pile-up noise [77].

- **Hadronic calorimeter**

The hadronic calorimeter uses different structure for different η regions. In the region of $|\eta| < 1.7$, the Tile Calorimeter uses steel absorbers and scintillating tiles as the active medium. It measures the energies of hadrons and jets. In the end-cap and forward regions, liquid-argon calorimeters are used. The hadronic end-cap calorimeter consists of two wheels per end-cap, covers the range $1.5 < |\eta| < 3.2$, and uses copper absorbers. The forward calorimeter extends the coverage to $3.1 < |\eta| < 4.9$ and uses




Fig. 4.7 Photograph of the EM barrel (left) and end-cap (right) LAr modules. The accordion absorber structure is visible [73].




Fig. 4.8 Cut-away view of an ATLAS LAr electromagnetic calorimeter module, showing the longitudinal sampling layers and the segmentation of the readout cells [73].

1085 dense absorber materials suitable for the high-radiation environment close to the beam
1086 line.

1087 4.2.6 Muon Spectrometer

1088 The Muon Spectrometer (MS) forms the outermost major detector system of ATLAS. It is
1089 embedded in the magnetic field produced by the barrel and end-cap air-core toroid magnets.
1090 The bending of muon trajectories in this field is measured by precision chambers, and the
1091 track sagitta is used to determine the muon momentum. The MS provides precision muon
1092 measurements in the range $|\eta| < 2.7$, while the dedicated muon trigger chambers cover the
1093 region up to approximately $|\eta| < 2.4$ [73].

1094 During Run 2, the MS consisted of Monitored Drift Tubes (MDTs), Cathode Strip
1095 Chambers (CSCs), Resistive Plate Chambers (RPCs) and Thin Gap Chambers (TGCs). A
1096 view of these chambers is shown in Figure 4.9.

1097 • *Monitored Drift Tubes (MDTs)*

1098 The MDT chambers provide precision tracking over most of the MS acceptance. They
1099 consist of aluminium drift tubes with a diameter of about 30 mm, each containing a
1100 central sense wire. A muon crossing a tube ionises the gas, and the drift time of the
1101 electrons to the wire is measured to determine the track position. Several tube layers
1102 are grouped into chamber stations, providing precise measurements in the bending
1103 plane of the toroidal magnetic field. MDT chambers are used over most of the region
1104 $|\eta| < 2.7$, except for the innermost forward stations at $2.0 < |\eta| < 2.7$, where CSCs
1105 are used.

1106 • *Cathode Strip Chambers (CSCs)*

1107 The CSCs are multi-wire proportional chambers used in the innermost forward muon
1108 stations, where the particle rates are highest. They cover approximately $2.0 < |\eta| < 2.7$
1109 and provide precision measurements in the forward region. Their small drift distance
1110 and segmented cathode-strip readout allow them to operate efficiently in the high-rate
1111 environment close to the beam line.

1112 • *Resistive Plate Chambers (RPCs)*

1113 The RPCs provide fast muon trigger information in the barrel region, covering approx-
1114 imately $|\eta| < 1.05$. They are gaseous parallel-plate detectors operated in avalanche
1115 mode. Their fast time response allows the correct bunch crossing to be identified
1116 for each muon candidate and provides the Level-1 muon trigger in the barrel. The

RPCs also provide coordinate measurements complementary to the precision MDT chambers.

- *Thin Gap Chambers (TGCs)*

The TGCs are multi-wire proportional chambers designed to provide fast signals for the Level-1 trigger. They extend to approximately $|\eta| < 2.7$ and provide trigger coverage in the end-cap region up to $|\eta| < 2.4$. The TGC system identifies high- p_T muon candidates and assigns them to the correct bunch crossing. It also provides a second coordinate measurement, complementing the precision tracking information from MDTs and CSCs.




Fig. 4.9 Cross-section of the ATLAS Muon Spectrometer, showing the barrel and end-cap muon chambers. The green and cyan chambers represent the MDT system, while the dashed lines indicate infinite-momentum muon trajectories [73].

During the second long shutdown, the innermost end-cap muon stations were replaced by the New Small Wheels (NSWs) for Run 3 operation. This upgrade was introduced to maintain good muon trigger and reconstruction performance under higher luminosity and pile-up conditions. The NSW detectors are therefore not relevant to the Run 2 dataset analysed in this thesis.

4.2.7 Data Acquisition and Trigger System

At the LHC, proton bunches can cross at a rate of up to 40 MHz for a bunch spacing of 25 ns. This rate is far above the available detector readout and permanent-storage bandwidth. The ATLAS trigger and data acquisition (TDAQ) system is therefore designed to select potentially interesting events online and reduce the event rate to a manageable level. ATLAS uses a two-level trigger system consisting of a hardware-based Level-1 (L1) trigger and a software-based High-Level Trigger (HLT) [78].

Level-1 (L1) trigger

The L1 trigger is the first stage of the ATLAS online event selection. It is implemented in custom hardware and uses reduced-granularity detector information to make a fast decision with a latency of approximately $2.5 \mu\text{s}$. It reduces the input rate from up to 40 MHz to at most 100 kHz. The L1 trigger was based on information from the calorimeter and muon trigger systems, together with topological selections applied to L1 trigger objects. A schematic overview of the Run 2 TDAQ system is shown in Figure 4.10.




Fig. 4.10 Schematic overview of the ATLAS Run-2 trigger and data acquisition system [78].

• *L1 calorimeter trigger (L1Calo)*

L1Calo uses reduced-granularity information from the electromagnetic and hadronic calorimeters to reconstruct energy deposits within the Level-1 latency. Signals from neighbouring calorimeter cells are combined into trigger towers in the (η, ϕ) plane, and dedicated hardware algorithms search for local maxima in the transverse-energy distribution by comparing the energies of neighbouring towers. The same calorimeter information is also summed over larger regions to reconstruct jet-like energy deposits and global quantities such as the total and missing transverse energy.

• *L1 muon trigger (L1Muon)*

L1Muon identifies muon candidates using fast signals from the RPCs and the TGCs. Hits in different trigger-chamber layers are required to form spatial coincidence patterns consistent with a muon trajectory, while their timing information is used to assign the candidate to the correct bunch crossing. The amount of bending in the magnetic field is used to determine whether the candidate satisfies predefined transverse-momentum thresholds.

• *L1 topological trigger (L1Topo)*

L1Topo receives compact trigger-object information reconstructed by L1Calo and L1Muon, including their transverse momentum or energy and (η, ϕ) positions. Dedicated FPGA hardware combines these objects and calculates event-level topological quantities. And these quantities are compared with predefined selection requirements.

The trigger decisions from L1Calo, L1Muon, and L1Topo are then sent to the Central Trigger Processor (CTP), which combines them according to the predefined L1 trigger menu.

If at least one enabled trigger item is satisfied, the CTP issues a Level-1 Accept (L1A) signal for the corresponding bunch crossing. The L1A signal is distributed to the detector front-end electronics and initiates the transfer of the detector data for that bunch crossing from the front-end buffers to the readout system.

For accepted events, the L1 trigger also provides Regions of Interest (RoIs), which identify detector regions containing the trigger objects responsible for the L1 decision. These RoIs provide seed information for the HLT and allow detailed reconstruction to be performed initially in limited detector regions, reducing the required processing time.

High-Level Trigger

The HLT is the second stage of the ATLAS trigger system. It is implemented in a large computing farm and runs software algorithms using detector information with substantially

1178 finer granularity than is available at L1. The HLT reconstruction is guided by RoIs identified
1179 by the L1 trigger, while full-event reconstruction can also be used for signatures such
1180 as jets and missing transverse momentum. The HLT algorithms are designed to provide
1181 reconstruction performance close to that of the offline algorithms while remaining sufficiently
1182 fast for online operation. As a result, the HLT reduced the L1 output rate to an average
1183 recording rate of approximately 1 kHz for permanent storage.

1184 The trigger menu comprises the active L1 and HLT selections used during data taking with
1185 prescale factors, which are applied to selected trigger items to control their rates according to
1186 the instantaneous luminosity and available bandwidth.

Chapter 5

Data and Monte Carlo Simulations

Monte Carlo (MC) simulation is used to model the signal and Standard Model background processes and to predict their event yields and kinematic distributions in this search. Event generation begins by sampling the incoming partons and their momentum fractions from proton PDFs and evaluating the matrix element for the hard-scattering process. The outgoing partons are then evolved through the parton shower and hadronisation, followed by the decays of unstable particles. Finally, pile-up interactions and the ATLAS detector response are simulated, after which the events are reconstructed with the same algorithms used for collision data.

This chapter describes the collision data and simulated event samples used in the search. The Run 2 dataset is presented first, followed by the simulated signal and Standard Model background samples. The trigger selections are summarised at the end of the chapter.

5.1 Dataset selection

The search uses the full LHC Run 2 proton–proton collision dataset recorded by the ATLAS detector between 2015 and 2018 at a centre-of-mass energy of $\sqrt{s} = 13$ TeV. Only data recorded during stable-beam periods and passing the data-quality requirements are retained [72]. These periods are defined by the ATLAS good-runs lists (GRLs).

The selected dataset corresponds to an integrated luminosity of $140.1 \pm 1.2 \text{ fb}^{-1}$, with a relative uncertainty of 0.83% [70]. The integrated luminosity recorded in each year is summarised in Table 5.1.

The mean pile-up during Run 2 was $\langle\mu\rangle \simeq 34$, as described in Section 4.1.

Table 5.1 Integrated luminosity of the Run 2 proton–proton collision data used in this analysis.

Year	Integrated luminosity [fb^{-1}]
2015	3.24 ± 0.04
2016	33.40 ± 0.30
2017	44.63 ± 0.50
2018	58.79 ± 0.64
Total	140.07 ± 1.17

5.2 Signal samples

The signal samples implement the U_1 vector-leptoquark benchmark defined in Section 3.1.5. The Yang–Mills scenario with $\kappa_c = \kappa_Y = 0$ and $\beta_L^{33} = 1$ is used. The simulated signal covers both the pure-left-handed scenario with $\beta_R^{33} = 0$ and the right-handed scenario with $|\beta_R^{33}| = 1$. Apart from their coupling parameters, the same event-generation, parton-shower and detector-simulation settings are used for both scenarios.

Signal events are generated at leading order (LO) in QCD using MADGRAPH5_AMC@NLO 3.3.1 [79], interfaced to PYTHIA 8.308 [80]. The hard-scattering process produces a τ -lepton, a ν_τ and up to two additional partons, which can be gluons or u -, d -, s -, c - or b -quarks. The NNPDF3.0nlo PDF set [81] is used for the hard-scattering calculation.

The generation is performed in the five-flavour scheme, in which the b -quark is treated as an active parton in the proton. The proton PDFs therefore allow processes with an initial-state b -quark. All allowed U_1 diagrams contributing to this final state are retained in the matrix-element calculation. The resonant single-production and non-resonant t -channel processes shown in Figure 3.4 are therefore included coherently and cannot be separated at the generation stage. Other allowed contributions, including leptoquark pair production where kinematically accessible, are also retained.

PYTHIA subsequently simulates the parton shower and hadronisation. It also models the underlying event using the A14 tune [82] and the NNPDF2.3lo PDF set [83]. The CKKW–L merging procedure [84] prevents the same radiation from being included in both the matrix-element and parton-shower calculations. The Durham $k_\perp$ merging scale is set to 15 GeV and the D parameter to 0.4. Together, they determine whether an emission is treated as an additional matrix-element parton or as part of the shower.

For the pure-left-handed interpretation, dedicated signal samples are generated for 152 combinations of (M_U, g_U, β_L^{23}) satisfying

$$1.5 \text{ TeV} \leq M_U \leq 3.0 \text{ TeV}, \quad (5.1)$$

$$0.5 \leq g_U \leq 3.0, \quad (5.2)$$

$$0 \leq \beta_L^{23} \leq 2.2, \quad (5.3)$$

with $\beta_L^{33} = 1$ and $\beta_R^{33} = 0$. The lower boundary of the mass scan covers the region around the exclusion limits obtained from previous leptoquark pair-production searches [57, 58]. For $M_U > 3 \text{ TeV}$, the selected signal is dominated by non-resonant exchange, where the sensitivity is governed mainly by the ratio g_U/M_U . The primary mass scan is therefore restricted to $M_U \leq 3 \text{ TeV}$.

For the right-handed interpretation with $|\beta_R^{33}| = 1$, one baseline sample is generated at each of the six mass points $M_U \in \{1.5, 2.0, 2.5, 3.0, 4.0, 5.0\} \text{ TeV}$ with $g_U = 1$ and $\beta_L^{23} = 0.2$. Predictions at other values of g_U and β_L^{23} are obtained from these baseline samples using corrected MADGRAPH5_AMC@NLO internal event weights, rather than from separately generated samples at every parameter point. The correction includes an adjustment of the normalisation and a two-dimensional truth-level shape correction based on the leading-jet p_T and $m_{b\tau}$. Details of the correction and its validation are given in Appendix A. This strategy avoids repeating event generation and detector simulation for every coupling configuration, thereby reducing the required computing resources.

The U_1 amplitude can interfere with the SM amplitude for production of the same $\tau\nu_\tau + \text{jets}$ final state. The total matrix element can be expressed as

$$|\mathcal{M}_{\text{SM}} + \mathcal{M}_{U_1}|^2 = |\mathcal{M}_{\text{SM}}|^2 + |\mathcal{M}_{U_1}|^2 + 2 \text{Re}(\mathcal{M}_{\text{SM}}\mathcal{M}_{U_1}^*). \quad (5.4)$$

The first term represents the SM contribution, the second represents the pure U_1 contribution, corresponding to the BSM-only component, and the third represents their interference. Unlike the BSM-only component, the interference term can be negative and can reduce the predicted yield relative to the SM-only expectation in some regions of phase space.

To separate the BSM-only and interference terms in Eq. (5.4), the two contributions are generated independently with MADGRAPH5_AMC@NLO using the configuration described above. The BSM-only component is selected by requiring exactly two new-physics couplings at the amplitude level, denoted by NP==2, whereas the interference component is isolated at the squared-matrix-element level by requiring $\text{NP}^2 = 2$. The interference sample therefore contains only the third term in Eq. (5.4). It does not include the pure-SM term, which is already represented by the nominal SM background samples.

All τ -lepton decay modes are retained in the signal simulation. Although the signal regions target hadronically decaying τ -leptons, leptonic decays can contribute to regions containing electrons or muons. The inclusive decay configuration allows this signal contamination to be evaluated.

The decays of bottom- and charm-hadrons are modelled with EVTGEN [85]. All signal samples used in this thesis are processed with the ATLAS fast detector simulation ATLFast3 [86], which provides a computationally efficient simulation of the calorimeter response.

5.3 Background samples

The simulated SM backgrounds considered in this analysis are $W + \text{jets}$, $Z + \text{jets}$, $t\bar{t}$, single-top and diboson production. The dominant contribution is $W \rightarrow \tau\nu + \text{jets}$ production. The QCD multijet background is negligible after event cleaning and is therefore neglected, as discussed in Appendix B.

The $W/Z + \text{jets}$ samples are generated with SHERPA 2.2.11 [87], except for $Z \rightarrow \tau\tau$, which uses version 2.2.14. Matrix elements include up to two additional partons at next-to-leading order (NLO) in QCD and up to five at LO. They are merged with the SHERPA parton shower [88] using the MEPS@NLO prescription [89–92].

Only a small fraction of inclusive $W \rightarrow \tau\nu$ events passes the high- E_T^{miss} requirement, leaving limited MC statistics. Two dedicated filtered samples are therefore used to reduce the MC statistical uncertainty in the high- E_T^{miss} tail. Their respective generator-level requirements are

$$70 \text{ GeV} < E_T^{\text{miss}} < 150 \text{ GeV} \quad \text{and} \quad p_T^\tau < 150 \text{ GeV},$$

$$E_T^{\text{miss}} > 150 \text{ GeV} \quad \text{or} \quad p_T^\tau > 150 \text{ GeV}. \quad (5.5)$$

Top-quark pair and single-top production are generated at NLO accuracy in QCD using POWHEG BOX v2 [93–96], interfaced to PYTHIA 8.230 [97] for parton showering and hadronisation. The single-top samples include s -channel, t -channel and associated Wt production. The predicted $t\bar{t}$ yield is normalised to the inclusive production cross-section calculated at next-to-next-to-leading order (NNLO) in QCD with next-to-next-to-leading-logarithmic (NNLL) soft-gluon resummation [98]. A dilepton filter is applied to the $t\bar{t}$ sample, requiring both W bosons to decay leptonically.

Diboson production is also modelled with SHERPA.

The hard-scattering calculations use the NNPDF3.0nnlo PDF set for $W/Z + \text{jets}$ and NNPDF3.0nlo for $t\bar{t}$ and single-top production [81]. The underlying event is modelled using the A14 tune for PYTHIA and the default tune for SHERPA. For samples not generated with SHERPA, the decays of bottom- and charm-hadrons are modelled with EVTGEN [85]. The full ATLAS detector simulation based on GEANT4 [99, 100] is used for all background samples.

The principal MC generator configurations are summarised in Table 5.2.

Table 5.2 Summary of the principal MC generator configurations used for the signal and SM background processes. The quoted perturbative order refers to the hard-scattering matrix-element calculation. For the $W/Z + \text{jets}$ samples, matrix elements with up to two additional partons are calculated at NLO and those with up to five additional partons at LO. The symbol ℓ denotes an electron or muon.

Process	Generator	ME order	Parton shower	ME PDF set	Tune
U_1 signal	MADGRAPH5_AMC@NLO 3.3.1	LO	PYTHIA 8	NNPDF3.0nlo	A14
$W \rightarrow \ell\nu + \text{jets}$	SHERPA 2.2.11	NLO	SHERPA	NNPDF3.0nnlo	SHERPA default
$W \rightarrow \tau\nu + \text{jets}$	SHERPA 2.2.11	NLO	SHERPA	NNPDF3.0nnlo	SHERPA default
$Z \rightarrow ee/\mu\mu/\nu\nu + \text{jets}$	SHERPA 2.2.11	NLO	SHERPA	NNPDF3.0nnlo	SHERPA default
$Z \rightarrow \tau\tau + \text{jets}$	SHERPA 2.2.14	NLO	SHERPA	NNPDF3.0nnlo	SHERPA default
$t\bar{t}$	POWHEG BOX v2	NLO	PYTHIA 8	NNPDF3.0nlo	A14
Single- t	POWHEG BOX v2	NLO	PYTHIA 8	NNPDF3.0nlo	A14
Diboson	SHERPA	NLO	SHERPA	NNPDF3	SHERPA default

5.4 Triggers

The ATLAS Run 2 trigger architecture was described in Chapter 4. This analysis uses only single-electron, single-muon and E_T^{miss} triggers.

The signal-region event selection uses the lowest-threshold unprescaled E_T^{miss} trigger, which is fully efficient for at least offline $E_T^{\text{miss}} > 200$ GeV applied in signal regions and the corresponding validation regions.

Single-electron and single-muon triggers are used in regions containing an electron or muon. The lowest online thresholds range from 20 to 26 GeV [101, 102].

Table 5.3 lists the single-electron, single-muon and E_T^{miss} trigger chains used in this analysis.


In an HLT chain name, e, mu and xe represent an electron, a muon and E_T^{miss} , respectively. The following number gives the online threshold in GeV. An explicit L1 suffix identifies the first-level trigger that seeds the chain. The electron labels lhloose, lhmedium and lhtight specify likelihood-based identification requirements, while nod0 indicates that transverse-impact-parameter information is omitted. The labels iloose, ivarloose and ivarmedium

specify isolation requirements. For E_T^{miss} , mht denotes a calculation based on the negative vector sum of jet transverse momenta, whereas pufit denotes a calorimeter-based calculation with a fit-based pile-up correction [103].


Figures 5.1 and 5.2 show representative Run 2 trigger-efficiency measurements. Differences in trigger efficiency between collision data and simulation are corrected using data-to-simulation efficiency scale factors.

Table 5.3 Single-electron, single-muon and E_T^{miss} trigger chains used in the Run 2 analysis. Single-lepton chains listed for the same flavour and year are combined with a logical OR, while the E_T^{miss} chain is selected according to the data-taking period. The extra xe70 or xe65 in the 2018 names denotes an additional calorimeter-cell E_T^{miss} requirement of 70 or 65 GeV.


Trigger	Year	HLT chain
Single-electron	2015	HLT_e24_lhmedium_L1EM20VH
		HLT_e60_lhmedium
		HLT_e120_lhloose
	2016–2018	HLT_e26_lhtight_nod0_ivarloose
		HLT_e60_lhmedium_nod0
		HLT_e140_lhloose_nod0
Single-muon	2015	HLT_mu20_iloose_L1MU15
		HLT_mu40
	2016–2018	HLT_mu26_ivarmedium
		HLT_mu50
E_T^{miss}	2015	HLT_xe70_mht
	2016	HLT_xe90_mht_L1XE50
		HLT_xe100_mht_L1XE50
		HLT_xe110_mht_L1XE50
	2017	HLT_xe110_pufit_L1XE55
		HLT_xe110_pufit_L1XE50
	2018	HLT_xe110_pufit_xe70_L1XE50
		HLT_xe110_pufit_xe65_L1XE50



(a) Single-electron trigger combination.



(b) Single-muon triggers, barrel.



(c) Single-muon triggers, end-cap.

Fig. 5.1 Single-lepton trigger efficiencies measured in Run 2 $Z \rightarrow \ell\ell$ events. (a) Electron efficiency versus offline E_T for electrons with Tight identification and FCTight isolation [101]. (b,c) Muon efficiency versus offline p_T in the barrel and end-cap, respectively [102]. Lower panels give data-to-simulation efficiency ratios. Electron uncertainties include statistical and systematic contributions, while muon uncertainties are statistical only.




Fig. 5.2 Combined L1 and HLT E_T^{miss} trigger efficiencies in $Z \rightarrow \mu\mu$ events for each year of Run 2 [103]. Since muons are treated as invisible by these calorimeter-based triggers, the dimuon transverse momentum $p_T(\mu\mu)$ provides a reference for the missing transverse momentum measured by the trigger. The curves use the lowest unprescaled trigger as it changes within each year. Error bars show statistical uncertainties.

1324 **Chapter 6**

1325 **Object Reconstruction and Identification**

1326 **6.1 Track**

1327 **6.2 Primary vertex**

1328 **6.3 Electron**

1329 **6.4 Muon**

1330 **6.5 Hadronic tau**

1331 **6.6 Jet**

1332 **6.7 Jet flavor tagging**

1333 **6.8 Overlap removal**

1334 **6.9 Missing transverse momentum**

Chapter 7

Event Selection

The event selection separates the four signal topologies introduced in Section 3.1.5 according to the presence of a b -tagged jet and the kinematics of resonant or non-resonant production. The analysis variables and common preselection are introduced first, followed by the optimisation of the one- b -jet regions and the definition of the zero- b -jet regions.

7.1 Preselection and variables definition

The analysis variables are constructed from the calibrated objects defined in Chapter 6, after overlap removal. Throughout this chapter, τ denotes the reconstructed visible decay products of the selected hadronically decaying τ -lepton. For signal events, the neutrino from the τ decay is not included in this four-momentum and contributes to $\vec{p}_T^{\text{miss}}$ together with the neutrino produced in the hard scattering.

7.1.1 Analysis variables

The transverse momenta, angular separations and reconstructed masses characterise the energy scale and geometry of the $\tau\nu$ + jets final state. They distinguish the signal from W +jets and top-quark production, and separate the resonant and non-resonant signal contributions.

- **The τ transverse momentum and E_T^{miss}** measure the visible and invisible transverse activity in the event. Heavy-leptoquark production can produce an energetic τ -lepton and neutrinos, whereas most SM events populate lower values of these quantities. Requiring large p_T^τ and E_T^{miss} therefore selects the high-energy part of the $\tau\nu$ spectrum in which a leptoquark contribution can become distinguishable [64]. The two quantities need not be equal because the jets carry transverse momentum and the τ -decay neutrino contributes to $\vec{p}_T^{\text{miss}}$.

- **The azimuthal separation** $\Delta\phi(\tau, E_T^{\text{miss}})$ is the angle between the visible τ momentum and $\vec{p}_T^{\text{miss}}$ in the transverse plane, with values between zero and π . In non-resonant production with a relatively soft recoiling jet, the τ and the invisible system tend to be back-to-back. A lower requirement on $\Delta\phi(\tau, E_T^{\text{miss}})$ favours this geometry and reduces top-quark backgrounds with a more complex final state. Resonant production contains an energetic jet from the leptoquark decay, so a back-to-back requirement on the τ and $\vec{p}_T^{\text{miss}}$ can also remove the signal of interest.

- **The transverse mass** m_T combines the two transverse momentum scales with their angular separation:

$$m_T = \sqrt{2p_T^\tau E_T^{\text{miss}} [1 - \cos \Delta\phi(\tau, E_T^{\text{miss}})]}. \quad (7.1)$$

This definition neglects the small visible τ mass. It is useful because the longitudinal momenta of the neutrinos are unmeasured and the full $\tau\nu$ invariant mass cannot be reconstructed. For an on-shell $W \rightarrow \tau\nu$ decay with correctly reconstructed momenta and no other source of $\vec{p}_T^{\text{miss}}$, m_T is bounded by the W -boson mass. Off-shell W production, additional neutrinos and detector resolution populate the high- m_T tail of the SM background. A heavy virtual leptoquark can enhance this tail, making m_T particularly useful for non-resonant production. Equation 7.1 also shows that its selection is correlated with those on p_T^τ , E_T^{miss} and $\Delta\phi(\tau, E_T^{\text{miss}})$.

- **The visible invariant masses** $m_{b\tau}$ and $m_{j\tau}$ are reconstructed from the selected b -jet or leading jet and the visible τ :

$$m_{b\tau}^2 = (p_b + p_\tau)^2, \quad m_{j\tau}^2 = (p_{j_1} + p_\tau)^2, \quad (7.2)$$

where p_b , p_{j_1} and p_τ are four-momenta, and j_1 is the jet with the largest p_T . In resonant $U_1 \rightarrow b\tau$ production, $m_{b\tau}$ retains information about the leptoquark mass. The neutrino from the τ decay shifts the visible mass below M_U and broadens its distribution, with further smearing from the reconstructed jet and τ energies. A lower mass threshold therefore selects heavy-particle decays without requiring a narrow peak at M_U . The analogous variable $m_{j\tau}$ is used in the zero- b -jet regions, where the leading jet can originate from $U_1 \rightarrow s\tau$.

- **The selected-jet transverse momentum** distinguishes the two production topologies. The b -jet from a resonant $U_1 \rightarrow b\tau$ decay is typically energetic because it shares the decay energy of a heavy particle. A b -jet radiated in non-resonant production is

typically softer. The b -jet p_T is therefore used to divide the one- b -jet sample, while the leading-jet p_T is used in the zero- b -jet sample. These reconstructed categories enrich the corresponding production modes without separating them exactly.

- **The object multiplicities** specify the reconstructed final state. The numbers of hadronic τ candidates, electrons, muons, jets and b -tagged jets are denoted by N_τ , N_e , N_μ , N_j and N_b , respectively, with $N_\ell = N_e + N_\mu$. All jets, including b -tagged jets, enter N_j , so $N_j - N_b$ counts jets that are not b -tagged. An upper requirement on this multiplicity suppresses $t\bar{t}$ events, whose two top-quark decays often produce several jets.

7.1.2 Multijet suppression

Multijet events can enter the selected sample when a jet is misidentified as a hadronic τ and a mismeasured jet produces large E_T^{miss} . In particular, an underestimated jet momentum produces a momentum imbalance pointing towards that jet. The minimum azimuthal separation and the corresponding momentum ratio are therefore defined as

$$\Delta\phi_{\min} = \min_j \Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_T^j), \quad R_{\text{miss}/j} = \frac{E_T^{\text{miss}}}{p_{T,j_{\min}}}, \quad (7.3)$$

where $j_{\min}$ is the jet with the smallest azimuthal separation from $\vec{p}_T^{\text{miss}}$. It need not be the leading jet. Mismeasured multijet events tend to have small $\Delta\phi_{\min}$ and a modest $R_{\text{miss}/j}$. By contrast, the neutrinos in signal events can produce large E_T^{miss} even when a much softer jet happens to lie nearby in azimuth.

Events are retained if

$$\Delta\phi_{\min} > 0.4 \quad \text{or} \quad R_{\text{miss}/j} > 6. \quad (7.4)$$

The second condition retains signal events that would be lost with the angular requirement alone. In the simulated study described in Appendix B, this selection rejects approximately 86% of the multijet events while retaining 96.4% of the signal after the loose study selection. These efficiencies describe the cleaning requirement at that stage, before the full SR selections.

7.1.3 Common preselection

The common preselection selects events with one hadronic τ , missing transverse momentum and at least one jet. The data-quality requirements in Section 5.1 and the primary-vertex requirement in Chapter 6 are applied first. Events must pass the lowest-threshold unprescaled

E_T^{miss} trigger for the relevant data-taking period and satisfy $E_T^{\text{miss}} > 200$ GeV, in the trigger-efficiency plateau described in Section 5.4. Exactly one baseline τ candidate is required, and it must also pass the Tight identification working point. Events containing a selected electron or muon are vetoed to suppress backgrounds with additional leptonic decays. At least one jet is required to define the jet-based variables, and the multijet suppression in Eq. 7.4 is applied. Table 7.1 summarises these requirements. The common preselection does not fix N_b or the final p_T^τ threshold, so it can be used for both flavour categories and as the starting point for the kinematic optimisation.

Table 7.1 Common preselection for the signal-region optimisation. Object definitions and overlap removal are given in Chapter 6.

Requirement	Selection
Data quality	Accepted data-taking periods
Primary vertex	At least two selected tracks
Trigger	Lowest-threshold unprescaled E_T^{miss} trigger
Missing transverse momentum	$E_T^{\text{miss}} > 200$ GeV
Hadronic τ multiplicity	$N_\tau = 1$ at baseline level
Selected τ identification	Tight
Electron and muon veto	$N_\ell = 0$
Jet multiplicity	$N_j \geq 1$
Multijet suppression	$\Delta\phi_{\text{min}} > 0.4$ or $R_{\text{miss}/j} > 6$

7.2 One b-jet signal region (SR1b) optimisation and definition

The SR1b regions select exactly one b -tagged jet to target the $b\tau$ interaction in the U_1 model. The non-resonant selection is motivated by the high- m_T $\tau\nu + b$ strategy of Ref. [64]: an energetic τ and large E_T^{miss} select the hard scattering, the b -tag reduces the inclusive W +jets background, and a restriction on additional jets suppresses top-quark production. The analysis also retains a resonant category to select an on-shell leptoquark decaying into $b\tau$. The sample is divided at $p_T^b = 250$ GeV, with $p_T^b > 250$ GeV defining SR1b-Res and $50 < p_T^b < 250$ GeV defining SR1b-NonRes.

7.2.1 Scan variables and optimisation criterion

The kinematic thresholds are scanned together because the discriminating variables are correlated. For example, increasing E_T^{miss} also raises m_T at fixed p_T^τ and $\Delta\phi(\tau, E_T^{\text{miss}})$, so the effect of an additional m_T requirement depends on the other cuts. Lower thresholds on p_T^τ , E_T^{miss} and m_T are scanned in both categories, together with an upper limit on $N_j - N_b$. A lower threshold on $\Delta\phi(\tau, E_T^{\text{miss}})$ is tested for the non-resonant topology. For the resonant topology, upper thresholds on this angle and a lower threshold on $m_{b\tau}$ are tested. The complete scan grid is given in Table 7.2.

Table 7.2 Thresholds tested in the SR1b optimisation. Each set lists the trial values for the indicated inequality. A zero mass threshold or an angular upper threshold of π represents no effective requirement on that variable.

Variable	SR1b-Res	SR1b-NonRes
$\Delta\phi(\tau, E_T^{\text{miss}})$	$< \{0.3, 1.2, 2.4, 2.6, \pi\}$	$> \{0.3, 1.2, 2.4, 2.6\}$
$m_{b\tau}$ [GeV]	$> \{0, 800\}$	Not scanned
E_T^{miss} [GeV]	$> \{200, 250, 300, 350, 400, 500, 600\}$	
p_T^τ [GeV]	$> \{200, 250, 300, 350, 400, 500, 600\}$	
m_T [GeV]	$> \{100, 200, 300, 400, 500, 600, 700, 800, 900, 1000\}$	
$N_j - N_b$		$\leq \{1, 2, 3\}$

The mass scan uses only $m_{b\tau} > 0$ and 800 GeV. The visible mass distribution is broad because of the unmeasured τ -decay neutrino, while the simulated background has limited statistics in its high-mass tail. These features favour an inclusive mass requirement over a finely scanned mass window. The resonant optimisation targets $M_U = 1.5$ TeV, where on-shell production is important. The non-resonant optimisation uses $(M_U, g_U, \beta_L^{23}) = (2.5 \text{ TeV}, 2.5, 1.0)$, where virtual exchange provides the dominant signal contribution.

The expected sensitivity of each cut combination is ranked using an approximate counting-experiment significance [104]:

$$Z^2 = 2 \left[n \ln \left(\frac{n(b + \sigma_b^2)}{b^2 + n\sigma_b^2} \right) - \frac{b^2}{\sigma_b^2} \ln \left(1 + \frac{\sigma_b^2(n - b)}{b(b + \sigma_b^2)} \right) \right], \quad (7.5)$$

where s and b are the expected signal and background event yields, $n = s + b$, and σ_b is the absolute uncertainty in the background expectation. A fractional background uncertainty of 30%, corresponding to $\sigma_b = 0.30b$, is assumed for this scan. The signal yield contains the

pure BSM contribution. Interference with the SM is treated separately in the final statistical analysis. The quantity Z is used to compare selections, while the final confidence limits are obtained with the likelihood analysis in Chapter 10.

Cut combinations with $b \leq 2$ are discarded to avoid selecting apparently background-free regions caused by limited MC statistics. If a particular simulated background has a negative weighted yield after a trial selection, its contribution is set to zero when evaluating the scan criterion. This is a prescription for comparing trial cuts in sparse simulated samples. It does not imply a negative physical event rate.

7.2.2 Signal efficiency and selected regions

The signal efficiency measures how much signal survives the selection. For the optimisation cutflow, the cumulative efficiency is defined with respect to the one- b -jet baseline, before the additional kinematic cuts:

$$\epsilon_{\text{rel}}(\text{SR}) = \frac{N_s(\text{SR})}{N_s(\text{one-}b\text{-jet baseline})}. \quad (7.6)$$

Both yields are weighted predictions for the same signal parameter point. This definition differs from the acceptance times selection efficiency, $A\epsilon$, whose denominator is the generated signal sample. The acceptance describes the fraction of generated events in the required kinematic and geometric range, and the selection efficiency describes the fraction of those events that are reconstructed and pass the selection. Their product relates the selected yield to the generated signal cross section through $N_s = \mathcal{L}\sigma_s A\epsilon$, using a consistent convention for the generated decay modes and any generator filters.

For the resonant benchmark $(M_U, g_U, \beta_L^{23}) = (1.5 \text{ TeV}, 1.5, 0.6)$, the baseline yield of 35.2 events is reduced to 4.2 events, corresponding to $\epsilon_{\text{rel}} \simeq 11.9\%$. For the non-resonant optimisation benchmark, the yield changes from 111.0 to 6.6 events, giving $\epsilon_{\text{rel}} \simeq 5.9\%$. The corresponding total background prediction decreases from approximately 4027 baseline events to 2.4 and 2.1 events, respectively. These numbers illustrate the trade-off between signal retention and background rejection in the optimisation and are evaluated before the background fit.

The selected SR1b-Res requirements are $p_T^\tau > 200 \text{ GeV}$, $E_T^{\text{miss}} > 200 \text{ GeV}$, $m_T > 200 \text{ GeV}$ and $m_{b\tau} > 800 \text{ GeV}$, with at most three jets that are not b -tagged. No additional $\Delta\phi(\tau, E_T^{\text{miss}})$ requirement is retained. The visible-mass requirement provides strong rejection of backgrounds without a heavy $b\tau$ resonance, while the energetic b -jet selects the two-body decay topology. The region is treated as a single event-counting bin because

the available background simulation does not support a finely binned fit to the broad $m_{b\tau}$ distribution.

The selected SR1b-NonRes requirements are $p_T^\tau > 200$ GeV, $E_T^{\text{miss}} > 400$ GeV, $m_T > 600$ GeV and $\Delta\phi(\tau, E_T^{\text{miss}}) > 1.2$, with at most one jet that is not b -tagged. The tighter E_T^{miss} and m_T requirements select the high-energy tail, and the smaller allowed jet multiplicity reduces the top-quark background. No $m_{b\tau}$ requirement is imposed because the b -jet and τ do not reconstruct an on-shell leptoquark in the target topology. This region also contributes a single event count to the fit.

Figure 7.1 shows the visible-mass distribution with the mass requirement removed, conventionally called an $N - 1$ distribution, and the transverse-mass distribution in the selected high- m_T range. The red arrow marks the $m_{b\tau} > 800$ GeV threshold, while the m_T panel displays events above 600 GeV. The background normalisations in these illustrations have been constrained by the background-only fit to the control regions described in Chapter 10. The optimisation itself uses the simulated expectations described above.




Fig. 7.1 Distributions of (a) $m_{b\tau}$ in SR1b-Res before the mass requirement and (b) m_T in SR1b-NonRes above 600 GeV. Background normalisations are obtained from the background-only fit to the control regions. The dotted red and violet signal curves correspond to $(M_U, g_U, \beta_L^{23}) = (1.5 \text{ TeV}, 1.5, 0.6)$ and $(3.0 \text{ TeV}, 2.5, 1.0)$, respectively. The red arrow marks the mass threshold, and overflow events enter the last bin. Lower panels show data divided by the background prediction. The hatched band includes statistical and systematic uncertainties, and upward triangles indicate ratios above the displayed range. The plot label τ_{had} denotes the visible τ candidate.

7.3 b-veto signal region definition (SR0b)

The SR0b regions select signal events with no reconstructed b -tagged jet and extend the sensitivity to large β_L^{23} . As shown by the flavour current and coupling texture in Eqs. 3.6 and 3.7, β_L^{23} controls the $s\tau$ interaction and contributes $V_{cs}\beta_L^{23}$ to the $c\nu_\tau$ interaction. The resulting zero- b -jet topologies were introduced in Section 3.1.5. In particular, $U_1 \rightarrow s\tau$ produces a resonant signal with a genuine non- b jet, so SR0b retains a physically distinct signal contribution in addition to events in which a b -jet is not tagged or reconstructed. A leading jet in this category is not explicitly identified as a strange-quark jet.

The two SR0b regions are defined by replacing the one- b -jet requirement with $N_b = 0$ and using the leading jet in the topology-dependent selections. The resonant region requires $p_T^{j_1} > 250$ GeV and $m_{j\tau} > 800$ GeV, while the non-resonant region requires $50 < p_T^{j_1} < 250$ GeV. The p_T^τ , E_T^{miss} , m_T and $\Delta\phi(\tau, E_T^{\text{miss}})$ thresholds follow the corresponding one- b -jet regions. The upper limits are imposed on the total jet multiplicity: $N_j \leq 4$ for SR0b-Res and $N_j \leq 2$ for SR0b-NonRes. These preserve the total jet limits used in SR1b after replacing the tagged jet with an untagged leading jet.

The $E_T^{\text{miss}} > 200$ GeV requirement in SR0b-Res is retained to keep its kinematic selection aligned with SR1b-Res. A supplementary study of tighter E_T^{miss} thresholds is presented in Appendix C. The SR0b-NonRes threshold remains $E_T^{\text{miss}} > 400$ GeV, which also maximises the significance in that study.

Signal efficiency in SR0b is evaluated for the pure BSM contribution, with $A\epsilon$ defined as in Section 7.2.2. For illustration, at $(M_U, g_U, \beta_L^{23}) = (1.5 \text{ TeV}, 1.0, 0.2)$ with $\beta_R^{33} = 0$, $A\epsilon$ is 0.30% and 0.13% in SR0b-Res and SR0b-NonRes, compared with 0.64% and 0.069% in the corresponding SR1b regions. These efficiencies depend on the mass and flavour couplings because the relative contributions of the production and decay channels change. The interference contribution can have either sign and is included separately in the predicted event yield rather than interpreted as an efficiency.

Figure 7.2 shows the discriminating variables in the two zero- b -jet regions. The high- $m_{j\tau}$ requirement selects the visible products of a heavy resonance, while the high- m_T requirement selects non-resonant production. Together with the one- b -jet regions, these selections retain sensitivity to both $b\tau$ and $s\tau$ final states over the coupling range studied in this thesis.

The final requirements for all four mutually exclusive SRs are summarised in Table 7.3. Each region contributes one event count to the simultaneous statistical analysis.




Fig. 7.2 Distributions of (a) $m_{j\tau}$ in SR0b-Res before the mass requirement and (b) m_T in SR0b-NonRes above 600 GeV. Background normalisations are obtained from the background-only fit to the control regions. The dotted red and violet signal curves correspond to $(M_U, g_U, \beta_L^{23}) = (1.5 \text{ TeV}, 1.5, 0.6)$ and $(3.0 \text{ TeV}, 2.5, 1.0)$, respectively. The red arrow marks the mass threshold, and overflow events enter the last bin. Lower panels show data divided by the background prediction, with the combined statistical and systematic uncertainty shown by the hatched band. Upward triangles indicate ratios outside the displayed range.

Table 7.3 Final signal-region selections, applied in addition to the common preselection in Table 7.1. A dash denotes no additional requirement. The jet multiplicity includes b -tagged jets.

Requirement	Resonant		Non-resonant	
	SR1b-Res	SR0b-Res	SR1b-NonRes	SR0b-NonRes
N_b	1	0	1	0
p_T^b [GeV]	> 250	–	$50 < p_T^b < 250$	–
$p_T^{j_1}$ [GeV]	–	> 250	–	$50 < p_T^{j_1} < 250$
p_T^τ [GeV]			> 200	
E_T^{miss} [GeV]	> 200			> 400
m_T [GeV]	> 200			> 600
$m_{b\tau}$ [GeV]	> 800	–	–	–
$m_{j\tau}$ [GeV]	–	> 800	–	–
$\Delta\phi(\tau, E_T^{\text{miss}})$		–		> 1.2
N_j		≤ 4		≤ 2

1535 **Chapter 8**

1536 **Background Estimation Strategy**

1537 **8.1 Strategy overview**

1538 **8.2 W +jets estimation**

1539 **8.3 $t\bar{t}$ estimation**

1540 **8.4 Fake- τ estimation**

1541 **8.5 Multi-jet cleaning**

1542 **8.6 VRs**

1543 **Chapter 9**

1544 **Systematics Uncertainties**

1545 **9.1 Experimental uncertainties**

1546 **9.2 Theoretical uncertainties**

1547

Chapter 10

1548

Statistical Analysis and Results

1549

10.1 Statistical analysis

1550

10.1.1 Hypothesis test

1551

10.1.2 Likelihood function

1552

10.2 Results

1553

10.2.1 Background-only fit

1554

10.2.2 Unblinded S+B fit in CR+SRs

1555

10.2.3 Exclusion limits with toy models

1556 **Chapter 11**

1557 **Discussion and Conclusion**

References

- 1559 [1] Sheldon L. Glashow. Partial-symmetries of weak interactions. *Nucl. Phys.*, 22:
1560 579–588, 1961. doi: 10.1016/0029-5582(61)90469-2.
- 1561 [2] Steven Weinberg. A model of leptons. *Phys. Rev. Lett.*, 19:1264–1266, 1967. doi:
1562 10.1103/PhysRevLett.19.1264.
- 1563 [3] Abdus Salam. Weak and electromagnetic interactions. In N. Svartholm, editor,
1564 *Elementary Particle Theory: Relativistic Groups and Analyticity*, pages 367–377,
1565 Stockholm, 1968. Almqvist and Wiksell.
- 1566 [4] Wolfgang Pauli. The connection between spin and statistics. *Phys. Rev.*, 58:716–722,
1567 1940. doi: 10.1103/PhysRev.58.716.
- 1568 [5] Gerard 't Hooft. Renormalization of massless yang–mills fields. *Nucl. Phys. B*, 33:
1569 173–199, 1971. doi: 10.1016/0550-3213(71)90395-6.
- 1570 [6] Gerard 't Hooft and Martinus J. G. Veltman. Regularization and renormalization of
1571 gauge fields. *Nucl. Phys. B*, 44:189–213, 1972. doi: 10.1016/0550-3213(72)90279-9.
- 1572 [7] F. Englert and R. Brout. Broken symmetry and the mass of gauge vector mesons.
1573 *Phys. Rev. Lett.*, 13:321–323, 1964. doi: 10.1103/PhysRevLett.13.321.
- 1574 [8] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. *Phys. Rev. Lett.*,
1575 13:508–509, 1964. doi: 10.1103/PhysRevLett.13.508.
- 1576 [9] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble. Global conservation laws and
1577 massless particles. *Phys. Rev. Lett.*, 13:585–587, 1964. doi: 10.1103/PhysRevLett.13.
1578 585.
- 1579 [10] Particle Data Group. Review of particle physics. *Prog. Theor. Exp. Phys.*, 2025.
- 1580 [11] ATLAS Collaboration. Observation of a new particle in the search for the standard
1581 model higgs boson with the atlas detector at the lh. *Phys. Lett. B*, 716:1–29, 2012.
1582 doi: 10.1016/j.physletb.2012.08.020.
- 1583 [12] CMS Collaboration. Observation of a new boson at a mass of 125 gev with the cms
1584 experiment at the lh. *Phys. Lett. B*, 716:30–61, 2012. doi: 10.1016/j.physletb.2012.
1585 08.021.
- 1586 [13] ATLAS and CMS Collaborations. Combined measurement of the higgs boson mass in
1587 pp collisions at $\sqrt{s} = 7$ and 8 tev with the atlas and cms experiments. *Phys. Rev. Lett.*,
1588 114:191803, 2015. doi: 10.1103/PhysRevLett.114.191803.

- [14] Nicola Cabibbo. Unitary symmetry and leptonic decays. *Phys. Rev. Lett.*, 10:531–533, 1963. doi: 10.1103/PhysRevLett.10.531.
- [15] Makoto Kobayashi and Toshihide Maskawa. Cp-violation in the renormalizable theory of weak interaction. *Prog. Theor. Phys.*, 49:652–657, 1973. doi: 10.1143/PTP.49.652.
- [16] Super-Kamiokande Collaboration. Evidence for oscillation of atmospheric neutrinos. *Phys. Rev. Lett.*, 81:1562–1567, 1998. doi: 10.1103/PhysRevLett.81.1562.
- [17] SNO Collaboration. Direct evidence for neutrino flavor transformation from neutral-current interactions in the sudbury neutrino observatory. *Phys. Rev. Lett.*, 89:011301, 2002. doi: 10.1103/PhysRevLett.89.011301.
- [18] Planck Collaboration. Planck 2018 results. vi. cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. doi: 10.1051/0004-6361/201833910.
- [19] A. D. Sakharov. Violation of cp invariance, c asymmetry, and baryon asymmetry of the universe. *Pisma Zh. Eksp. Teor. Fiz.*, 5:32–35, 1967.
- [20] M. B. Gavela, P. Hernandez, J. Orloff, and O. Pene. Standard model cp-violation and baryon asymmetry. *Mod. Phys. Lett. A*, 9:795–810, 1994. doi: 10.1142/S0217732394000629.
- [21] Andrew G. Cohen, D. B. Kaplan, and A. E. Nelson. Progress in electroweak baryogenesis. *Ann. Rev. Nucl. Part. Sci.*, 43:27–70, 1993. doi: 10.1146/annurev.ns.43.120193.000331.
- [22] C. A. Baker et al. An improved experimental limit on the electric dipole moment of the neutron. *Phys. Rev. Lett.*, 97:131801, 2006. doi: 10.1103/PhysRevLett.97.131801.
- [23] LHCb Collaboration. Test of lepton universality in $b \rightarrow s\ell^+\ell^-$ decays. *Phys. Rev. Lett.*, 131:051803, 2023. doi: 10.1103/PhysRevLett.131.051803.
- [24] LHCb Collaboration. Measurement of lepton universality parameters in $b^+ \rightarrow k^+\ell^+\ell^-$ and $b^0 \rightarrow k^{*0}\ell^+\ell^-$ decays. *Phys. Rev. D*, 108:032002, 2023. doi: 10.1103/PhysRevD.108.032002.
- [25] Heavy Flavor Averaging Group (HFLAV). Average of $R(D)$ and $R(D^*)$ for CKM 2025. <https://hflav-eos.web.cern.ch/hflav-eos/semi/ckm25/html/RDsDsstar/RDRDs.html>, 2025. Last update: 28 September 2025.
- [26] BaBar Collaboration. Evidence for an excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ decays. *Phys. Rev. Lett.*, 109:101802, 2012. doi: 10.1103/PhysRevLett.109.101802.
- [27] BaBar Collaboration. Measurement of an Excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ Decays and Implications for Charged Higgs Bosons. *Phys. Rev. D*, 88(7):072012, 2013. doi: 10.1103/PhysRevD.88.072012.
- [28] Belle Collaboration. Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ decays with hadronic tagging at Belle. *Phys. Rev. D*, 92(7):072014, 2015. doi: 10.1103/PhysRevD.92.072014.

- [29] Belle Collaboration. Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^* \tau^- \bar{\nu}_\tau$. *Phys. Rev. Lett.*, 118(21):211801, 2017. doi: 10.1103/PhysRevLett.118.211801.
- [30] Belle Collaboration. Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^* \tau^- \bar{\nu}_\tau$ with one-prong hadronic τ decays at Belle. *Phys. Rev. D*, 97(1):012004, 2018. doi: 10.1103/PhysRevD.97.012004.
- [31] Belle Collaboration. Measurement of $R(D)$ and $R(D^*)$ with a semileptonic tagging method. *Phys. Rev. Lett.*, 124(16):161803, 2020. doi: 10.1103/PhysRevLett.124.161803.
- [32] Belle-II Collaboration. Test of lepton flavor universality with a measurement of $R(D^*)$ using hadronic B tagging at the Belle II experiment. *Phys. Rev. D*, 110(7):072020, 2024. doi: 10.1103/PhysRevD.110.072020.
- [33] Belle-II Collaboration. Test of lepton flavor universality with measurements of $R(D^+)$ and $R(D^{*+})$ using semileptonic B tagging at the Belle II experiment. *Phys. Rev. D*, 112(3):032010, 2025. doi: 10.1103/fmn3-h8fy.
- [34] LHCb Collaboration. Measurement of the ratios of branching fractions $\mathcal{R}(D^*)$ and $\mathcal{R}(D^0)$. *Phys. Rev. Lett.*, 131:111802, 2023. doi: 10.1103/PhysRevLett.131.111802.
- [35] LHCb Collaboration. Test of lepton flavor universality using $B^0 \rightarrow D^{*-} \tau^+ \nu_\tau$ decays with hadronic τ channels. *Phys. Rev. D*, 108(1):012018, 2023. doi: 10.1103/PhysRevD.108.012018. [Erratum: *Phys.Rev.D* 109, 119902 (2024)].
- [36] LHCb Collaboration. Measurement of the Branching Fraction Ratios $R(D^+)$ and $R(D^{*+})$ Using Muonic τ Decays. *Phys. Rev. Lett.*, 134(6):061801, 2025. doi: 10.1103/PhysRevLett.134.061801.
- [37] J. Wess and B. Zumino. Supergauge transformations in four dimensions. *Nucl. Phys. B*, 70:39–50, 1974. doi: 10.1016/0550-3213(74)90355-1.
- [38] Stephen P. Martin. A supersymmetry primer. 1997.
- [39] Savas Dimopoulos and Howard Georgi. Softly broken supersymmetry and $su(5)$. *Nucl. Phys. B*, 193:150–162, 1981. doi: 10.1016/0550-3213(81)90522-8.
- [40] Glennys R. Farrar and Pierre Fayet. Phenomenology of the production, decay, and detection of new hadronic states associated with supersymmetry. *Phys. Lett. B*, 76: 575–579, 1978. doi: 10.1016/0370-2693(78)90858-4.
- [41] H. Goldberg. Constraint on the photino mass from cosmology. *Phys. Rev. Lett.*, 50: 1419–1422, 1983. doi: 10.1103/PhysRevLett.50.1419.
- [42] John R. Ellis, J. S. Hagelin, D. V. Nanopoulos, Keith A. Olive, and M. Srednicki. Supersymmetric relics from the big bang. *Nucl. Phys. B*, 238:453–476, 1984. doi: 10.1016/0550-3213(84)90461-9.
- [43] Savas Dimopoulos, Stuart Raby, and Frank Wilczek. Supersymmetry and the scale of unification. *Phys. Rev. D*, 24:1681–1683, 1981. doi: 10.1103/PhysRevD.24.1681.

- [44] Ugo Amaldi, Wim de Boer, and Hermann Fürstenau. Comparison of grand unified theories with electroweak and strong coupling constants measured at lep. *Phys. Lett. B*, 260:447–455, 1991. doi: 10.1016/0370-2693(91)91641-8.
- [45] ATLAS Collaboration. The quest to discover supersymmetry at the atlas experiment. *Phys. Rep.*, 1116:1–76, 2025. doi: 10.1016/j.physrep.2025.01.004.
- [46] R. D. Peccei and Helen R. Quinn. Cp conservation in the presence of pseudoparticles. *Phys. Rev. Lett.*, 38:1440–1443, 1977. doi: 10.1103/PhysRevLett.38.1440.
- [47] R. D. Peccei and Helen R. Quinn. Constraints imposed by cp conservation in the presence of pseudoparticles. *Phys. Rev. D*, 16:1791–1797, 1977. doi: 10.1103/PhysRevD.16.1791.
- [48] Steven Weinberg. A new light boson? *Phys. Rev. Lett.*, 40:223–226, 1978. doi: 10.1103/PhysRevLett.40.223.
- [49] Frank Wilczek. Problem of strong p and t invariance in the presence of instantons. *Phys. Rev. Lett.*, 40:279–282, 1978. doi: 10.1103/PhysRevLett.40.279.
- [50] Jihn E. Kim and Gianpaolo Carosi. Axions and the strong cp problem. *Rev. Mod. Phys.*, 82:557–601, 2010. doi: 10.1103/RevModPhys.82.557.
- [51] Jogesh C. Pati and Abdus Salam. Lepton number as the fourth color. *Phys. Rev. D*, 10:275–289, 1974. doi: 10.1103/PhysRevD.10.275.
- [52] Howard Georgi and Sheldon L. Glashow. Unity of all elementary-particle forces. *Phys. Rev. Lett.*, 32:438–441, 1974. doi: 10.1103/PhysRevLett.32.438.
- [53] I. Dorsner, S. Fajfer, A. Greljo, J. F. Kamenik, and N. Kosnik. Physics of leptoquarks in precision experiments and at particle colliders. *Phys. Rept.*, 641:1–68, 2016. doi: 10.1016/j.physrep.2016.06.001.
- [54] Claudia Cornella, Javier Fuentes-Martin, and Gino Isidori. Revisiting the vector leptoquark explanation of the b -physics anomalies. *JHEP*, 07:168, 2019. doi: 10.1007/JHEP07(2019)168.
- [55] Michael J. Baker, Javier Fuentes-Martin, Gino Isidori, and Matthias König. High- p_t signatures in vector-leptoquark models. *Eur. Phys. J. C*, 79:334, 2019. doi: 10.1140/epjc/s10052-019-6853-x.
- [56] Jason Aebischer, Gino Isidori, Marko Pesut, Ben A. Stefanek, and Felix Wilsch. Confronting the vector leptoquark hypothesis with new low- and high-energy data. *Eur. Phys. J. C*, 83:153, 2023. doi: 10.1140/epjc/s10052-023-11304-5.
- [57] ATLAS Collaboration. Search for leptoquarks decaying into the $b\tau$ final state in pp collisions at $\sqrt{s} = 13$ tev with the atlas detector. *JHEP*, 10:001, 2023. doi: 10.1007/JHEP10(2023)001.
- [58] CMS Collaboration. Search for singly and pair-produced leptoquarks coupling to third-generation fermions in proton-proton collisions at $\sqrt{s} = 13$ tev. *Phys. Lett. B*, 819:136446, 2021. doi: 10.1016/j.physletb.2021.136446.

- [59] CMS Collaboration. Search for a third-generation leptoquark coupled to a τ lepton and a b quark through single, pair, and nonresonant production in proton-proton collisions at $\sqrt{s} = 13$ tev. *JHEP*, 05:311, 2024. doi: 10.1007/JHEP05(2024)311.
- [60] ATLAS Collaboration. A measurement of the high-mass $\tau\bar{\tau}$ production cross-section at $\sqrt{s} = 13$ tev with the atlas detector and constraints on new particles and couplings. *JHEP*, 10:054, 2025. doi: 10.1007/JHEP10(2025)054.
- [61] CMS Collaboration. Searches for additional higgs bosons and for vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ tev. *JHEP*, 07:073, 2023. doi: 10.1007/JHEP07(2023)073.
- [62] ATLAS Collaboration. Leptoquark summary plots for scalar or vector models. Technical Report ATL-PHYS-PUB-2025-013, CERN, 2025.
- [63] CMS Collaboration. Search for new physics in the τ lepton plus missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 13$ tev. *JHEP*, 09:051, 2023. doi: 10.1007/JHEP09(2023)051.
- [64] Motoi Endo, Syuhei Iguro, Teppei Kitahara, Michihisa Takeuchi, and Ryoutaro Watanabe. Non-resonant new physics search at the lhc for the $b \rightarrow c\tau\nu$ anomalies. *JHEP*, 02:106, 2022. doi: 10.1007/JHEP02(2022)106.
- [65] CERN. About CERN. <https://home.cern/about/>, . Accessed: 2026-06-11.
- [66] Lyndon Evans and Philip Bryant. LHC Machine. *JINST*, 3:S08001, 2008. doi: 10.1088/1748-0221/3/08/S08001.
- [67] CERN. The Large Hadron Collider. <https://home.cern/science/accelerators/large-hadron-collider>, . Accessed: 2026-06-11.
- [68] Esma Mobs. The CERN accelerator complex - August 2018. Complexe des accélérateurs du CERN - Août 2018. CERN Document Server, 2018. URL <https://cds.cern.ch/record/2636343>. General Photo, CERN-GRAPHICS-2018-005.
- [69] Werner Herr and B. Muratori. Concept of luminosity. 2006. doi: 10.5170/CERN-2006-002.361. URL <https://cds.cern.ch/record/941318>.
- [70] ATLAS Collaboration. Luminosity determination in pp collisions at $\sqrt{s} = 13$ tev using the atlas detector at the lhc. *Eur. Phys. J. C*, 83:982, 2023. doi: 10.1140/epjc/s10052-023-11747-w.
- [71] ATLAS Collaboration. LHC Run 3: physics at record energy starts tomorrow. 2022. URL <https://atlas.cern/Updates/Press-Statement/LHC-Run3-Starts>. Accessed: 2026-06-11.
- [72] ATLAS Collaboration. ATLAS data quality operations and performance for 2015–2018 data-taking. *JINST*, 15:P04003, 2020. doi: 10.1088/1748-0221/15/04/P04003.
- [73] ATLAS Collaboration. The atlas experiment at the cern large hadron collider. *JINST*, 3:S08003, 2008. doi: 10.1088/1748-0221/3/08/S08003.

- [74] ATLAS Collaboration. The atlas experiment at the cern large hadron collider: A description of the detector configuration for run 3. *JINST*, 19:P05063, 2024. doi: 10.1088/1748-0221/19/05/P05063.
- [75] Fabian Kuger. Signal formation processes in micromegas detectors and quality control for large size detector construction for the atlas new small wheel, 2017.
- [76] B. Abbott et al. Production and integration of the atlas insertable b-layer. *JINST*, 13:T05008, 2018. doi: 10.1088/1748-0221/13/05/T05008.
- [77] ATLAS Collaboration. The Phase-I Trigger Readout Electronics Upgrade of the ATLAS Liquid Argon Calorimeters. *JINST*, 17(05):P05024, 2022. doi: 10.1088/1748-0221/17/05/P05024.
- [78] ATLAS Collaboration. Operation of the atlas trigger system in run 2. *JINST*, 15:P10004, 2020. doi: 10.1088/1748-0221/15/10/P10004.
- [79] Johan Alwall et al. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014. doi: 10.1007/JHEP07(2014)079.
- [80] Christian Bierlich et al. A comprehensive guide to the physics and usage of PYTHIA 8.3. *SciPost Phys. Codebases*, page 8, 2022. doi: 10.21468/SciPostPhysCodeb.8.
- [81] Richard D. Ball et al. Parton distributions for the LHC run II. *JHEP*, 04:040, 2015. doi: 10.1007/JHEP04(2015)040.
- [82] ATLAS Collaboration. ATLAS Pythia 8 tunes to 7 tev data. Technical Report ATL-PHYS-PUB-2014-021, CERN, 2014. URL <https://cds.cern.ch/record/1966419>.
- [83] Richard D. Ball et al. Parton distributions with LHC data. *Nucl. Phys. B*, 867:244–289, 2013. doi: 10.1016/j.nuclphysb.2012.10.003.
- [84] Leif Lönnblad and Stefan Prestel. Matching tree-level matrix elements with interleaved showers. *JHEP*, 03:019, 2012. doi: 10.1007/JHEP03(2012)019.
- [85] David J. Lange. The EvtGen particle decay simulation package. *Nucl. Instrum. Meth. A*, 462:152–155, 2001. doi: 10.1016/S0168-9002(01)00089-4.
- [86] ATLAS Collaboration. AtlFast3: The next generation of fast simulation in ATLAS. *Comput. Softw. Big Sci.*, 6:7, 2022. doi: 10.1007/s41781-021-00079-7.
- [87] Enrico Bothmann et al. Event generation with Sherpa 2.2. *SciPost Phys.*, 7:034, 2019. doi: 10.21468/SciPostPhys.7.3.034.
- [88] Steffen Schumann and Frank Krauss. A parton shower algorithm based on Catani–Seymour dipole factorisation. *JHEP*, 03:038, 2008. doi: 10.1088/1126-6708/2008/03/038.
- [89] Stefan Höche, Frank Krauss, Marek Schönherr, and Frank Siegert. A critical appraisal of NLO+PS matching methods. *JHEP*, 09:049, 2012. doi: 10.1007/JHEP09(2012)049.

- [90] Stefan Höche, Frank Krauss, Marek Schönherr, and Frank Siegert. QCD matrix elements + parton showers. the NLO case. *JHEP*, 04:027, 2013. doi: 10.1007/JHEP04(2013)027.
- [91] Stefano Catani, Frank Krauss, Bryan R. Webber, and Ralf Kuhn. QCD matrix elements + parton showers. *JHEP*, 11:063, 2001. doi: 10.1088/1126-6708/2001/11/063.
- [92] Stefan Höche, Frank Krauss, Steffen Schumann, and Frank Siegert. QCD matrix elements and truncated showers. *JHEP*, 05:053, 2009. doi: 10.1088/1126-6708/2009/05/053.
- [93] Stefano Frixione, Paolo Nason, and Giovanni Ridolfi. A positive-weight next-to-leading-order monte carlo for heavy flavour hadroproduction. *JHEP*, 09:126, 2007. doi: 10.1088/1126-6708/2007/09/126.
- [94] Paolo Nason. A new method for combining NLO QCD with shower monte carlo algorithms. *JHEP*, 11:040, 2004. doi: 10.1088/1126-6708/2004/11/040.
- [95] Stefano Frixione, Paolo Nason, and Carlo Oleari. Matching NLO QCD computations with parton shower simulations: the POWHEG method. *JHEP*, 11:070, 2007. doi: 10.1088/1126-6708/2007/11/070.
- [96] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. A general framework for implementing NLO calculations in shower monte carlo programs: the POWHEG BOX. *JHEP*, 06:043, 2010. doi: 10.1007/JHEP06(2010)043.
- [97] Torbjörn Sjöstrand et al. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.*, 191:159–177, 2015. doi: 10.1016/j.cpc.2015.01.024.
- [98] Michał Czakon and Alexander Mitov. Top++: A program for the calculation of the top-pair cross-section at hadron colliders. *Comput. Phys. Commun.*, 185:2930–2938, 2014. doi: 10.1016/j.cpc.2014.06.021.
- [99] ATLAS Collaboration. The ATLAS simulation infrastructure. *Eur. Phys. J. C*, 70: 823–874, 2010. doi: 10.1140/epjc/s10052-010-1429-9.
- [100] S. Agostinelli et al. Geant4—a simulation toolkit. *Nucl. Instrum. Meth. A*, 506: 250–303, 2003. doi: 10.1016/S0168-9002(03)01368-8.
- [101] ATLAS Collaboration. Performance of electron and photon triggers in ATLAS during LHC run 2. *Eur. Phys. J. C*, 80:47, 2020. doi: 10.1140/epjc/s10052-019-7500-2.
- [102] ATLAS Collaboration. Performance of the ATLAS muon triggers in run 2. *JINST*, 15: P09015, 2020. doi: 10.1088/1748-0221/15/09/P09015.
- [103] ATLAS Collaboration. Performance of the missing transverse momentum triggers for the ATLAS detector during run-2 data taking. *JHEP*, 08:080, 2020. doi: 10.1007/JHEP08(2020)080.
- [104] ATLAS Collaboration. Formulae for estimating significance. ATL-PHYS-PUB-2020-025, 2020. URL <https://cds.cern.ch/record/2736148>.

1812 **Appendix A**

1813 **Correction of the signal internal event** 1814 **weights**

1815 For the right-handed interpretation, the dependence of the signal prediction on g_U and
1816 β_L^{23} is obtained by reweighting the baseline sample generated at each M_U value. The
1817 MADGRAPH5_AMC@NLO internal event weights encode the change in the matrix element
1818 when these coupling parameters are varied. Their performance is evaluated by comparing the
1819 reweighted predictions with samples generated independently at selected parameter points.

1820 The internal event weights alone do not fully reproduce the reference predictions in
1821 regions where the relative contributions of resonant and non-resonant production change. The
1822 overall normalisation is therefore adjusted to that of the corresponding reference prediction.
1823 A bin-by-bin two-dimensional correction in the leading-jet p_T and $m_{b\tau}$ is then applied to
1824 account for the remaining shape differences. These observables provide sensitivity to the
1825 resonant and non-resonant components and allow both contributions to be corrected within
1826 the same sample.

1827 The corrected predictions are validated using truth-level selections that reproduce the main
1828 signal-region requirements. At each stage of the selection, the corrected and independently
1829 generated event yields agree within 10% for both the resonant and non-resonant signal
1830 regions. This difference is smaller than the theoretical uncertainties assigned to the signal
1831 prediction.

1832 Appendix B

1833 Multijet background study

1834 The multijet suppression used in Section 7.1.2 is studied with simulated dijet and leptoquark
1835 events. The purpose is to identify a selection that removes events with instrumental E_T^{miss}
1836 while retaining the signal with neutrinos. The multijet production cross section is large, so
1837 a small probability for a jet to be misidentified as a τ , combined with a large jet-energy
1838 mismeasurement, can produce events with the selected final state.

1839 B.1 Origin of the discriminating variables

1840 Jet-momentum mismeasurement produces a characteristic direction for the apparent missing
1841 momentum. For example, if the transverse momentum of one jet is underestimated by $\delta \vec{p}_T^j$,
1842 the reconstructed momentum imbalance receives a contribution in the direction of the missing
1843 part of that jet momentum. The resulting $\vec{p}_T^{\text{miss}}$ is therefore approximately aligned with the
1844 jet. In a signal event, the neutrinos produce real missing momentum, whose direction is not
1845 fixed by a jet-energy mismeasurement.

1846 The variables $\Delta\phi_{\text{min}}$ and $R_{\text{miss}/j}$ defined in Eq. 7.3 test these two features. The first
1847 measures the angular alignment, and the second compares the missing momentum with the
1848 momentum of the closest jet in azimuth. The combination is useful because a low- p_T jet can
1849 lie close to the direction of real missing momentum in a signal event. Such an event can have
1850 small $\Delta\phi_{\text{min}}$ but large $R_{\text{miss}/j}$.

1851 B.2 Study selection and distributions

1852 The study selection is kept loose to retain enough simulated multijet events to examine the
1853 relevant distributions. It requires the E_T^{miss} trigger, exactly one hadronic τ candidate, no

selected electron or muon, and at least one jet, as listed in Table B.1. The final SR thresholds on E_T^{miss} , p_T^τ , m_T and the visible invariant mass are not imposed at this stage. No b -tag multiplicity is required.

Table B.1 Loose selection used to study the multijet suppression. This selection defines the denominator of the rejection fractions shown below.

Requirement	Selection
Trigger	Lowest-threshold unprescaled E_T^{miss} trigger
Hadronic τ multiplicity	$N_\tau = 1$
Electron and muon veto	$N_\ell = 0$
Jet multiplicity	$N_j \geq 1$

Figure B.1 shows the two-dimensional distributions after this selection. The multijet sample is concentrated at small $\Delta\phi_{\text{min}}$ and relatively small $R_{\text{miss}/j}$. The signal occupies a broader angular range and extends to larger momentum ratios. Approximately 90% of the multijet events have $R_{\text{miss}/j} < 6$, compared with approximately 55% of the signal in the study sample. The ratio requirement alone would therefore reject a substantial fraction of the signal. It is instead combined with the angular requirement so that either a large angle or a large ratio is sufficient to retain an event.




Fig. B.1 Correlation between the minimum jet- $\vec{p}_T^{\text{miss}}$ azimuthal separation and the ratio of E_T^{miss} to the p_T of the closest jet in azimuth, for (a) simulated signal with $M_U = 1.5$ TeV and (b) simulated multijet background after the loose study selection. The concentration of multijet events at small values of both variables motivates a combined requirement.

B.3 Threshold scan and signal retention

The angular threshold is scanned while the momentum-ratio threshold is fixed at six. For an angular threshold X , the rejected fraction is defined as

$$f_{\text{rej}}(X) = \frac{N(\Delta\phi_{\text{min}} < X \text{ and } R_{\text{miss}/j} < 6)}{N(\text{loose study selection})}. \quad (\text{B.1})$$

The numerator requires both small angular separation and a small momentum ratio. Its complement is the accepted region, defined by the logical OR in Eq. 7.4. The rejected fraction is evaluated separately for signal and multijet events, using each sample's yield after the loose selection as its denominator.

Figure B.2 shows the resulting rejection fractions. At $X = 0.4$, approximately 86% of the multijet events are rejected, while the signal loss is 3.6%. The corresponding signal efficiency for the cleaning cut is therefore $1 - 0.036 = 96.4\%$. Increasing the angular threshold rejects more multijet events but also removes more signal. The selected threshold provides substantial multijet rejection with a small signal loss in the studied sample.




Fig. B.2 Rejected fractions of simulated signal and multijet events as a function of the minimum jet- $\vec{p}_T^{\text{miss}}$ angular threshold X , with $R_{\text{miss}/j} < 6$ also required in the rejected sample. The red and black curves show signal loss and multijet rejection, respectively. The vertical dotted line marks the adopted threshold $X = 0.4$. The denominator is the loose selection in Table B.1.

B.4 Residual multijet contribution

The residual multijet contribution is examined after applying the cleaning cut and the full SR requirements. No simulated multijet events survive those selections in the samples used for the study. This supports neglecting the multijet contribution in the nominal SR prediction, but zero surviving simulated events does not establish an exactly zero physical rate or a quantitative upper bound without the effective sample size.

Jets misidentified as τ candidates can still occur in other background processes. In particular, $Z \rightarrow \nu\nu$ production with jets contains genuine missing momentum and can pass the multijet cleaning even when the selected τ candidate is a jet. This contribution is included in the simulated Z+jets background. A supplementary selection requiring Medium τ identification while failing Tight also gives a negligible multijet contribution in the simulated study, supporting the effectiveness of the cleaning beyond the nominal Tight selection.

Appendix C

Zero- b -jet signal-region optimisation

The E_T^{miss} threshold in SR0b is studied to assess whether tighter selections can improve the expected sensitivity at large β_L^{23} . Removing the b -tag requirement admits a larger SM background than in SR1b and increases the relevance of interference between the leptoquark and SM amplitudes. A tighter E_T^{miss} selection can change the balance between the retained signal, background and interference contributions. The study therefore examines the significance as the lower E_T^{miss} threshold is varied.

The threshold is scanned from 200 to 1500 GeV in 100 GeV steps. The signal parameter points are $(M_U, g_U, \beta_L^{23}) = (1.5 \text{ TeV}, 0.5, 1.8)$ for SR0b-Res and $(3.0 \text{ TeV}, 2.0, 1.8)$ for SR0b-NonRes. The large value of β_L^{23} makes these benchmarks relevant to the zero- b -jet categories discussed in Section 7.3. The scan outcomes and adopted thresholds are summarised in Table C.1.

Table C.1 Signal benchmarks and preferred thresholds in the supplementary SR0b E_T^{miss} scan. The last column gives the thresholds used in the final analysis.

Region	$(M_U [\text{TeV}], g_U, \beta_L^{23})$	Scan maximum	Adopted E_T^{miss} cut
SR0b-Res	(1.5, 0.5, 1.8)	600 GeV	$> 200 \text{ GeV}$
SR0b-NonRes	(3.0, 2.0, 1.8)	400 GeV	$> 400 \text{ GeV}$

For SR0b-NonRes, the significance is maximised at $E_T^{\text{miss}} > 400 \text{ GeV}$. The scan therefore supports retaining the threshold used in the final region definition. For SR0b-Res, the maximum occurs at $E_T^{\text{miss}} > 600 \text{ GeV}$, giving an improvement of approximately 20% in the significance at the chosen benchmark. The final resonant selection nevertheless retains $E_T^{\text{miss}} > 200 \text{ GeV}$ to keep its kinematic requirements aligned with SR1b-Res.

