

Variable	Description
n_{PV}	Number of primary vertices in the event.
μ	Average number of interactions per bunch crossing.
$\langle \lambda_{\text{centre}} \rangle$	Mean distance of the cluster shower centre from the calorimeter front face measured along the shower axis
$\langle \lambda^2 \rangle$	Mean distance of a cell from the shower centre along the shower axis
$\langle \rho^1 \rangle$	Mean of the first moment in the cluster energy density, $\rho = E^{\text{cluster}}/V^{\text{cluster}}$, where E^{cluster} and V^{cluster} represent the energy and volume of the cluster, respectively
$f_{\text{presampler}}$	Mean fraction of cluster energy deposited in the barrel and endcap presamplers
P_{EM}	Classification probability of the cluster to be EM-like, as described in Ref. [45]
n_{track}	Number of tracks associated with the $\tau_{\text{had-vis}}$
n_{π^0}	Number of reconstructed neutral pions associated with the $\tau_{\text{had-vis}}$
γ_{π}	Relative difference of the total charged pion energy, E_{charged} , and the total neutral pion energy, E_{neutral} : $\gamma_{\pi} = (E_{\text{charged}} - E_{\text{neutral}})/(E_{\text{charged}} + E_{\text{neutral}})$
$\eta_{\text{calo}}^{\text{Comb}}$	Calorimeter-based (baseline) pseudorapidity
p_{T}	Transverse momentum interpolated from calorimetric corrections to energy measurement and TPF reconstruction
$p_{\text{T}}^{\text{EM}}/p_{\text{T}}^{\text{LC}}$	Ratio of the calorimetric energy at the EM scale to the local hadron calibration transverse momentum p_{T}^{LC}
$p_{\text{T}}^{\text{LC}}/p_{\text{T}}^{\text{Comb}}$	Ratio of the local hadron calibration transverse momentum to $p_{\text{T}}^{\text{Comb}}$
$p_{\text{T}}^{\text{TPF}}/p_{\text{T}}^{\text{Comb}}$	Ratio of the TPF reconstruction transverse momentum, $p_{\text{T}}^{\text{TPF}}$, to $p_{\text{T}}^{\text{Comb}}$
TPF BDT Values	BDT values of the TPF algorithm for 1p0n vs 1p1n, 1p1n vs 1pXn, and 3p0n vs 3pXn discrimination