

Solution Manual to
A Course in Mathematical Statistics
and Large Sample Theory

Kisung You

kyou@nd.edu

Contents

Chapter 1.	Introduction	3
Chapter 2.	Decision Theory	4
Chapter 3.	Introduction to General Methods of Estimation	5
Chapter 4.	Sufficient Statistics, Exponential Families, and Estimation	6
Chapter 5.	Testing Hypotheses	7
Chapter 6.	Consistency and Asymptotic Distributions of Statistics	8

CHAPTER 1

Introduction

EX. 1.1. From a population of size N a simple random sample of size n is drawn *without replacement*, and a real-valued characteristic X measured to yield observations X_j ($j = 1, 2, \dots, n$). Show that

- (a) The way we deal
- (b) the expected squared error of \bar{X} as an estimator of m , i.e., the variance of \bar{X} , is smaller than that of the mean of a simple random sample of the same size n drawn *with replacement*, and
- (c) the difference between the expected squared errors of the two estimators is $O(n/N)$, as n/N goes to zero.

CHAPTER 2

Decision Theory

EX. 2.1. From a population of size N a simple random sample of size n is drawn *without replacement*, and a real-valued characteristic X measured to yield observations X_j ($j = 1, 2, \dots, n$). Show that

- (a) The way we deal
- (b) the expected squared error of \bar{X} as an estimator of m , i.e., the variance of \bar{X} , is smaller than that of the mean of a simple random sample of the same size n drawn *with replacement*, and
- (c) the difference between the expected squared errors of the two estimators is $O(n/N)$, as n/N goes to zero.

CHAPTER 3

Introduction to General Methods of Estimation

EX. 3.1. From a population of size N a simple random sample of size n is drawn *without replacement*, and a real-valued characteristic X measured to yield observations X_j ($j = 1, 2, \dots, n$). Show that

- (a) The way we deal
- (b) the expected squared error of \bar{X} as an estimator of m , i.e., the variance of \bar{X} , is smaller than that of the mean of a simple random sample of the same size n drawn *with replacement*, and
- (c) the difference between the expected squared errors of the two estimators is $O(n/N)$, as n/N goes to zero.

CHAPTER 4

Sufficient Statistics, Exponential Families, and Estimation

EX. 4.1. From a population of size N a simple random sample of size n is drawn *without replacement*, and a real-valued characteristic X measured to yield observations X_j ($j = 1, 2, \dots, n$). Show that

- (a) The way we deal
- (b) the expected squared error of \bar{X} as an estimator of m , i.e., the variance of \bar{X} , is smaller than that of the mean of a simple random sample of the same size n drawn *with replacement*, and
- (c) the difference between the expected squared errors of the two estimators is $O(n/N)$, as n/N goes to zero.

CHAPTER 5

Testing Hypotheses

EX. 5.1. From a population of size N a simple random sample of size n is drawn *without replacement*, and a real-valued characteristic X measured to yield observations X_j ($j = 1, 2, \dots, n$). Show that

- (a) The way we deal
- (b) the expected squared error of \bar{X} as an estimator of m , i.e., the variance of \bar{X} , is smaller than that of the mean of a simple random sample of the same size n drawn *with replacement*, and
- (c) the difference between the expected squared errors of the two estimators is $O(n/N)$, as n/N goes to zero.

CHAPTER 6

Consistency and Asymptotic Distributions of Statistics

EX. 6.1. **ERRATA**

- (a)
- (b) errata cascades

EX. 6.2. (a)

- (b) errata cascades