

Algebraic Topology II Homework 2

James Harbour

September 6, 2023

1 Problem 1

Assuming as known the cup product structure on the 2-torus $\mathbb{T}^2 = S^1 \times S^1$, compute the cup product structure in $H^*(M_g)$ for M_g the closed orientable surface of genus g using the quotient map from M_g to a wedge sum of g tori (see photo).

Proof. We take as known that the cohomology ring of \mathbb{T}^2 with coefficients in a ring R is given by

$$H^*(\mathbb{T}^2; R) = \Lambda_R[\alpha_1, \alpha_2]$$

$$\begin{array}{ccc} H^1(\bigvee_g \mathbb{T}^2) \oplus H^1(\bigvee_g \mathbb{T}^2) & \longrightarrow & H^2(\bigvee_g \mathbb{T}^2) \\ \downarrow & & \downarrow \\ H^1(M_g) \oplus H^1(M_g) & \longrightarrow & H^2(M_g) \end{array}$$

□

2 Problem 3

(a): Using the cup product structure, show that there is no map $\mathbb{R}P^n \rightarrow \mathbb{R}P^m$ inducing a nontrivial map $H^1(\mathbb{R}P^m; \mathbb{Z}_2) \rightarrow H^1(\mathbb{R}P^n; \mathbb{Z}_2)$ if $n > m$. What is corresponding result for maps $\mathbb{C}P^n \rightarrow \mathbb{C}P^m$.

(b): Prove the Borsuk-Ulam theorem.

3 Problem 7

Use cup products to show that $\mathbb{R}P^3$ is not homotopy equivalent to $\mathbb{R}P^2 \vee S^1$.

4 Problem 8

Let X be $\mathbb{C}P^2$ with a cell e^3 attached by a map $S^2 \rightarrow \mathbb{C}P^1 \subseteq \mathbb{C}P^2$ of degree p , and let $Y := M(\mathbb{Z}_p, 2) \vee S^4$. Thus X and Y have the same 3-skeleton but differ in the way their 4-cells are attached. Show that X and Y have isomorphic cohomology rings with coefficients in \mathbb{Z} , but not with \mathbb{Z}_p coefficients.

5 Problem 9

Show that if $H_n(X; \mathbb{Z})$ free for each n , then $H^*(X; \mathbb{Z}_p)$ and $H^*(X; \mathbb{Z}) \otimes \mathbb{Z}_p$ are isomorphic as rings, so in particular the ring structure with \mathbb{Z} coefficients determines the ring structure with \mathbb{Z}_p coefficients.