

**Reading:**

- For this homework: 1.3-1.4
- For Wednesday, February 2: 1.5 up to Theorem 1.11
- For Monday, February 7 : 2.1

**Problem 1.**

Folland Chapter 1, Problem 11

**Problem 2.**

Folland Chapter 1, Problem 12 (we will later see that this metric space is complete).

**Problem 3.**

Folland Chapter 1, Problem 23

**Problem 4.**

Let  $\mathcal{A}$  be an algebra, and let  $\mu: \mathcal{A} \rightarrow [0, +\infty]$  be a finitely additive measure.

- (i) Suppose  $(A_j)_{j=1}^\infty$  are pairwise disjoint subsets of  $\mathcal{A}$ , and that  $A = \bigcup_{j=1}^\infty A_j \in \mathcal{A}$ . Show that

$$\mu(A) \geq \sum_{j=1}^\infty \mu(A_j).$$

- (ii) Show that the following are equivalent:

- $\mu$  is a premeasure,
- $\mu\left(\bigcup_{j=1}^\infty A_j\right) \leq \sum_{j=1}^\infty \mu(A_j)$  for any sequence  $(A_j)_{j=1}^\infty$  with  $\bigcup_{j=1}^\infty A_j \in \mathcal{A}$ ,
- for any increasing sequence  $(E_j)_{j=1}^\infty$  in  $\mathcal{A}$  with  $\bigcup_{j=1}^\infty E_j \in \mathcal{A}$ , we have

$$\mu\left(\bigcup_j E_j\right) = \lim_{n \rightarrow \infty} \mu(E_n).$$

- (iii) If  $\mu(X) < +\infty$ , show that  $\mu$  is a premeasure if and only if for every decreasing sequence  $(E_n)_{n=1}^\infty$  of sets with  $\bigcap_{n=1}^\infty E_n = \emptyset$ , we have

$$\lim_{n \rightarrow \infty} \mu(E_n) = 0.$$

**Problem 5.**

A *metric measure space* is a triple  $(X, d, \mu)$  where  $(X, d)$  is a metric space and  $\mu: \mathcal{B}_{(X, d)} \rightarrow [0, +\infty]$  is a measure. We say that  $E \subseteq X$  is a *continuity set*, if  $\mu(\overline{E} \setminus \text{Int}(E)) = 0$ . Here  $\text{Int}(E)$  is the interior of  $E$ . For this problem, fix a metric measure space  $(X, d, \mu)$ .

- Show that the collection of continuity sets forms an algebra of sets.
- Show that if  $x \in X$ ,  $r > 0$  and  $\mu(B_r(x, d)) < +\infty$ , then there is an  $s \in (0, r)$  so that  $B_s(x, d)$  is a continuity set.
- Suppose that  $(X, d)$  is separable and that for every  $x \in X$ , there is an  $r > 0$  so that  $\mu(B_r(x, d)) < +\infty$ . Show that there is a countable basis consisting of open continuity sets. (Hint: given a countable, dense  $D \subseteq X$  and  $x \in D$ , use the preceding part to choose a countable set  $J_x \subseteq (0, +\infty)$  with the property that  $\inf_{t \in J_x} t = 0$  and so that  $B_t(x, d)$  is a continuity set for all  $t \in J_x$ ).

**Problem 6.**

Let  $(X, d)$  be a metric space and  $\mu, \nu$  be finite, Borel measures on  $X$  with  $\mu(X) = \nu(X)$ . Let  $\mathcal{A} = \{E \in \mathcal{B}_{(X,d)} : \mu(E) = \nu(E)\}$ .

- (i) Show that if  $F \subseteq E$  and  $F, E \in \mathcal{A}$ , then  $E \setminus F \in \mathcal{A}$ . Also show that if  $(E_n)_{n=1}^\infty$  is an increasing sequence of elements of  $\mathcal{A}$ , then  $\bigcup_{n=1}^\infty E_n \in \mathcal{A}$ .
- (ii) Given a nonempty  $F \subseteq X$  closed, and  $x \in X$ , define  $d(x, F) = \inf_{y \in F} d(x, y)$ . Show that  $x \mapsto d(x, F)$  is continuous and  $F = \{x \in X : d(x, F) = 0\}$ .
- (iii) Show that  $\{U \subseteq X : U \text{ is open}\} \subseteq \mathcal{A}$  if and only if  $\{F \subseteq X : F \text{ is closed}\} \subseteq \mathcal{A}$ . (Hint: use the metric to write a closed subset of  $X$  as a countable intersection of open sets. Similarly, use the metric to write an open set as a countable union of closed sets).

We'll see later that either of the conditions in (iii) imply that  $\mathcal{A} = \mathcal{B}_X$ .

**Problems to think about, do not turn in:**

**Problem 7.**

Folland Chapter 1, Problem 3. (Problem 4 on the last homework, and Folland Chapter 1, Problem 5 might be helpful).

**Problem 8.**

Suppose that  $X$  is a set, that  $\mathcal{A} \subseteq \mathcal{P}(X)$  is an algebra and that  $\mu: \mathcal{A} \rightarrow [0, +\infty]$  is a finitely additive measure with  $\mu(X) = 1$ . Suppose that  $T_j: X \rightarrow X, j = 1, \dots, k$  are bijections so that  $\mathcal{A} = \{T_j(A) : A \in \mathcal{A}\}$  for all  $j = 1, \dots, k$ . Suppose that there exists  $B \subseteq X$  with  $0 < \mu(B) < +\infty$ , an integer  $1 \leq s \leq k$ , and sets  $A_1, \dots, A_k \in \mathcal{A}$  so that

$$B \supseteq \bigcup_{i=1}^k A_i,$$

and so that

$$B \sqcup T_s(B) \subseteq \bigcup_{i=1}^k T_i(A_i).$$

Show that there is an integer  $1 \leq l \leq k$ , and an  $A \in \mathcal{A}$  with  $\mu(T_l(A)) \neq \mu(A)$ .

(Note: the existence of such a set where  $B$  is a unit ball in  $\mathbb{R}^3$ ,  $\mu$  is a finitely additive measure defined on  $\mathcal{P}(\mathbb{R}^3)$  extending Lebesgue measure, and  $T_1, \dots, T_k$  are isometries is the Banach-Tarski paradox).