

MATH 7752 Homework 1

James Harbour

January 25, 2022

Problem 1

Let R be a ring and M an R -module.

(a) Prove that for every $m \in M$, the map $r \mapsto rm$ from R to M is a homomorphism of R -modules.

Proof. Fix $m \in M$ and let φ denote the map $r \mapsto rm$. Fix $x, y \in R$ and $r \in R$. Observe that

$$\varphi(x + y) = (x + y)m = xm + ym = \varphi(x) + \varphi(y)$$

and

$$\varphi(rx) = (rx)m = r(xm) = r\varphi(x),$$

so φ is an R -module homomorphism. □

(b) Assume that R is commutative and M an R -module. Prove that there is an isomorphism $\text{Hom}_R(R, M) \simeq M$ as R -modules.

Proof. For $m \in M$, let φ_m denote the R -module homomorphism in part (a).

Consider the map $\psi : M \rightarrow \text{Hom}_R(R, M)$ given by $\psi(m) = \varphi_m$. For $m, n \in M$ and $r, x \in R$,

$$\psi(m + n)(x) = \varphi_{m+n}(x) = x(m + n) = xm + xn = \varphi_m(x) + \varphi_n(x) = (\psi(m) + \psi(n))(x)$$

so $\psi(m + n) = \psi(m) + \psi(n)$, and

$$\psi(rm)(x) = \varphi_{rm}(x) = x(rm) = r(xm) = r\varphi_m(x) = (r\psi(m))(x)$$

so $\psi(rm) = r\psi(m)$.

Suppose $\psi(m) = \psi(n)$. Then $m = \varphi_m(1) = \psi(m)(1) = \psi(n)(1) = \varphi_n(1) = n$, so ψ is injective.

Suppose $\varphi \in \text{Hom}_R(R, M)$. For $r \in R$,

$$\psi_{\varphi(1)}(r) = r\varphi(1) = \varphi(r),$$

so $\psi_{\varphi(1)} = \varphi$, i.e. ψ is surjective. □

Problem 2

Give an explicit example of a map $f : A \rightarrow B$ with the following properties:

- A, B are R -modules.
- f is a group homomorphism.
- f is not an R -module homomorphism.

Solution. Consider $A = B = \mathbb{C}$ viewed as \mathbb{C} -modules over themselves. Let $f : A \rightarrow B$ be complex conjugation. For $z, w \in A$, $f(z + w) = \overline{z + w} = \bar{z} + \bar{w} = f(z) + f(w)$, so f is a group homomorphism. However, for $z \in A \setminus \{0\}$, $f(iz) = -i\bar{z} \neq i\bar{z} = if(z)$, so f is not an R -module homomorphism. \square

Problem 3

Let R be a ring and M an R -module.

(a) Let N be a subset of M . The *annihilator* of N is defined to be the set

$$\text{Ann}_R(N) := \{r \in R : rn = 0, \text{ for all } n \in N\}.$$

Prove that $\text{Ann}_R(N)$ is a left ideal of R .

Proof. Let $x, y \in I$ and $r \in R$. Fix $n \in N$. Noting that $xn = 0 = yn$, it follows that

$$(x + ry)n = xn + (ry)n = xn + r(yn) = 0.$$

Thus $x + ry \in \text{Ann}_R(N)$. Since all elements chosen were arbitrary, $\text{Ann}_R(N)$ is a left ideal of R . \square

(b) Show that if N is an R -submodule of M , then $\text{Ann}_R(N)$ is an ideal of R (i.e. it is two-sided ideal).

Proof. By part (a), it suffices to show that $\text{Ann}_R(N)$ is a right ideal of R . Moreover, part (a) shows *a fortiori* that $\text{Ann}_R(N)$ is already an abelian group, so we need only address its multiplicative structure. Let $y \in \text{Ann}_R(N)$ and $r \in R$. Fix $n \in N$. As N is an R -submodule of M , $yn \in N$, whence $(yr)n = y(rn) = 0$ by definition. Hence $\text{Ann}_R(N)$ is a two-sided ideal of R . \square

(c) For a subset I of R the *annihilator* of I in M is defined to be the set,

$$\text{Ann}_M(I) := \{m \in M : xm = 0, \text{ for all } x \in I\}.$$

Find a natural condition on I that guarantees that $\text{Ann}_M(I)$ is a submodule of M .

Claim. $\text{Ann}_M(I)$ is an R -submodule of M if I is a right ideal of R .

Proof. Suppose I is a right ideal of R . As $x \cdot 0 = 0$ for all $x \in I$, $\text{Ann}_M(I) \neq \emptyset$. Suppose $m, n \in \text{Ann}_M(I)$ and $r \in R$. Fix $x \in I$. By definition $x \cdot m = 0$. As I is a right ideal, $xr \in I$, so $x \cdot (m + r \cdot n) = x \cdot m + (xr) \cdot n = 0$. Thus $\text{Ann}_M(I)$ is an R -submodule of M . \square

(d) Let R be an integral domain. Prove that every finitely generated torsion R -module has a nonzero annihilator.

Proof. Let M be a finitely generated torsion R -module. Taking a generating set $m_1, \dots, m_n \in M$ of M , for each $k \in \{1, \dots, n\}$ there exists an $x_k \in R^\times = R \setminus \{0\}$ such that $x_k m_k = 0$. As R^\times is closed under multiplication, $r := x_1 \cdots x_n \in R^\times$ whence $r \neq 0$.

Now suppose that $m \in M$. Then there exist $r_1, \dots, r_n \in R$ such that $m = r_1 m_1 + \cdots + r_n m_n$. Observe that, by the commutativity of R ,

$$rm = (x_1 \cdots x_n)(r_1 m_1 + \cdots + r_n m_n) = \sum_{k=1}^n \left(\prod_{i \neq k} x_i \right) (x_k m_k) = 0.$$

Thus $0 \neq r \in \text{Ann}_R(M)$, so M has nonzero annihilator. □

Problem 4

In class we obtained a simple characterization of R -modules when $R = \mathbb{Z}$, and $R = F[x]$, with F a field. Imitate the method to find similar characterizations for R -modules in the following cases:

(a) $R = \mathbb{Z}/n\mathbb{Z}$, for some $n \geq 2$.

Proof. □

(b) $R = \mathbb{Z}[x]$.

(c) $R = F[x, y]$.

Problem 5

An R -module M is called *simple* (or *irreducible*) if its only submodules are $\{0\}$ and M . An R -module M is called *indecomposable* if M is not isomorphic to $N \oplus Q$ for some non-zero submodules N, Q . Show that every simple R -module is indecomposable, but the converse is not true.

Proof. Let M be a simple R -module. Then $M \neq 0$ as otherwise there would only be one submodule. Suppose, for the sake of contradiction, that M is not indecomposable. Then there exist some nonzero submodules $N, Q \subseteq M$ such that $M \cong N \oplus Q$. By simplicity of M , it follows that $N, Q = M$. But then $M \cong M \oplus M$. Moreover, $0 \oplus M$ is then a nonzero proper submodule of $M \oplus M$, whence via the isomorphism $M \oplus M \cong M$ we obtain a nonzero proper submodule of M , contradicting the simplicity of M .

To see that the converse does not hold, consider $R = \mathbb{Z}$ and $M = \mathbb{Z}$ considered as a \mathbb{Z} -module over itself. Note that $2\mathbb{Z} \subseteq \mathbb{Z}$ is a nonzero proper submodule of \mathbb{Z} , so M is not simple. To see that M is indecomposable, note that all nonzero submodules of M are of the form $a\mathbb{Z}$ for some $a \in \mathbb{Z} \setminus \{0\}$, and for any $a, b \in \mathbb{Z} \setminus \{0\}$, $ab \in a\mathbb{Z} \cap b\mathbb{Z}$. Hence, no sum of the required form would be direct. □

Problem 6

Let R be a ring. An R -module M is called *cyclic* if it is generated as an R -module by a single element.

(a) Prove that every cyclic R -module is of the form R/I for some left ideal I of R .

Proof. Let M be a cyclic R -module. Then there exists an $m \in M$ such that $M = Rm$. Consider the map $\varphi : R \rightarrow M$ given by $\varphi(r) = rm$ for $r \in R$. By problem 1 part (a), φ is an R -module homomorphism; moreover, φ is surjective since m generates M . Let $I = \ker(\varphi)$, a left ideal of R (actually two-sided, but we are identifying R with its left regular module over itself so a priori I is just a left R -submodule). Then, by the first isomorphism theorem, $M = \varphi(R) \cong R/\ker(\varphi) = R/I$. \square

(b) Show that the simple R -modules are precisely the ones which are isomorphic to R/\mathfrak{m} for some maximal left ideal \mathfrak{m} .

Proof. On one hand, \mathfrak{m} be a maximal left ideal of R . By the correspondence theorem applied to the natural projection, the only R -submodules of R/\mathfrak{m} are $\{0\}$ and R/\mathfrak{m} , so R/\mathfrak{m} is simple (and so is every R -module isomorphic to it).

On the other hand, suppose M is a nonzero simple R -module. Take $m \in M \setminus \{0\}$. Then by the simplicity of M , $Rm = M$ i.e. M is a cyclic module generated by m . Part (a) implies that there is some left ideal \mathfrak{m} of R such that $M \cong R/\mathfrak{m}$. Suppose that I is a proper left ideal of R such that $\mathfrak{m} \subseteq I \subsetneq R$. Applying the natural projection, we see that $0 \subseteq I/\mathfrak{m} \subsetneq R/\mathfrak{m}$, whence simplicity of R/\mathfrak{m} implies that I/\mathfrak{m} is trivial i.e. $I = \mathfrak{m}$. Thus by definition \mathfrak{m} is a maximal left ideal. \square

(c) Show that any non-zero homomorphism of simple R -modules is an isomorphism. Deduce that if M is simple, its endomorphism ring $\text{End}_R(M) := \text{Hom}_R(M, M)$ is a division ring. This result is known as *Schur's Lemma*.

Proof. Suppose that M, N are simple R -modules and let $f : M \rightarrow N$ be a nonzero R -module homomorphism. As $f(M) \neq 0$ is a submodule of N , by simplicity $f(M) = N$ i.e. f is surjective. As f is nonzero, $\ker(f)$ is a nonzero submodule of M whence $\ker(f) = 0$ i.e. f is injective. Hence f is an isomorphism.

Suppose M is simple and $f \in \text{End}_R(M) \setminus \{0\}$. Then f is an isomorphism, so the set-theoretic inverse f^{-1} is in fact an R -module isomorphism and $f^{-1} \in \text{End}_R(M)$. Hence $\text{End}_R(M)$ is a division ring. \square

Problem 7

Show that \mathbb{Q} is not a free \mathbb{Z} -module, that is \mathbb{Q} is not isomorphic to a direct sum of the form $\bigoplus_I \mathbb{Z}$, for any index set I . More generally, let R be a PID which is not a field and $K = \text{frac}(R)$ be its fraction field. Show that K is not a free R -module.

Proof. \square

Problem 8

Let R be a commutative ring. Recall that an ideal I of R is called *nilpotent* if there exists some $n \in \mathbb{N}$ such that $I^n = 0$.

(a) Let $i \in I$. Show that the element $r = 1 - i$ is invertible in R .

Proof. As I is a nilpotent ideal, there exists an $n \in \mathbb{N}$ such that $I^n = 0$. Then $i^n = 0$, so

$$1 = 1 - i^n = (1 - i)(1 + i + \cdots + i^{n-1}),$$

whence $1 - i \in R^\times$. \square

(b) Let M, N be R -modules and let $\varphi : M \rightarrow N$ be an R -module homomorphism. Show that φ induces an R -module homomorphism, $\bar{\varphi} : M/IM \rightarrow N/IN$.

Proof. Let $\pi_M : M \rightarrow M/IM$ and $\pi_N : N \rightarrow N/IN$ be the natural projections. Define a map $\bar{\varphi} : M/IM \rightarrow N/IN$ by $\bar{\varphi}(m + IM) := \varphi(m) + IN = (\pi_N \circ \varphi)(m)$. To see that this map is well defined, suppose that $m + IM = m' + IM$. Then there exist $i_1, \dots, i_s \in I$ and $m_1, \dots, m_s \in M$ such that $m - m' = i_1 m_1 + \dots + i_s m_s$. So

$$\varphi(m - m') = \varphi(i_1 m_1 + \dots + i_s m_s) = i_1 \varphi(m_1) + \dots + i_s \varphi(m_s) \in IN,$$

whence $\pi_N(\varphi(m)) - \pi_N(\varphi(m')) = \pi_N(\varphi(m - m')) = 0$, so $\pi_N(\varphi(m)) = \pi_N(\varphi(m'))$. □

(c) Prove that if $\bar{\varphi}$ is surjective, then φ is itself surjective.

Proof. Suppose that $\bar{\varphi}$ is surjective. Then $N/IN = \bar{\varphi}(M/IM) = (\bar{\varphi} \circ \pi_M)(M) = (\pi_N \circ \varphi)(M) = \varphi(M)/IN$. Take $n \in N$. Then there exists an $m \in M$ such that $n + IN = \bar{\varphi}(m + IM) = \varphi(m) + IN$, whence $n - \varphi(m) \in IN$. It follows that $n = \varphi(m) + (n - \varphi(m)) \in \varphi(M) + IN$, whence $N = \varphi(M) + IN$. As I is a nilpotent ideal, there exists a $k \in \mathbb{N}$ such that $I^k = 0$. Observe that

$$N = \varphi(M) + IN = \varphi(M) + I(\varphi(M) + IN) = \varphi(M) + I^2 N = \dots = \varphi(M) + I^k N = \varphi(M),$$

so φ is surjective. (Note: I couldn't see a way to use part (a) for this proof). □

Problem 9

Let G be a finite group and k a field. Consider the group ring $k[G]$.

(a) Let M be a k -vector space with a G -action. Show that M becomes a $k[G]$ -module. Conversely, if M is a $k[G]$ -module, show that M is a G -set.

(b) Let M, N be two $k[G]$ -modules. Show that $\text{Hom}_k(M, N)$ becomes a $k[G]$ -module with the following G -action: For $g \in G$ and $\varphi : M \rightarrow N$ a $k[G]$ -homomorphism define

$$(g \cdot \varphi)(m) := g\varphi(g^{-1}m), \text{ for } m \in M.$$