

Problem 1

Let $X = Y$ be an uncountable linearly ordered set such that for each $x \in X$, $\{y \in X : y < x\}$ is countable. Let $\mathcal{M} = \mathcal{N}$ be the σ -algebra of countable or co-countable sets, and let $\mu = \nu$ be defined on \mathcal{M} by $\mu(A) = 0$ if A is countable and $\mu(A) = 1$ if A is co-countable. Let $E = \{(x, y) \in X \times X : y < x\}$. Prove that E_x and E^y are measurable for all x, y , and that $\int \int \mathbb{1}_E d\mu d\nu$ and $\int \int \mathbb{1}_E d\nu d\mu$ exist but are not equal.

Problem 3

(a): Suppose (X, Σ, μ) is a σ -finite measure space and $f \in L^+(X)$. Let

$$G_f = \{(x, y) \in X \times [0, +\infty] : y \leq f(x)\}.$$

Show that G_f is $\Sigma \times \mathcal{B}_{\mathbb{R}}$ -measurable and $\mu \times m(G_f) = \int f d\mu$. Show also that the same is true if the inequality in the definition of G_f is made strict.

(b): Let (X, μ) be a σ -finite measure space. Fix $p \in [1, +\infty)$. Show that if $f \in L^p(X, \mu)$, then

$$\|f\|_p^p = p \int_0^\infty t^{p-1} \mu(\{x : f(x) > t\}) dt.$$

(c): Let (X, μ) be a σ -finite measure space. Show that if $f, g \in L^1(X, \mu)$ with $0 \leq f, g$ a.e., then

$$\|f - g\|_1 = \int_0^\infty \mu(\{x : f(x) > t\}) \Delta \mu(\{x : g(x) > t\}) dt.$$

Suggestion: it might be helpful to first show that for $a, b \in [0, +\infty)$ we have

$$|a - b| = \int_0^\infty |\mathbb{1}_{(t, \infty)}(a) - \mathbb{1}_{(t, \infty)}(b)| dt$$

Problem 4

If f is Lebesgue integrable on $(0, a)$ and $g(x) = \int_x^a t^{-1} f(t) dt$, then g is integrable on $(0, a)$ and $\int_0^a g(x) dx = \int_0^a f(x) dx$.

Problem 5

Let \mathcal{E}_q be the set of products of the form $\prod_{j=1}^d I_j$ where each I_j is an h -interval with the property that all of its finite endpoints are rational.

(a): Show that \mathcal{E}_q is an elementary family which generates the Borel sets.

(b): Suppose that μ is a Borel measure on \mathbb{R}^d with $0 < \mu((0, 1]^d) < +\infty$. If $\mu(E + x) = \mu(E)$ for every $x \in \mathbb{R}^d$, show that $\mu(E) = \mu((0, 1]^d) m(E)$ for every Borel $E \subseteq \mathbb{R}^d$.

Problem 6

Fix $d \in \mathbb{N}$.

(a): Let $s : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ be the map $s(x, y) = x + y$. Let μ, ν be finite, Borel measures on \mathbb{R}^d . Define $\mu * \nu = s_*(\mu \otimes \nu)$. Show that for every Borel $E \subseteq \mathbb{R}^d$ we have

$$\mu * \nu(E) = \int \int \mathbb{1}_E(x + y) d\mu(x) d\nu(y)$$

and

$$\int \mu(E - y) d\nu(y) = \mu * \nu(E) = \int \nu(E - x) d\mu(x).$$

Show as a consequence that

$$\mu * \nu(X) = \mu(X)\nu(X).$$

(b): Show that for finite, Borel measures μ, ν, η on \mathbb{R}^d we have

$$(\mu * \nu) * \eta = \mu * (\nu * \eta).$$

(c): For $f, g \in L^1(\mathbb{R}^d)$ show that

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |f(y)g(x - y)| dx dy = \|f\|_1 \|g\|_1.$$

Explain why this implies that $y \mapsto f(y)g(x - y)$ is in $L^1(\mathbb{R}^d)$ for almost every $x \in \mathbb{R}^d$ and why if we set $f * g(x) = \int_{\mathbb{R}^d} f(y)g(x - y) dy$ then we have that $f * g \in L^1(\mathbb{R}^d)$ and

$$\|f * g\|_1 \leq \|f\|_1 \|g\|_1.$$

(d): Adopt notation as in Problem 1 of HW5. Show that if $f, g \in L^1(\mathbb{R}^d)$ are nonnegative then $(f dm) * (g dm) = f * g dm$ with m being the Lebesgue measure.

(e): Show that for $f, g, k \in L^1(\mathbb{R}^d)$ we have that

$$(f * g) * k = f * (g * k) \text{ almost everywhere.}$$