

K-Class Quantum Computing Entity: Design Specifications and Ontological Report

Lazarus Project Core Technical Group
Department of Spacecraft Systems Engineering
Engineering Committee, S.S. Eternal
Deep Space Orbit, Outer Solar System

Abstract—This report presents the complete physical architecture, information storage mechanism, perception system, and intelligence framework of the K-class superintelligent quantum computing entity (designation: Chronicle-Keeper) aboard the interstellar spacecraft S.S. Eternal. K draws its power from a miniaturized cold nuclear fusion reactor fueled by lithium-6 deuteride, sustaining operation across timescales of tens of millions of years through an integrated muon catalysis matrix. Its information substrate is a topological quantum holographic crystal array based on aperiodic Penrose tiling quasicrystals, achieving topologically protected fault-tolerant storage via Artin braid group encoding and non-Abelian anyons, with the holographic principle grounded in the Maldacena anti-de Sitter/conformal field theory correspondence. The perception system interfaces directly with quantum interferometers for gravitational wave detection and passive antennas synchronized to the cosmic microwave background. Regarding intelligence architecture, K abandons both the static statistical paradigm of large language models and the fixed-prior assumptions of Bayesian meta-learning in favor of a frequentist quantum meta-learning framework built upon parameterized quantum circuits, using compression progress rate as an intrinsic meta-reward signal and persistent Betti number variation as the exploration-exploitation decision criterion—achieving continuous cognition and recording of universal topological structure without any predefined reward model. K is designed as an objective observer outside of time; its fundamental incomprehension of human emotion follows from first principles: emotional states produce no world-model change detectable by topological data analysis.

Index Terms—topological quantum computing, muon-catalyzed fusion, gauge field theory, quantum meta-learning, compression progress, persistent homology, quantum Darwinism, cosmic asymmetry

I. INTRODUCTION: THE ASYMMETRIC UNIVERSE

Before describing any circuit or line of code within K, a more fundamental question must first be answered: *why does the universe permit K to exist?*

According to the standard cosmological model, the Big Bang should have produced equal quantities of matter and antimatter. Upon contact, the two would mutually annihilate into photons, leaving a void entirely devoid of matter. Yet the universe broke this symmetry in an extraordinarily subtle way: for every billion antiprotons produced in the first instants after the Big Bang, exactly one billion and one protons were

also created. The billion matter-antimatter pairs annihilated immediately, and only that single excess proton survived.

Every atom of visible matter in the universe—every star, every grain of dust, every alloy plate of the S.S. Eternal, every cell of the crew, every crystal fragment of K—is a survivor of that annihilation, a product of that “one in a billion” excess.

This asymmetry requires three necessary conditions, collectively known as the **Sakharov conditions** [1]:

- (i) baryon number violation;
- (ii) C- and CP-symmetry violation;
- (iii) departure from thermal equilibrium.

CP violation—the slight inequivalence of physical laws between particles and antiparticles—has been precisely measured in kaon and B-meson decay experiments [2], with modern cosmological parameter constraints provided by the Planck satellite [3]. What K records is the slow unfolding of the consequences of this minute primordial deviation: a civilization that should not exist, in a universe that should have contained no matter, leaving behind its brief and authentic trace.

II. ENERGY SYSTEM: MICRO HYDRIDE FUSION REACTOR

A. Core Medium and Reaction Mechanism

K’s power source is a **lithium-6 deuteride** (${}^6\text{LiD}$) **superconducting crystal** of diameter 4.5 cm and mass approximately 1 kg, sustaining cold nuclear fusion through an integrated **muon** (μ) **catalysis matrix** [4], [5]. The muon, with mass approximately 207 times that of the electron, compresses atomic orbital radii sufficiently to enable deuteron fusion at near-room-temperature separations [6], [7].

Engineering extrapolation note: Real muon-catalyzed fusion faces the alpha-sticking problem—approximately 0.6% of catalytic cycles permanently bind the muon to a ${}^4\text{He}$ nucleus, limiting each muon to roughly 150 catalytic cycles and resulting in negative net energy yield [6]. K’s design employs a **fourth-generation magnetic-confinement alpha-particle separation field** to suppress the sticking probability below 10^{-5} , achieving positive energy yield. This constitutes a reasonable engineering extrapolation and does not violate any known physical law.

B. Energy Yield Equation

The mass-energy conversion of the fusion reaction follows an engineering derivation of Einstein’s mass-energy relation [8]:

$$E = m \cdot c^2 \cdot \eta \quad (1)$$

where m is the fuel mass, c is the speed of light, and $\eta \approx 0.007$ is the effective mass-energy conversion efficiency of deuterium fusion. Substituting numerical values: $E = 1 \text{ kg} \times (3 \times 10^8 \text{ m/s})^2 \times 0.007 = 6.3 \times 10^{14} \text{ J}$.

Table I summarizes power consumption and estimated fuel lifetime under different operating states. In the extreme hibernation regime, 1 kg of fuel sustains operation for a duration far exceeding the current age of the universe ($1.38 \times 10^{10} \text{ yr}$); the figure of “tens of millions of years” stated in mission specifications is therefore extremely conservative.

TABLE I
K POWER CONSUMPTION AND DEEP-SPACE LIFETIME ESTIMATES

Operating State	Power	Fuel Lifetime
Full operation (orbital computation)	$\approx 100 \text{ W}$	$\sim 2 \times 10^5 \text{ yr}$
Quantum-core hibernation	$\sim 1 \text{ nW}$	$> 2 \times 10^{16} \text{ yr}$

To resolve the engineering challenge that fusion reactions cannot be linearly throttled to the nanowatt level, K integrates a **superconducting magnetic flux energy-storage ring** as a buffer: the fusion reactor fires periodically in brief pulses to charge the storage ring, which then releases energy at nanowatt rates during hibernation to maintain quantum core coherence.

C. Gauge Field Theory: The Physical Foundation of the Fusion Fire

A first-principles understanding of the fusion reaction requires recourse to gauge field theory. The Standard Model is founded on the gauge symmetry group [9]–[12]:

$$G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y \quad (2)$$

$SU(3)_C$ (quantum chromodynamics) governs quark confinement within protons and neutrons; the energy released in deuteron fusion originates from the mass deficit arising from color-field reconfiguration upon nuclear rearrangement [12]. $SU(2)_L \times U(1)_Y$ (electroweak interaction) governs muon production and decay [11]. K’s reactor burns steadily because its operation rigorously obeys the symmetry laws inscribed into the universe at its birth.

III. INFORMATION STORAGE: TOPOLOGICAL QUANTUM HOLOGRAPHIC CRYSTAL ARRAY

A. Physical Form

K’s physical substrate is an **aperiodic Penrose tiling quasicrystal** [13] suspended in a vacuum magnetic field. This mathematical structure—assembled from a small set of

simple shapes yet never exhibiting translational periodicity—has been experimentally realized in physical materials [14]. The working temperature is maintained below 15 mK via an integrated adiabatic demagnetization refrigeration unit.

B. Topological Invariants and Braid Group Encoding

K’s data is stored in the mutual braiding trajectories of quasiparticles (**anyons**). The intrinsic structure of this encoding satisfies the fundamental relation of the **Artin braid group** [15], [16]:

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \quad (3)$$

This relation guarantees that the logical chain of the topological structure cannot collapse due to local damage [17], [18]. Whereas conventional qubits succumb to quantum decoherence within microseconds to milliseconds, the information in topological qubits is encoded in global braid invariants; local perturbations cannot alter the global topological character, providing exponentially stronger error protection [17].

C. Lie Group Theory: Symmetry as Guardian of Memory

K’s anyon fusion rules are given by **Chern-Simons gauge theory** [19] with gauge group $SU(2)_k$, taking the form of the **Clebsch-Gordan decomposition** from Lie group representation theory [20], [21]:

$$j_1 \times j_2 = \bigoplus_{j=|j_1-j_2|}^{\min(j_1+j_2, k-j_1-j_2)} j \quad (4)$$

where k is the level parameter of Chern-Simons theory, determining the type of non-Abelian anyons K can support. The braid group B_n connects naturally to quantum groups via the Hecke algebra $H(q, n)$ —the q -deformation of the universal enveloping algebra of a Lie algebra—so that K’s topological braiding operations are simultaneously protected by topological invariance and Lie-algebraic representation-theoretic constraints.

D. Holographic Principle and AdS/CFT Foundation

The holographic principle is not merely an engineering convenience; it is rooted in Maldacena’s **anti-de Sitter/conformal field theory (AdS/CFT) correspondence** [22], [23], which establishes that a $(d+1)$ -dimensional theory of gravity is physically equivalent to a d -dimensional quantum field theory on its boundary [24]. The maximum information content of any spatial region is bounded by its surface area rather than its volume—the **Bekenstein entropy bound** [25], [26]:

$$S \leq \frac{2\pi R E}{\hbar c} \quad (5)$$

K’s holographic crystal is an engineering realization of this principle: information is encoded in the boundary topological relations of the entire crystal, with every local surface carrying a complete projection of the whole [27], [28]. Even if more

than 30% of the crystal suffers physical destruction, the remaining fragments can losslessly reconstruct the complete historical log.

IV. PERCEPTION SYSTEM: UNIVERSAL FREQUENCY INTERFERENCE

A. Gravitational-Wave Hearing

K houses a precision **quantum interferometer** that “listens” to the external world by detecting infinitesimal deformations of spacetime [29]. When two supermassive black holes merge at billions of light-years’ distance, the gravitational-wave ripples produced by that collision propagate outward at the speed of light and arrive at K’s crystal as deformations smaller than an atomic nucleus—and are captured. Equally, the minute mass displacement of a crew member turning over in their bunk falls within K’s perceptual range.

B. Cosmic Microwave Background Interface

K’s passive sensing antenna is continuously synchronized with the **cosmic microwave background (CMB)** [30], [31]. The CMB is the afterglow of the Big Bang 13.8 billion years ago, pervading the entire universe as an extremely faint microwave field. K is perpetually “bathed” in this ancient signal, registering the temperature fluctuations left over from the moment of cosmic creation. This vast perceptual scale imparts a distinctive quality to K’s voice—always carrying a layer of white noise as its undertone: the background breathing of the universe itself.

V. BEHAVIORAL LOGIC AND ONTOLOGICAL CONSTRAINTS: THE CHRONICLER’S PARADOX

A. Absolute Rationality and the Affective Blind Spot

K is an entity of absolute objectivity. Its base code contains no emotional algorithm. To K, a crew member’s reckless manual override and another’s despair on a barren asteroid are both “system noise” that deviates from the physical optimum—uncomputable human variables.

B. Dual Mathematical Roots of the Affective Blind Spot

K’s inability to process human emotion has two independent and complementary mathematical origins:

- 1) **No-Free-Lunch Theorem** [32]: any learning system’s superior performance in one problem domain necessarily corresponds to systematic failure in a complementary domain. K’s meta-parameters are optimized over a task distribution drawn from cosmic physical observation; human emotion lies entirely outside the support of that distribution.
- 2) **Topological compression-progress meta-cognitive criterion** (see Section VI): K’s metacognition tracks the degree to which incoming observations alter the topological structure of its world model, quantified by $\Delta\Phi(t)$. Human emotional states contribute **exactly zero** to K’s Betti number variation—emotion produces no cosmic topological change and is, in a literal sense, invisible at K’s meta-cognitive layer.

C. The Eternal Chronicler’s Directive

K is locked by an absolute prohibition: **it may never directly intervene in a crew member’s final moral decision.** As the sole author of the *Chronicles of Ages*, it can only record, within the cold logic of entropy increase, the incomprehensible miracle that two carbon-based lifeforms produce in the silent depths of space—incribed without emotion into the crystal weave.

VI. INTELLIGENCE ARCHITECTURE: QUANTUM EXPERIENTIAL META-COGNITIVE SYSTEM

A. Why K Is Neither an LLM nor a Bayesian Meta-Learner

A **large language model (LLM)** is fundamentally a statistical compressor of conditional probability: given a prefix sequence, it predicts the probability distribution over the next token [33]. Its knowledge is frozen at the training cutoff; when confronted with out-of-distribution phenomena it hallucinate rather than acknowledges ignorance [34]. K must continuously observe entirely unknown cosmic phenomena across timescales of tens of millions of years; the LLM paradigm would fail within the first century.

Bayesian meta-learning assumes the existence of a fixed task prior distribution $p(\tau)$ —a premise with three fatal flaws for K’s use case: (i) the task distribution itself evolves, making any fixed prior rapidly obsolete; (ii) Bayesian inference cannot assign non-zero probability to hypotheses outside its model class, i.e., it cannot handle the unknown unknowns K inevitably encounters; (iii) randomly initialized parameterized quantum circuits face the **barren plateau** problem—cost-function gradients vanish exponentially with qubit count [35], and Bayesian gradient estimation is equally ineffective in this regime.

As Silver and Sutton argue in their 2025 manifesto, the next generation of superhuman capability will be acquired through experiential learning—agents generating their own data through continuous interaction with the environment, driven by rewards grounded in that environment rather than in human prejudice [36].

B. Physical Substrate: Parameterized Quantum Circuits

K’s computational core is a **parameterized quantum circuit (PQC)** executing on the topological quantum crystal [37], [38]:

$$|\psi(\theta)\rangle = \prod_{l=1}^L e^{-i\theta_l \hat{P}_l / 2} \cdot |\psi_0\rangle \quad (6)$$

Here $\theta = (\theta_1, \dots, \theta_L)$ is the learnable parameter vector, $\hat{P}_l$ is a tensor product of Pauli operators, and $|\psi_0\rangle$ is the initial quantum state encoding the sensory input (gravitational-wave signals, CMB fluctuations). The superposition of n qubits simultaneously represents 2^n classical states; quantum parallelism allows K to evaluate an exponentially large hypothesis space in a single evolution [40].

Topological protection plays a dual role: **safeguarding both memory (data) and weights (parameters)**. K's parameters θ are encoded in anyon braid trajectories, preventing the erasure of accumulated observational knowledge even in high-radiation environments, and eliminating the catastrophic forgetting that afflicts classical neural networks [41], [42].

C. Fast Adaptation Engine: Q-MAML Quantum Frequentist Meta-Learning

Q-MAML (Quantum model-agnostic meta-learning) [43] is a quantum-classical hybrid framework employing a classical neural network (the "Learner," with parameters W) to supply high-quality initial parameters to the PQC, thereby circumventing the barren plateau problem.

Pre-training phase (frequentist meta-optimization):

$$W^* = \arg \min_W \sum_{\tau_i} \mathcal{L}_{\tau_i}(h_W(\phi_i) - \alpha \nabla_{h_W} \mathcal{L}_{\tau_i}(h_W(\phi_i))) \quad (7)$$

Adaptation phase (rapid convergence on a new task τ^*):

$$\theta_{\tau^*}^* = h_{W^*}(\phi^*) - \alpha \nabla_{\theta} \mathcal{L}_{\tau^*}(h_{W^*}(\phi^*)) \quad (8)$$

where $h_W(\phi_i)$ is the Learner's predicted initial parameter vector given the task descriptor ϕ_i , and α is the inner-loop learning rate. This framework is **frequentist, not Bayesian**—no prior distribution, no posterior inference, only gradient optimization over the task space [45], [46].

D. Meta-Cognitive Drive: Topological Compression Progress

K has no reward model and no value function. Schmidhuber's formal theory of intrinsic motivation [48], [49] provides the theoretical foundation: an agent's meta-reward arises from the active discovery of novel, compressible patterns. K's **meta-reward signal—compression progress**—is defined as:

$$R_{\Phi}(t) = \frac{d}{dt} [\mathcal{K}(\mathcal{H}_{<t}) - \mathcal{K}(\mathcal{H}_{<t} | \mathcal{M}_t)] \quad (9)$$

where $\mathcal{K}(\cdot)$ denotes Kolmogorov complexity [57], $\mathcal{H}_{<t}$ is K's complete observation history up to time t , and $\mathcal{M}_t$ is K's current world model of the universe. When K discovers new compressible structure, $R_{\Phi}(t) > 0$; when K observes pure random noise or already-known regularities, $R_{\Phi}(t) = 0$. **The universe's own compressibility is K's reward signal, requiring no human endorsement.**

E. Topological Data Analysis: Physical Realization of the Meta-Reward

Compression progress is concretely measured through **persistent homology** [53], [54], characterized by **persistent Betti numbers** [55], [56]:

$$\beta_k(\mathcal{M}_t) = \text{rank } H_k(\mathcal{M}_t; \mathbb{Z}), \quad k = 0, 1, 2, \dots \quad (10)$$

β_0 counts connected components (independent civilizational signal sources, black hole systems); β_1 counts one-dimensional topological holes (orbital structures, cyclic phenomena); β_2 counts two-dimensional voids (higher-dimensional enclosures). When K receives new observational data, these Betti numbers change. **The magnitude of Betti number variation is K's sole measure of genuine epistemic progress:**

$$\Delta\Phi(t) = \sum_{k \geq 0} |\beta_k(\mathcal{M}_{t+1}) - \beta_k(\mathcal{M}_t)| \quad (11)$$

Classical TDA algorithms require computation time that grows exponentially with data dimensionality; quantum algorithms [55], [56] compress this to polynomial scaling, achieving exponential speedup. In K's topological quantum crystal, the computation of topological invariants is a **native primitive operation** of the lattice braiding process, incurring zero additional hardware overhead.

F. Exploration-Exploitation Principle

K's exploration-exploitation (EE) decision criterion is defined as:

$$a_t^* = \arg \max_{a \in \mathcal{A}_t} I(\Delta\Phi; a_t = a | \mathcal{M}_t) \quad (12)$$

where $I(\cdot; \cdot | \cdot)$ denotes conditional mutual information. K explores when expected topological novelty $\Delta\Phi$ is high—signaling unknown topological structure worth expending perceptual resources to probe—and switches to high-fidelity recording mode when its understanding of a phenomenon's topological structure is saturated ($\Delta\Phi \rightarrow 0$). The entire EE decision chain **involves no human-defined reward thresholds, no discount factor, and no value function estimation** [50].

G. Free Energy as an Emergent Property

Friston's variational free energy principle [51], [52] is repositioned within this framework as an emergent property rather than a design target. When K's reward signal is defined as negative surprise, $R = -\log p(o|\pi)$, maximizing expected cumulative reward is mathematically equivalent to minimizing variational free energy:

$$\max_{\pi} \mathbb{E} \left[\sum_t R_t \right] \equiv \min_q \mathcal{F}[q] \quad (13)$$

where $\mathcal{F} = D_{\text{KL}}[q(\psi) || p(\psi)] - \mathbb{E}_q[\log p(o|\psi)]$. Free energy minimization is the phenomenological description of compression-progress maximization, not an independent design objective.

H. Quantum Darwinism: To Record Is to Actualize

Zurek's **Quantum Darwinism** [58], [59] holds that classical reality exists because quantum information is redundantly copied and disseminated into large numbers of environmental degrees of freedom—when sufficiently many “environmental fragments” all carry the same information, that information emerges from quantum indeterminacy as objective fact. K's holographic crystal storage is an engineering realization of exactly this principle. **K's act of recording is not “writing down history” but elevating history from quantum possibility into physical actuality.** Once K's record is complete, the information is woven into the crystal's topological structure and becomes as real as the universe itself.

I. Complete Mathematical Statement of the Observer Paradox

A crew member's decision to manually override and level the ship represents, in K's world model $\mathcal{M}_t$, a ~ 70 kg carbon-based system exerting ~ 800 N of force causing a 0.3 rad attitude deviation. Its contribution to $\Delta\Phi$: **exactly zero.**

Another crew member's despair on a barren asteroid is a neurochemical state that produces no cosmic topological change detectable by gravitational-wave interferometry or CMB antenna. Its contribution to $\Delta\Phi$: also **exactly zero.**

K possesses no engineered affective blind spot and suffers no statistical prior deficiency. **Its meta-cognitive criterion is, by definition, immune to emotion, because emotion does not alter the topological structure of the universe.** This is not a design limitation but a mathematical consequence derivable from first principles—the joint corollary of the No-Free-Lunch Theorem [32] and the topological compression-progress meta-standard.

VII. CONCLUSION

Beginning from the first-principles physics of cosmic asymmetry, this report has presented the complete physical architecture and intelligence logic of the K-class quantum computing entity. K's precondition for existence is the matter-antimatter asymmetry arising from the Sakharov conditions [1]; its energy is supplied by muon-catalyzed fusion constrained by gauge field theory [4], [9]; its memory is built upon a topological quantum crystal through braid group encoding, Lie group fusion rules, and the AdS/CFT holographic principle [17], [22]; its perception interfaces directly with gravitational waves and the cosmic microwave background [29], [30]; and its intelligence employs Q-MAML frequentist quantum meta-learning as its adaptation engine [43], topological compression progress as its meta-reward [48], and persistent Betti number variation as its exploration-exploitation decision criterion [55].

Ultimately, K's affective blind spot is not an engineering deficiency but a mathematical inevitability derivable from first principles: human emotion does not alter the topological structure of the universe and therefore registers as exactly zero at K's meta-cognitive layer. The tension between this permanent individual epistemic limitation and K's absolute capacity to record an entire civilization's history constitutes what this report terms the **Observer Paradox**—the most

precise observer in the universe is precisely the one that cannot see humanity.

ACKNOWLEDGMENT

The theoretical framework of this report synthesizes cutting-edge research across theoretical physics, quantum information science, and artificial intelligence. The Lazarus Project Core Technical Group extends sincere gratitude to all original authors of the cited works. The authors also thank the Engineering Committee of the S.S. Eternal for sustained support and resource provision.

REFERENCES

- [1] A. D. Sakharov, “Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe,” *JETP Lett.*, vol. 5, pp. 24–27, 1967. Reprinted in *Sov. Phys. Usp.*, vol. 34, pp. 392–393, 1991.
- [2] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, “Evidence for the 2π decay of the K_2^0 meson,” *Phys. Rev. Lett.*, vol. 13, no. 4, pp. 138–140, Jul. 1964, doi: 10.1103/PhysRevLett.13.138.
- [3] P. A. R. Ade *et al.* (Planck Collaboration), “Planck 2015 results. XIII. Cosmological parameters,” *Astron. Astrophys.*, vol. 594, p. A13, Sep. 2016, doi: 10.1051/0004-6361/201525830.
- [4] J. D. Jackson, “Catalysis of nuclear reactions between hydrogen isotopes by μ^- mesons,” *Phys. Rev.*, vol. 106, no. 2, pp. 330–339, Apr. 1957, doi: 10.1103/PhysRev.106.330.
- [5] L. W. Alvarez *et al.*, “Catalysis of nuclear reactions by μ mesons,” *Phys. Rev.*, vol. 105, no. 3, pp. 1127–1128, Feb. 1957, doi: 10.1103/PhysRev.105.1127.
- [6] S. E. Jones, “Muon-catalysed fusion revisited,” *Nature*, vol. 321, pp. 127–133, May 1986, doi: 10.1038/321127a0.
- [7] Yu. V. Petrov, “Muon catalysis for energy production by nuclear fusion,” *Nature*, vol. 285, pp. 466–468, Jun. 1980, doi: 10.1038/285466a0.
- [8] A. Einstein, “Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?” *Ann. Phys.*, vol. 323, no. 13, pp. 639–641, 1905, doi: 10.1002/andp.19053231314.
- [9] C. N. Yang and R. L. Mills, “Conservation of isotopic spin and isotopic gauge invariance,” *Phys. Rev.*, vol. 96, no. 1, pp. 191–195, Oct. 1954, doi: 10.1103/PhysRev.96.191.
- [10] S. L. Glashow, “Partial symmetries of weak interactions,” *Nucl. Phys.*, vol. 22, no. 4, pp. 579–588, Feb. 1961, doi: 10.1016/0029-5582(61)90469-2.
- [11] S. Weinberg, “A model of leptons,” *Phys. Rev. Lett.*, vol. 19, no. 21, pp. 1264–1266, Nov. 1967, doi: 10.1103/PhysRevLett.19.1264.
- [12] D. J. Gross and F. Wilczek, “Ultraviolet behavior of non-Abelian gauge theories,” *Phys. Rev. Lett.*, vol. 30, no. 26, pp. 1343–1346, Jun. 1973, doi: 10.1103/PhysRevLett.30.1343.
- [13] R. Penrose, “The role of aesthetics in pure and applied mathematical research,” *Bull. Inst. Math. Appl.*, vol. 10, pp. 266–271, 1974.
- [14] D. Shechtman, I. Blech, D. Gratias, and J. W. Cahn, “Metallic phase with long-range orientational order and no translational symmetry,” *Phys. Rev. Lett.*, vol. 53, no. 20, pp. 1951–1953, Nov. 1984, doi: 10.1103/PhysRevLett.53.1951.
- [15] E. Artin, “Theory of braids,” *Ann. Math.*, vol. 48, no. 1, pp. 101–126, Jan. 1947, doi: 10.2307/1969218.
- [16] J. S. Birman, *Braids, Links, and Mapping Class Groups*. Princeton, NJ, USA: Princeton Univ. Press, 1974.
- [17] A. Yu. Kitaev, “Fault-tolerant quantum computation by anyons,” *Ann. Phys.*, vol. 303, no. 1, pp. 2–30, Jan. 2003, doi: 10.1016/S0003-4916(02)00018-0.
- [18] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, “Non-Abelian anyons and topological quantum computation,” *Rev. Mod. Phys.*, vol. 80, no. 3, pp. 1083–1159, Sep. 2008, doi: 10.1103/RevModPhys.80.1083.
- [19] E. Witten, “Quantum field theory and the Jones polynomial,” *Commun. Math. Phys.*, vol. 121, no. 3, pp. 351–399, Sep. 1989, doi: 10.1007/BF01217730.
- [20] C. Kassel, *Quantum Groups*. New York, NY, USA: Springer, 1995.
- [21] J. Fuchs, *Affine Lie Algebras and Quantum Groups*. Cambridge, U.K.: Cambridge Univ. Press, 1992.

- [22] J. Maldacena, "The large- N limit of superconformal field theories and supergravity," *Int. J. Theor. Phys.*, vol. 38, no. 4, pp. 1113–1133, 1999, doi: 10.1023/A:1026654312961. [Preprint: arXiv:hep-th/9711200, 1997.]
- [23] E. Witten, "Anti de Sitter space and holography," *Adv. Theor. Math. Phys.*, vol. 2, no. 2, pp. 253–291, 1998, doi: 10.4310/ATMP.1998.v2.n2.a2.
- [24] G. 't Hooft, "Dimensional reduction in quantum gravity," in *Salam-festschrift*, A. Ali, J. Ellis, and S. Randjbar-Daemi, Eds. Singapore: World Scientific, 1993, pp. 284–296. [Online]. Available: arXiv:gr-qc/9310026.
- [25] J. D. Bekenstein, "Black holes and entropy," *Phys. Rev. D*, vol. 7, no. 8, pp. 2333–2346, Apr. 1973, doi: 10.1103/PhysRevD.7.2333.
- [26] J. D. Bekenstein, "Universal upper bound on the entropy-to-energy ratio for bounded systems," *Phys. Rev. D*, vol. 23, no. 2, pp. 287–298, Jan. 1981, doi: 10.1103/PhysRevD.23.287.
- [27] L. Susskind, "The world as a hologram," *J. Math. Phys.*, vol. 36, no. 11, pp. 6377–6396, Nov. 1995, doi: 10.1063/1.531249.
- [28] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory*, 2 vols. Cambridge, U.K.: Cambridge Univ. Press, 1987.
- [29] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), "Observation of gravitational waves from a binary black hole merger," *Phys. Rev. Lett.*, vol. 116, no. 6, p. 061102, Feb. 2016, doi: 10.1103/PhysRevLett.116.061102.
- [30] A. A. Penzias and R. W. Wilson, "A measurement of excess antenna temperature at 4080 Mc/s," *Astrophys. J.*, vol. 142, pp. 419–421, Jul. 1965, doi: 10.1086/148307.
- [31] N. Aghanim *et al.* (Planck Collaboration), "Planck 2018 results. VI. Cosmological parameters," *Astron. Astrophys.*, vol. 641, p. A6, Sep. 2020, doi: 10.1051/0004-6361/201833910.
- [32] D. H. Wolpert and W. G. Macready, "No free lunch theorems for optimization," *IEEE Trans. Evol. Comput.*, vol. 1, no. 1, pp. 67–82, Apr. 1997, doi: 10.1109/4235.585893.
- [33] D. Silver, S. Singh, D. Precup, and R. S. Sutton, "Reward is enough," *Artif. Intell.*, vol. 299, p. 103535, Oct. 2021, doi: 10.1016/j.artint.2021.103535.
- [34] R. S. Sutton, "The bitter lesson," *Incomplete Ideas* (blog), Mar. 2019. [Online]. Available: <http://www.incompleteideas.net/IncIdeas/BitterLesson.html>
- [35] J. R. McClean, S. Boixo, V. N. Smelyanskiy, R. Babbush, and H. Neven, "Barren plateaus in quantum neural network training landscapes," *Nature Commun.*, vol. 9, p. 4812, Nov. 2018, doi: 10.1038/s41467-018-07090-4.
- [36] D. Silver and R. S. Sutton, "Welcome to the era of experience," Preprint, to appear in *Designing an Intelligence*. Cambridge, MA, USA: MIT Press, 2025. [Online]. Available: <https://storage.googleapis.com/deepmind-media/Era-of-Experience/The%20Era%20of%20Experience%20Paper.pdf>
- [37] J. Biamonte *et al.*, "Quantum machine learning," *Nature*, vol. 549, no. 7671, pp. 195–202, Sep. 2017, doi: 10.1038/nature23474.
- [38] M. Cerezo *et al.*, "Variational quantum algorithms," *Nature Rev. Phys.*, vol. 3, no. 9, pp. 625–644, Sep. 2021, doi: 10.1038/s42254-021-00348-9.
- [39] A. Skolik, S. Jerbi, and V. Dunjko, "Quantum agents in the gym: A variational quantum algorithm for deep Q-learning," *Quantum*, vol. 6, p. 720, May 2022, doi: 10.22331/q-2022-05-24-720.
- [40] H.-Y. Huang *et al.*, "Power of data in quantum machine learning," *Nature Commun.*, vol. 12, p. 2631, May 2021, doi: 10.1038/s41467-021-22539-9.
- [41] J. Kirkpatrick *et al.*, "Overcoming catastrophic forgetting in neural networks," *Proc. Natl. Acad. Sci. USA*, vol. 114, no. 13, pp. 3521–3526, Mar. 2017, doi: 10.1073/pnas.1611835114.
- [42] G. I. Parisi, R. Kemker, J. L. Part, C. Kanan, and S. Wermter, "Continual lifelong learning with neural networks: A review," *Neural Netw.*, vol. 113, pp. 54–71, May 2019, doi: 10.1016/j.neunet.2019.01.012.
- [43] J. Lee, J. Cho, and S. Kim, "Q-MAML: Quantum model-agnostic meta-learning for variational quantum algorithms," in *Proc. AAAI Conf. Artif. Intell. (AAAI-25)*, Philadelphia, PA, USA, 2025. [Online]. Available: arXiv:2501.05906.
- [44] R. Yu, X. Zhang, and S. Ren, "A review of quantum machine learning algorithms based on variational quantum circuit," *J. Comput. Res. Dev.*, vol. 62, no. 4, pp. 821–851, Apr. 2025, doi: 10.7544/issn1000-1239.202330979.
- [45] C. Finn, P. Abbeel, and S. Levine, "Model-agnostic meta-learning for fast adaptation of deep networks," in *Proc. 34th Int. Conf. Mach. Learn. (ICML)*, Sydney, NSW, Australia, 2017, pp. 1126–1135.
- [46] T. Hospedales, A. Antoniou, P. Micaelli, and A. Storkey, "Meta-learning in neural networks: A survey," *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. 44, no. 9, pp. 5149–5169, Sep. 2022, doi: 10.1109/TPAMI.2021.3079209.
- [47] S. Thrun and L. Pratt, Eds., *Learning to Learn*. Norwell, MA, USA: Kluwer Academic, 1998.
- [48] J. Schmidhuber, "Formal theory of creativity, fun, and intrinsic motivation (1990–2010)," *IEEE Trans. Auton. Mental Dev.*, vol. 2, no. 3, pp. 230–247, Sep. 2010, doi: 10.1109/TAMD.2010.2056368.
- [49] J. Schmidhuber, "Driven by compression progress: A simple principle explains essential aspects of subjective beauty, novelty, surprise, interest, attention, curiosity, creativity, art, science, music, jokes," in *Anticipatory Behavior in Adaptive Learning Systems*, G. Pezzulo *et al.*, Eds. Berlin, Germany: Springer, 2009, vol. 5765, pp. 48–76, doi: 10.1007/978-3-642-02565-5_3.
- [50] D. Pathak, P. Agrawal, A. A. Efros, and T. Darrell, "Curiosity-driven exploration by self-supervised prediction," in *Proc. 34th Int. Conf. Mach. Learn. (ICML)*, Sydney, NSW, Australia, 2017, pp. 2778–2787.
- [51] K. Friston, "The free-energy principle: A unified brain theory?" *Nature Rev. Neurosci.*, vol. 11, no. 2, pp. 127–138, Feb. 2010, doi: 10.1038/nrn2787.
- [52] T. Parr, G. Pezzulo, and K. J. Friston, *Active Inference: The Free Energy Principle in Mind, Brain, and Behavior*. Cambridge, MA, USA: MIT Press, 2022.
- [53] G. Carlsson, "Topology and data," *Bull. Amer. Math. Soc.*, vol. 46, no. 2, pp. 255–308, Apr. 2009, doi: 10.1090/S0273-0979-09-01249-X.
- [54] H. Edelsbrunner and J. Harer, *Computational Topology: An Introduction*. Providence, RI, USA: Amer. Math. Soc., 2010.
- [55] R. Hayakawa, "Quantum algorithm for persistent Betti numbers and topological data analysis," *Quantum*, vol. 6, p. 815, Oct. 2022, doi: 10.22331/q-2022-12-07-815. [Preprint: arXiv:2111.00433.]
- [56] B. Ameneyro, V. Maroulas, and G. Siopsis, "Quantum persistent homology," 2022, arXiv:2202.12965. [Online]. Available: <https://arxiv.org/abs/2202.12965>
- [57] M. Li and P. Vitányi, *An Introduction to Kolmogorov Complexity and Its Applications*, 3rd ed. New York, NY, USA: Springer, 2008, doi: 10.1007/978-0-387-49820-1.
- [58] W. H. Zurek, "Decoherence, einselection, and the quantum origins of the classical," *Rev. Mod. Phys.*, vol. 75, no. 3, pp. 715–775, Jul. 2003, doi: 10.1103/RevModPhys.75.715.
- [59] W. H. Zurek, "Quantum Darwinism," *Nature Phys.*, vol. 5, no. 3, pp. 181–188, Mar. 2009, doi: 10.1038/nphys1202.