



WCPFC Scientific Committee
Stock Assessment

South Pacific albacore 2024: KIMSA assessment

2024

Kyuhan Kim

DEMONSTRATION

Table of contents

1	Introduction	3
2	Background	3
2.1	Fisheries	3
2.2	Distribution and movement	4
2.3	Depth use and availability to fishing	4
2.4	Growth	4
2.5	Reproduction	4
2.6	Natural mortality	5
3	Data compilation	5
3.1	Assessment area and data period	5
3.2	Fisheries and catch	5
3.3	Abundance indices	5
3.4	Length composition	5
3.5	Age observations	6
4	Model description	6
4.1	Model framework	6
4.2	Population processes	6
4.2.1	Recruitment and initial population	6
4.2.2	Growth and natural mortality	6
4.2.3	Movement	7
4.2.4	Reproductive potential	7
4.3	Fishing processes	7
4.4	Observation models	7
4.5	Parameter estimation	7
4.6	Stepwise development	7
4.7	Model ensemble	8
5	Model diagnostics	8
5.1	Starting-value trials (jitter)	8
5.2	Simulation-estimation tests (self-test)	8
5.3	Retrospective analysis	8
5.4	Hindcast evaluation	8
5.5	Likelihood profiles and identifiability	9
5.6	Data fit	9
5.7	Population estimates	9
5.8	Stock status	9
6	Discussion	10
6.1	Interpretation	10
6.2	Limitations	10
6.3	Conclusions and research priorities	10
7	Acknowledgements	10
8	References	10

9 Tables	12
10 Figures	15
11 Supplementary outputs	49

This demonstration rebuilds the South Pacific albacore 2024 assessment with KIMSA. It combines one diagnostic fit, eight cumulative full-workflow model-development fits, 100 independently refitted ensemble members, 30 starting-value trials, 20 simulation-estimation refits, five retrospective peels, five one-year index hindcasts, an average-adult-biomass likelihood profile and 5,000 reference-catch projection paths. All supplied finite results are carried into the displays; recorded numerical checks remain available in the interactive outputs.

The fitted period ends in 2022 and projections cover 2023–2042. Management summaries use 2019–2022 for recent spawning quantities and 2018–2021 for recent fishing mortality. Report resources and interactive supplements are listed in Table 1.

1 Introduction

Assessing South Pacific albacore requires a link between what fisheries observe and how the population changes. Catch records describe removals, abundance indices track relative trends, and size samples help resolve the sizes and ages affected by fishing. Biological studies provide the context needed to interpret these observations.

An integrated assessment brings these sources together in a population model. The analysis can then examine changes in recruitment, spawning potential and fishing mortality, and assess stock status against defined reference points. Differences in data coverage and uncertainty in the underlying processes are central to that interpretation.

Each update revisits the population history as well as recent stock status. New observations and revised assumptions can both change earlier estimates. A stepwise comparison helps explain those changes, while sensitivity analyses show how strongly the conclusions depend on particular choices.

2 Background

2.1 Fisheries

Longline fleets take much of the South Pacific albacore catch, while southern surface fisheries target smaller fish. The fleets include distant-water vessels and vessels based in Pacific Island countries and territories. Troll fishing around New Zealand and the subtropical convergence zone, together with historical driftnet catches, forms part of this broader fishery history (Nikolic et al., 2017; Vidal et al., 2021).

These fisheries sample different parts of the population. Their catches and size distributions reflect where and how they operate, as well as changes in fish abundance. This provides a basis for defining fisheries in the assessment.

2.2 Distribution and movement

Albacore occupy a wide range of tropical and temperate habitats. North and South Pacific populations have generally been considered separate stocks, while structure within the South Pacific remains less certain (Anderson et al., 2019; Nikolic et al., 2017). Regional differences in growth, genetics and otolith chemistry help describe that structure (Anderson et al., 2019; Macdonald et al., 2013; Williams et al., 2012).

Tag recoveries and patterns in fish size suggest movement between southern feeding grounds and warmer waters as fish grow. Smaller albacore are common in cooler waters, whereas mature fish occur farther north during the spawning season (Farley et al., 2014; Nikolic et al., 2017). Tagging evidence is uneven across the region because releases, fishing effort and reporting vary in space and time. Seasonal catch patterns provide additional information on distribution (Langley, 2004).

Habitat-based models offer hypotheses about movement and the distribution of recruitment (Lehodey et al., 2015; Senina et al., 2020). These can inform assessment alternatives alongside evidence from fisheries and biological sampling. Region boundaries provide a way to represent spatial variation; they do not necessarily delimit closed populations.

2.3 Depth use and availability to fishing

The depths occupied by albacore affect their availability to different gears. Longline catches tend to come from deeper water in warmer areas, while troll fisheries sample fish near the surface (Bigelow et al., 2006). Electronic tagging has also shown changes in depth through the day and across latitudes (Williams et al., 2015). These behaviours help explain why catchability can vary even when abundance does not.

2.4 Growth

Otolith ages and length-frequency patterns provide complementary views of growth. Surface fisheries sample younger fish, and growth slows as albacore become larger (Farley et al., 2013a; Williams et al., 2012). Differences between sexes and across the South Pacific add variation around the mean growth curve.

The interpretation of age data also depends on ageing methods and the timing of sampling within the year (Farley et al., 2021). Comparisons of growth estimates therefore need to account for which fish were sampled and how their ages were assigned. This is particularly relevant when combining age data with modes in length-frequency samples.

2.5 Reproduction

South Pacific albacore spawn in warm waters during the austral summer. Maturity and reproductive activity vary with latitude and season (Farley et al., 2013b; Farley et al., 2014). A reproductive ogive summarises how the contribution to spawning changes with fish size or age.

Converting that ogive to population spawning potential depends on the treatment of maturity,

female proportion, body mass and, where used, fecundity. The resulting quantity may be expressed as biomass or as a relative measure of reproductive output. Its definition is given in Section [4.2.4](#).

2.6 Natural mortality

Natural mortality is difficult to estimate directly from fishery data. Life history and relationships between mortality and body size can help define plausible age-dependent patterns (Lorenzen, 1996). When those patterns depend on growth, the two sets of assumptions need to be considered together. Their effects on population scale and stock status can then be examined through sensitivity analyses.

3 Data compilation

3.1 Assessment area and data period

Spatial structure links the population being assessed to the areas sampled by fisheries. Population regions describe where fish live and move; fishery strata group observations from similar fishing operations. An areas-as-fleets approach can retain differences among fishing areas within a pooled population (Waterhouse et al., 2014).

Coverage by data type, fishery and year is summarised in Figure [1](#).

3.2 Fisheries and catch

Fishery definitions group observations with similar selectivity and fishing practices. Gear, area, fleet and changes in operations can all inform these groups. Extraction fisheries account for removals, while abundance-index series describe relative population trends.

Fishery definitions and observation units are listed in Table [2](#).

3.3 Abundance indices

Catch per unit effort (CPUE) reflects both fish abundance and the way a fleet operates. Standardisation aims to account for changes in fishing practices and sampling so that the remaining trend can inform relative abundance. Spatiotemporal methods also account for spatial dependence and combine predictions over a defined area (Thorson et al., 2015). Combining fleets has helped extend the coverage of earlier albacore analyses (Vidal et al., 2021).

3.4 Length composition

Length samples describe the size distribution of fish in the catch. Observer and port-sampling programmes provide much of this information, but their coverage varies among fleets and periods. Fish measured in the same sampling event may also be more alike than fish drawn independently from the catch. Sample construction and likelihood weighting therefore need to be considered together.

3.5 Age observations

Age-length data link measured fish sizes to population age structure. A length-at-age model describes $P(L | A)$; an age-at-length model describes $P(A | L)$. The second also depends on the age distribution of the sampled fish. Bayes' rule expresses the connection:

$$P(A = a | L = l) = \frac{P(L = l | A = a) \pi_a}{\sum_b P(L = l | A = b) \pi_b},$$

where π_a represents the sampled age distribution, including selectivity where appropriate.

4 Model description

4.1 Model framework

The model connects population change to fishery removals and the observations used in fitting. Population processes determine the numbers of fish available; fishing processes determine which fish are caught; observation models describe how the data relate to those quantities. MULTIFAN-CL is an established example of this integrated approach (Fournier et al., 1998; Hampton and Fournier, 2001; Kleiber et al., 2019).

4.2 Population processes

4.2.1 Recruitment and initial population

Recruitment adds fish to the youngest modelled age class. Its definition links that age class to the timing and spatial distribution of entry into the population. Variation in recruitment can be represented around a mean or a stock-recruitment relationship.

For a Beverton-Holt relationship, steepness describes recruitment at 20% of unfished equilibrium spawning potential relative to unfished equilibrium recruitment (Francis, 1992). This parameter influences the stock's assumed capacity to replenish itself as spawning potential declines.

4.2.2 Growth and natural mortality

Growth translates population age structure into a distribution of fish lengths. A weight-length relationship then connects fish size to biomass. Variation around mean length at age is important because fish of the same age can occur in several length classes.

Natural mortality governs survival in the absence of fishing. Its time unit must be consistent with the population model, and any dependence on growth must be carried through the biological inputs.

Growth inputs and estimates are shown in Figure 2. Natural mortality at age is shown in Figure 4.

4.2.3 Movement

In a spatial model, movement redistributes fish among regions. The resulting transitions depend on age, season and the sequence of population processes within a time step. Movement rates and transition probabilities describe different parts of that calculation.

Regional movement probabilities are shown in Figure 22.

4.2.4 Reproductive potential

Spawning potential combines the population at each age or size with its contribution to reproduction. The reproductive ogive determines that contribution and the units of the resulting population measure.

The reproductive ogive is shown in Figure 3.

4.3 Fishing processes

Selectivity describes the relative susceptibility of fish of different sizes or ages to a fishery. Catchability links an index to the population it samples or, in an effort-based formulation, links effort to fishing mortality. Their parameterisation determines how fishing operations enter the population model.

Fishery selectivity at age is shown in Figure 21.

4.4 Observation models

Each observation model connects a data source to its predicted quantity and specifies how deviations contribute to the objective. For composition data, the information assigned to a sample depends on its effective sample size and the chosen likelihood. These choices affect the balance among data sources.

4.5 Parameter estimation

Fitting minimises the negative log-likelihood and any additional penalties. Automatic differentiation supplies derivatives for optimisation (Fournier et al., 2012). Numerical convergence and the information available to estimate parameters are examined in Section 5.

4.6 Stepwise development

Eight cumulative model configurations were each fitted through their complete declared doital sequence. Colours identify configurations and all trajectories use solid lines. The four panels in Figure 5 show spawning depletion, aggregate fishing mortality, recruitment and spawning potential.

4.7 Model ensemble

The ensemble contains 100 member-specific official 2024 inputs. Each member was refitted independently with KIMSA before its derived quantities were combined.

5 Model diagnostics

5.1 Starting-value trials (jitter)

Jitter trials refit the same model and data from different starting values. Agreement in final objectives and population estimates indicates that the solution is reproducible across those trials. Gradients help distinguish converged solutions from runs that stopped before reaching an optimum.

The corresponding population estimates are shown in Figure 11. Regional depletion is compared in Figure 12.

5.2 Simulation-estimation tests (self-test)

A self-test generates data from a known model and then estimates that model from the simulated observations. Comparing the estimates with their generating values reveals bias and variability in recovery under the assumed processes. Where uncertainty intervals are available, their coverage can also be examined.

Recovery of selected quantities is summarised in Figure 13. Recovery through time is shown in Figure 14.

5.3 Retrospective analysis

Retrospective analysis examines how estimates change as terminal observations are removed. Each reduced-data fit is compared with the full-data fit over their shared years (Cadigan and Farrell, 2005). Repeated shifts in the same direction can reveal a systematic pattern in recent estimates.

Population trajectories for the retrospective fits are compared in Figure 15.

5.4 Hindcast evaluation

Hindcasts assess predictions against observations withheld from fitting. At each cutoff, the model is fitted using the information available up to that point and predicts subsequent observations. The comparison measures predictive performance at the specified horizons.

Predictions and withheld observations are compared in Figure 33. Prediction errors and skill measures are summarised in Table 7.

5.5 Likelihood profiles and identifiability

A profile follows the objective as one parameter or derived quantity is fixed at a sequence of values and the remaining parameters are re-estimated. Separating the objective by data source shows where information is concentrated and where sources favour different values. Total average biomass is one possible measure of population scale for this analysis (McKechnie et al., 2017; Tremblay-Boyer et al., 2017).

The total profile and contributions by data source are shown in Figure 16. Index contributions are shown in Figure 17. Length-composition contributions are shown in Figure 18. Age-data contributions are shown in Figure 19.

5.6 Data fit

Observed abundance indices, fitted values, supplied log-scale uncertainty and standardised residuals are paired by index in Figure 6. Composition figures use the observation and prediction records carried by the diagnostic fit.

5.7 Population estimates

The diagnostic population trajectories are shown in Figure 20. Regional shares are calculated only where regional quantities sum to the supplied all-region value; they are not inferred from fishery labels.

5.8 Stock status

Terminal ensemble uncertainty is shown in Kobe and Majuro coordinates in Figure 28. The diagnostic model alone supplies the annual path in Figure 29. Management quantities and reference-point probabilities use the explicit periods stated in the summary and are reported in Table 3 and Table 4.

The reference-catch projection follows the 2024 assessment specification and spans 2023–2042. Each of 100 independently refitted ensemble members contributes 50 stochastic recruitment paths. Figure 31 joins compatible historical ensemble summaries to the projected period, marks the transition, shows 2.5%–97.5% intervals and overlays three coherent individual paths selected across the terminal depletion distribution. Regional and catch-attainment results remain in the portable result tables. Terminal quantities are listed in Table 6.

6 Discussion

6.1 Interpretation

6.2 Limitations

6.3 Conclusions and research priorities

7 Acknowledgements

8 References

- Anderson, G., Hampton, J., Smith, N., and Rico, C. (2019). Indications of strong adaptive population genetic structure in albacore tuna (*Thunnus alalunga*) in the southwest and central Pacific Ocean. *Ecol Evol*, 9(18): 10354–10364 <https://doi.org/10.1002/ece3.5554>.
- Bigelow, K., Musyl, M. K., Poisson, F., and Kleiber, P. (2006). Pelagic longline gear depth and shoaling. *Fisheries Research*, 77(2): 173–183 <https://doi.org/10.1016/j.fishres.2005.10.010>.
- Cadigan, N. G., and Farrell, P. J. (2005). Local influence diagnostics for the retrospective problem in sequential population analysis. *ICES Journal of Marine Science: Journal du Conseil*, 62(2): 256–265 <https://doi.org/10.1016/j.icesjms.2004.11.015>.
- Farley, J. H., Hoyle, S. D., Eveson, J. P., Williams, A. J., Davies, C. R., and Nicol, S. J. (2014). Maturity Ogives for South Pacific Albacore Tuna (*Thunnus alalunga*) That Account for Spatial and Seasonal Variation in the Distributions of Mature and Immature Fish. *PLoS ONE*, 9(1): e83017 <https://doi.org/10.1371/journal.pone.0083017>.
- Farley, J., Krusic-Golub, K., and Eveson, P. (2021). Ageing of South Pacific Albacore: (Project 106). WCPFC-SC17-SA-IP-10.
- Farley, J. H., Williams, A. J., Clear, N. P., Davies, C. R., and Nicol, S. J. (2013a). Age estimation and validation for South Pacific albacore (*Thunnus alalunga*). *Journal of Fish Biology*, 82(5): 1523–1544 <https://doi.org/10.1111/jfb.12077>.
- Farley, J. H., Williams, A. J., Hoyle, S. D., Davies, C. R., and Nicol, S. J. (2013b). Reproductive Dynamics and Potential Annual Fecundity of South Pacific Albacore Tuna (*Thunnus alalunga*). *PLoS ONE*, 8(4): e60577.
- Fournier, D. A., Hampton, J., and Sibert, J. R. (1998). MULTIFAN-CL: A length-based, age-structured model for fisheries stock assessment, with application to South Pacific albacore, *Thunnus alalunga*. *Canadian Journal of Fisheries and Aquatic Sciences*, 55: 2105–2116.
- Fournier, D. A., Skaug, H. J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M. N., Nielson, A., and Sibert, J. (2012). AD Model Builder: Using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optimization Methods and Software*, 27(2): 233–249.
- Francis, R. I. C. C. (1992). Use of risk analysis to assess fishery management strategies: A case study using orange roughy (*Hoplostethus atlanticus*) on the Chatham Rise, New Zealand. *Canadian Journal of Fisheries and Aquatic Sciences*, 49: 922–930 <https://doi.org/10.1139/f92-102>.
- Hampton, J., and Fournier, D. A. (2001). A spatially-disaggregated, length-based, age-structured

- population model of yellowfin tuna (*Thunnus albacares*) in the western and central Pacific Ocean. *Marine and Freshwater Research*, 52: 937–963.
- Kleiber, P., Fournier, D., Hampton, J., Davies, N., Bouye, F., and Hoyle, S. (2019). MULTIFAN-CL User’s Guide.
- Langley, A. (2004). An examination of the influence of recent oceanographic conditions on the catch rate of albacore in the main domestic longline fisheries. Working Paper SA-4., 17th Standing Committee on Tuna; Billfish. 9–18 August 2004. Majuro, Republic of Marshall Islands.
- Lehodey, P., Senina, I., Nicol, S., and Hampton, J. (2015). Modelling the impact of climate change on South Pacific albacore tuna. *Deep Sea Research Part II: Topical Studies in Oceanography*, 113: 246–259 <https://doi.org/10.1016/j.dsr2.2014.10.028>.
- Lorenzen, K. (1996). The relationship between body weight and natural mortality in juvenile and adult fish: A comparison of natural ecosystems and aquaculture. *Journal of Fish Biology*, 49(4): 627–642 <https://doi.org/10.1111/j.1095-8649.1996.tb00060.x>.
- Macdonald, J. I., Farley, J. H., Clear, N. P., Williams, A. J., Carter, T. I., Davies, C. R., and Nicol, S. J. (2013). Insights into mixing and movement of South Pacific albacore *Thunnus alalunga* derived from trace elements in otoliths. *Fisheries Research*, 148: 56–63 <https://doi.org/10.1016/j.fishres.2013.08.004>.
- McKechnie, S., Pilling, G., and Hampton, J. (2017). Stock assessment of bigeye tuna in the western and central Pacific Ocean. WCPFC-SC13-2017/SA-WP-05, Rarotonga, Cook Islands, 9–17 August 2017.
- Nikolic, N., Morandeau, G., Hoarau, L., West, W., Arrizabalaga, H., Hoyle, S., Nicol, S. J., Bourjea, J., Puech, A., Farley, J. H., Williams, A. J., and Fonteneau, A. (2017). Review of albacore tuna, *Thunnus alalunga*, biology, fisheries and management. *Rev Fish Biol Fisheries*, 27(4): 775–810 <https://doi.org/10.1007/s11160-016-9453-y>.
- Senina, I. N., Lehodey, P., Hampton, J., and Sibert, J. (2020). Quantitative modelling of the spatial dynamics of South Pacific and Atlantic albacore tuna populations. *Deep Sea Research Part II: Topical Studies in Oceanography*, 175: 104667 <https://doi.org/10.1016/j.dsr2.2019.104667>.
- Thorson, J. T., Shelton, A. O., Ward, E. J., and Skaug, H. J. (2015). Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. *Ices Journal of Marine Science*, 72(5): 1297–1310 <https://doi.org/10.1093/icesjms/fsu243>.
- Tremblay-Boyer, L., McKechnie, S., Pilling, G., and Hampton, J. (2017). Stock assessment of yellowfin tuna in the Western and Central Pacific Ocean. WCPFC-SC13-2017/SA-WP-06, Rarotonga, Cook Islands, 9-17 August 2017.
- Vidal, T., Castillo Jordan, C., Peatman, T., Ducharme-Barth, N., Xu, H., Williams, P., Lennert-Cody, C., and Hamer, P. (2021). Background analysis and data inputs for the 2021 South Pacific albacore tuna stock assessment. WCPFC-SC17-SA-IP-03.
- Waterhouse, L., Sampson, D. B., Maunder, M., and Semmens, B. X. (2014). Using areas-as-fleets selectivity to model spatial fishing: Asymptotic curves are unlikely under equilibrium conditions. *Fisheries Research*, 158: 15–25 <https://doi.org/10.1016/j.fishres.2014.01.009>.

- Williams, A. J., Allain, V., Nicol, S. J., Evans, K. J., Hoyle, S. D., Dupoux, C., Vourey, E., and Dubosc, J. (2015). Vertical behavior and diet of albacore tuna (*Thunnus alalunga*) vary with latitude in the South Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 113: 154–169 <https://doi.org/10.1016/j.dsr2.2014.03.010>.
- Williams, A. J., Farley, J. H., Hoyle, S. D., Davies, C. R., and Nicol, S. J. (2012). Spatial and sex-specific variation in growth of albacore tuna (*Thunnus alalunga*) across the South Pacific Ocean. *PLoS ONE*, 7(6): e39318.

9 Tables

Table 1: Assessment resources and output provenance.

Resource	Link	Purpose
Assessment report and interactive outputs	https://kyuhank.github.io/assessment-workflow-demo	Published demonstration outputs
Official 2024 ensemble design	PacificCommunity/ofp-sam-alb-2024-ensemble	Read-only source of the 100 member input design
Official 2024 assessment write-up	PacificCommunity/ofp-sam-alb-2024-writeup	Read-only assessment context and projection specification

Table 2: Fishery definitions and observation units.

Fishery	Region	Role	Years	Unit
1	1	Catch	1954–2022	fish
2	1	Catch	1954–2022	fish
3	1	Catch	1954–2022	fish
4	1	Catch	1954–2022	fish
5	1	Catch	1992–2022	fish
6	1	Catch	1982–2022	fish
7	1	Catch	1987–2022	fish
8	1	Catch	1954–2022	fish
9	1	Catch	1984–2022	fish
10	1	Catch	1985–2022	fish
11	1	Catch	1982–2022	t
12	1	Catch	1987–2022	t
13	1	Catch	1983–1990	t
14	2	Catch	1957–2022	fish
15	2	Catch	1961–2021	fish
16	2	Catch	1963–2021	fish
17	2	Catch	1986–2006	t
18	1	Index	1954–2022	relative
19	1	Index	1992–2022	relative

Fishery	Region	Role	Years	Unit
20	2	Index	1954–2022	relative

Table 3: Management quantities and supplied uncertainty summaries.

Quantity	Period	Unit	Mean	Median	Minimum	10%	90%	Maximum
Aggregate F	2018–2021	per year	0.0728	0.0666	0.0222	0.0332	0.108	0.156
Aggregate F	2022	per year	0.0692	0.0619	0.0206	0.0314	0.104	0.151
BMSY	Reference point	t	785000	662000	330000	423000	1290000	2210000
Dynamic unfished spawning potential	2019–2022	t	678000	668000	601000	624000	748000	813000
Dynamic unfished spawning potential	2022	t	705000	690000	616000	651000	799000	901000
F/F_{MSY}	2018–2021	ratio	0.458	0.448	0.205	0.253	0.638	0.85
F/F_{MSY}	2022	ratio	0.435	0.429	0.189	0.232	0.613	0.818
FMSY	Reference point	per year	0.155	0.159	0.106	0.123	0.182	0.203
MSY	Reference point	t/year	115000	103000	62900	75100	177000	235000
$SB/SB_{F=0}$	2019–2022	ratio	0.513	0.512	0.274	0.39	0.668	0.742
$SB/SB_{F=0}$	2022	ratio	0.548	0.542	0.292	0.422	0.703	0.781
SB/SB_{MSY}	2019–2022	ratio	3.22	3.03	1.55	2.1	4.89	6.01
SB/SB_{MSY}	2022	ratio	3.58	3.32	1.67	2.28	5.47	6.97
SBMSY	Reference point	t	114000	114000	60600	83600	146000	167000
Spawning potential	2019–2022	t	349000	335000	222000	262000	488000	604000
Spawning potential	2022	t	389000	369000	238000	287000	546000	704000

Table 4: Stock-status probabilities for the declared reference points.

Condition	Period	Probability
$F/F_{MSY} > 1$	2018–2021	0
$SB/SB_{F=0} < 0.20$	2019–2022	0
$SB/SB_{MSY} < 1$	2019–2022	0
$F/F_{MSY} > 1$	2022	0
$SB/SB_{F=0} < 0.20$	2022	0
$SB/SB_{MSY} < 1$	2022	0

Table 5: Model reference points and their definitions.

Quantity	Estimate	Unit
MSY	86055	t/year
FMSY	0.14846	per year
BMSY	579660	t
SBMSY	136940	t

Table 6: Projected quantities by scenario and reporting period.

Scenario	Period	Quantity	Median	95% interval
Reference catch	2042	Spawning Biomass	273700	7.053e+04– 6.297e+05
Reference catch	2042	Dynamic Spawning Depletion	0.46049	0.1184–0.8391
Reference catch	2042	Fishing Mortality Proxy	0.084509	0.02722–0.2258
Reference catch	2042	Recruitment	90321000	1.728e+07– 5.534e+08
Reference catch	2042	Total Biomass	943750	2.862e+05– 2.997e+06
Reference catch	2042	Catch	79733	5.664e+04– 8.432e+04

Table 7: Held-out prediction bias (predicted minus observed), RMSE and MASE by series and forecast lead. N counts scored forecasts; MASE uses each cutoff’s training naive-error scale.

Series	Lead	N	Bias	RMSE	Unit
F18_R1_G18	1	5	0.013635	0.075785	index
F19_R1_G19	1	5	0.43361	0.70298	index
F20_R2_G20	1	5	-0.13701	0.21829	index

10 Figures

DEMONSTRATION




Figure 1: Observation availability by data type and fishery. Coloured cells indicate years with observations.




Figure 2: Mean length at the start of each model age class.




Figure 3: Model reproductive coefficient at age.




Figure 4: Annual instantaneous natural mortality at age.




Figure 5: Stepwise model trajectories.




Figure 6: Abundance-index fits and log residuals by series. Shading: central 95% conditional observation intervals. Residuals use the supplied log-scale standard deviations.




Figure 7: Abundance-index fits and log residuals by series. Shading: central 95% conditional observation intervals. Residuals use the supplied log-scale standard deviations.




Figure 8: Length-composition fits by fishery. Observed sample counts (bars) and predictions (lines), pooled over years.




Figure 9: Length-composition fits by region. Observed sample counts (bars) and predictions (lines), pooled over years.




Figure 10: Observed and predicted age distributions conditional on length. Cells: predictions; circles: observations; solid and dashed lines: predicted and observed mean age.




Figure 11: Population trajectories across models (thin lines) and the reference model (red).




Figure 12: Spawning depletion by model and region, relative to each model's dynamic unfished reference.




Figure 13: Relative recovery error in terminal quantities for simulated-data refits. All finite refits are shown.




Figure 14: Population trajectories across models (thin lines) and the reference model (red).




Figure 15: Population trajectories for the full-data fit and retrospective peels. Points mark terminal years; rho is Mohn's mean terminal relative difference for peels passing the recorded numerical checks.




Figure 16: Total negative log-likelihood and component profiles. Each curve is centered on its own minimum. Reproductively weighted biomass summed over regions and averaged across model periods in 1954-2022.




Figure 17: Negative log-likelihood profiles for abundance-index components. Each curve is centered on its own minimum. Reproductively weighted biomass summed over regions and averaged across model periods in 1954-2022.




Figure 18: Negative log-likelihood profiles for length-composition components. Each curve is centered on its own minimum. Reproductively weighted biomass summed over regions and averaged across model periods in 1954-2022.




Figure 19: Negative log-likelihood profiles for conditional age-at-length components. Each curve is centered on its own minimum. Reproductively weighted biomass summed over regions and averaged across model periods in 1954-2022.




Figure 20: Estimated population trajectories.




Figure 21: Fishery selectivity at age.




Figure 22: Probability of movement from source to destination region per movement event.




Figure 23: Annual recruitment.




Figure 24: Recruitment by model region.




Figure 25: Regional shares of population quantities.




Figure 26: Total biomass under estimated fishing (solid) and the dynamic no-fishing counterfactual (dashed), by region and overall.




Figure 27: Population trajectories across ensemble models (thin lines) and the reference model (red).




Figure 28: Kobe (left) and Majuro (right) status in 2022. Open points: model estimates; percentages: weighted status probabilities. Contours: 50%, 80% and 95% joint highest-density regions of structural model spread. Majuro probabilities distinguish depletion below the limit, and F above or below F_{MSY} when the depletion criterion is met.




Figure 29: Stock status relative to the supplied spawning and fishing reference points.




Figure 30: Equilibrium yield across fishing-mortality multipliers.




Figure 31: Historical and projected quantities by scenario with supplied uncertainty intervals. Central 95% intervals (2.5% to 97.5% quantiles across Monte Carlo ensemble members and recruitment replicates). Thin lines are a declared subset of individual projection paths.




Figure 32: Historical and projected quantities by scenario with supplied uncertainty intervals. Central 95% intervals (2.5% to 97.5% quantiles across Monte Carlo ensemble members and recruitment replicates). Thin lines are a declared subset of individual projection paths.




Figure 33: Observed values (black) and rolling 1-year-ahead predictions (teal); shading marks the held-out evaluation period. Right panels compare predictions with observations against the 1:1 line. n is the number of evaluated predictions.




Figure 34: Observed values (black) and rolling 1-year-ahead predictions (teal); shading marks the held-out evaluation period. Right panels compare predictions with observations against the 1:1 line. n is the number of evaluated predictions.

11 Supplementary outputs

[Diagnostic model — interactive viewer](#)

[Diagnostic model — figures and tables](#)

[ALB 2024 reconstructable model-choice stepwise — interactive viewer](#)

[ALB 2024 reconstructable model-choice stepwise — figures and tables](#)

[Monte Carlo model ensemble — interactive viewer](#)

[Monte Carlo model ensemble — figures and tables](#)

[Starting-value trials — interactive viewer](#)

[Starting-value trials — figures and tables](#)

[Simulation-estimation tests — interactive viewer](#)

[Simulation-estimation tests — figures and tables](#)

[Retrospective analysis — interactive viewer](#)

[Retrospective analysis — figures and tables](#)

[Index hindcast — interactive viewer](#)

[Index hindcast — figures and tables](#)

[Average adult biomass likelihood profile — interactive viewer](#)

[Average adult biomass likelihood profile — figures and tables](#)

[Reference catch projection — interactive viewer](#)

[Reference catch projection — figures and tables](#)

Prepared with KIMSA companion tools. Developed by Kyuhan Kim.

DEMONSTRATION