

Tail-Index Sure Independence Screening for Ultra-High-Dimensional Heavy-Tailed Regression

Anonymous manuscript

Abstract

We propose a marginal screen for the covariates that determine a positive conditional tail index, designed for problems where the dimension p is large relative to the number of observations informative about the tail. The screen rests on one geometric fact: after projection onto a single coordinate, the conditional distribution is a mixture whose tail is governed by its heaviest component, so the projected tail index is the upper envelope of the tail-index surface along the corresponding fibre. Inactive coordinates produce flat envelopes at the global maximum γ^* ; active coordinates are detectable exactly when their fibres miss the global argmax set on a set of positive measure. Ranking coordinates by the average projected index therefore separates detectable active variables from noise. We estimate the envelope with empirical-rank local Hill profiles, and prove that the maximal score error over p coordinates is $O_{\mathbb{P}}(\sqrt{\log(pn)/(nah)})$ plus a bias term negligible relative to the population gap, where α is the local tail fraction and h the rank bandwidth; nah is the effective local extreme sample size. Sure screening and exact recovery follow when $\log(pn) = o(nah\Delta_{\min}^2)$. The algorithm costs $O(pn \log n)$. A rank-aggregated version of the screen, taking the minimum rank over a block of nine neighbouring tunings, stabilizes the ranking against the choice of the smoothing parameters. Simulations across $p \in \{500, 1000, 2000\}$, including a model whose scale is driven by a correlated cluster of twenty inactive covariates, show when screening on the tail index differs from screening on a fixed high quantile: the scale cluster distracts fixed-quantile screens at every level, while the aggregated tail screen recovers the tail-index-active set reliably. An application to violent-crime rates in 1,993 U.S. communities, with $p = 99$ continuous predictors against an effective tail sample of about 200, illustrates the method in its intended regime. The uniform projected-quantile conditions are stated directly; deriving them from a fully primitive growing-dimensional model remains open.

1 Introduction

Let (X, Y) be a random pair with $Y > 0$ and $X \in \mathbb{R}^p$, and suppose that, conditionally on $X = x$,

$$\frac{\mathbb{P}(Y > ty \mid X = x)}{\mathbb{P}(Y > y \mid X = x)} \longrightarrow t^{-1/\gamma(x)}, \quad y \rightarrow \infty, \quad t > 0, \quad (1)$$

for a positive function γ , the conditional tail index. Estimating γ is the first step toward extreme conditional quantiles and tail probabilities [6, 12], and it suffers from a particularly severe curse of dimensionality: local estimation must find observations near x and then keep only an upper fraction of their responses. A one-dimensional window of width h combined with a tail fraction α leaves an effective sample of order nah , not n [5, 10]. When p is large, even moderate sample sizes are effectively small.

This paper asks the screening question: among p candidate covariates, possibly with p growing with n , which ones does γ actually depend on? Sure independence screening [7] answers such questions by ranking coordinates through inexpensive marginal statistics and keeping a moderate

number for subsequent joint analysis. Model-free variants exist for general dependence [13] and for fixed conditional quantiles [11], but the limiting exponent in (1) is a different target: a covariate can move the 0.95 quantile of Y without affecting γ , and conversely. Our simulations make this distinction concrete.

The screen we propose rests on a single geometric mechanism. Transform each continuous covariate to $U_j = F_j(X_j)$ and let $\xi_j(u)$ denote the tail index of Y given $U_j = u$ alone. Conditioning on one coordinate leaves the others free to range over the fibre $\{u\} \times [0, 1]^{p-1}$, so the projected distribution is a mixture, and a mixture of heavy tails is governed by its heaviest component. Under a full-support condition,

$$\xi_j(u) = \max_{v \in [0, 1]^{p-1}} \gamma(u, v) : \quad (2)$$

the projected index is the upper envelope of the tail-index surface along the fibre. If coordinate j is inactive, every fibre reaches the global maximum $\gamma^* = \max_x \gamma(x)$ and the envelope is flat at γ^* . If coordinate j is active and some of its fibres miss the global argmax set, the envelope dips below γ^* there. Averaging the envelope over a trimmed interval gives a score Ψ_j whose gap $\Delta_j = \gamma^* - \Psi_j$ is zero for every inactive coordinate and positive for every detectable active one. Ranking coordinates by increasing Ψ_j is the screen.

The same geometry states the method's limitation. An active coordinate is invisible when every one of its fibres still meets the global argmax set; this happens, for instance, for $\gamma(u_1, u_2) = \gamma_0 - (u_1 - u_2)^2$, whose argmax is the diagonal. Detectability is a property of the argmax projections, and we characterize it exactly rather than assuming a generic signal-strength condition.

Estimation follows the envelope. For each coordinate we slide a window over the empirical marginal ranks, apply a Hill statistic to the upper α -fraction of the responses in the window, and average the resulting profile. Empirical ranks give every coordinate the same deterministic window geometry and remove the marginal scales. The price is a rank-displacement term, which we control simultaneously over p coordinates by a Dvoretzky–Kiefer–Wolfowitz bound; it is negligible when $\log(pn) = o(nh^2)$. The main result bounds the maximal score error by

$$O_{\mathbb{P}} \left(\sqrt{\frac{\log(pn)}{n\alpha h}} \right) + o(\Delta_{\min}),$$

where $\Delta_{\min}$ is the weakest population gap, and yields sure screening when $\log(pn) = o(n\alpha h\Delta_{\min}^2)$: ultra-high dimension is affordable exactly when the local extreme count dominates the logarithm of the dimension. A blockwise implementation costs $O(pn \log n)$ time.

Our construction adapts the projected-tail framework of Gardes and Podgorny [9], who estimate a low-dimensional central tail-index subspace by minimizing an average of projected local Hill estimates over projection matrices, in fixed dimension. We keep their projection semantics and local Hill estimator, and change what is needed for screening: the projections are the p coordinates themselves, p may diverge, windows are built from empirical ranks with deterministic local counts, and uniformity over a compact matrix class is replaced by union bounds over coordinates and window states. Their density assumptions on the covariate are not needed here; the empirical-rank construction replaces them. Among screening proposals, Yoshida and Umezu [19] screen on covariate-dependent extreme-value indices through kernel conditional Pickands contrasts in a large-bandwidth regime, and parametric tail-index regression [18], extended to high dimensions with regularization by Sasaki et al. [17], exploits a parametric link when one is available. Tail dimension reduction methods [1, 2, 4, 8] target related but distinct conditional-tail objects.

The theoretical scope deserves a clear statement. The primitive model (1) identifies the projected index (2), but does not by itself provide the uniform projected-quantile regularity needed for growing-dimensional concentration: projection can create additional logarithmic slow variation even from

exact Pareto conditionals. We therefore state the required conditions directly on the projected quantiles and leave their derivation from primitive assumptions open.

Section 2 develops the population theory. Section 3 defines the estimator and states the screening results. Section 4 presents the simulation models, the tuning study together with a rank-aggregated version of the screen built on a block of nine neighbouring tunings, and a comparison with competing screens at $p \in \{500, 1000, 2000\}$ that covers, in the same tables, the sensitivity to the dimension and to the quantile level of fixed-quantile screening. Section 5 applies the aggregated screen to violent-crime rates in U.S. communities. Section 6 discusses limitations. Proofs are in Appendix A.

2 Population theory

2.1 Model and assumptions

All quantities below may depend on n : the dimension $p = p_n$, the tail-index function γ , the active set A , its size $s = |A|$, and later the bandwidth h , the tail fraction α and the retained size d . The index n is suppressed whenever no ambiguity arises.

Notation. A positive measurable function f is slowly varying at infinity if $f(ty)/f(y) \rightarrow 1$ as $y \rightarrow \infty$ for every $t > 0$, and a survival function $\bar{F}$ is regularly varying with index $-1/\xi < 0$ if $\bar{F}(ty)/\bar{F}(y) \rightarrow t^{-1/\xi}$ for every $t > 0$. Generalized inverses are $F^{-1}(t) = \inf\{y : F(y) \geq t\}$. For $u \in [0, 1]$ and $v \in [0, 1]^{p-1}$, $\iota_j(u, v) \in [0, 1]^p$ denotes the vector with u in coordinate j and v in the remaining coordinates, in order; we abbreviate $\gamma(u, v) = \gamma\{\iota_j(u, v)\}$ when j is clear from context. Norms on $[0, 1]^A$ are Euclidean. All asymptotic statements concern a triangular array indexed by n , and $Z_n = O_{\mathbb{P}}(a_n)$ means $\lim_{C \rightarrow \infty} \limsup_n \mathbb{P}(|Z_n| > Ca_n) = 0$.

The observed data are i.i.d. pairs (X_i, Y_i) with $Y_i > 0$ and $X_i \in \mathbb{R}^p$ with continuous marginal distribution functions $F_1, \dots, F_p$. The theory is formulated on the latent probability-integral transforms $U_{ij} = F_j(X_{ij})$, so that $U_i \in [0, 1]^p$ has uniform one-dimensional marginals and unrestricted cross-coordinate dependence; the transforms are never needed by the algorithm, because the ranks of $X_{1j}, \dots, X_{nj}$ coincide with the ranks of $U_{1j}, \dots, U_{nj}$ almost surely. With slight abuse of notation we write γ for the tail-index function on the cube, $\gamma(u) = \gamma_X\{F_1^{-1}(u_1), \dots, F_p^{-1}(u_p)\}$, where γ_X is the function appearing in (1).

(C1) *Conditional regular variation.* For every u , the conditional survival function factorizes as

$$\bar{F}(y \mid u) = \mathbb{P}(Y > y \mid U = u) = y^{-1/\gamma(u)} L(y, u),$$

where $L(\cdot, u)$ is slowly varying at infinity, $\gamma : [0, 1]^p \rightarrow (0, \infty)$ is continuous, and $0 < \underline{\gamma} \leq \gamma(u) \leq \Gamma < \infty$ uniformly in n and u .

(C2) *Uniform tail regularity.* There is a function c with $0 < c_- \leq c(u) \leq c_+ < \infty$ such that $\sup_u |L(y, u)/c(u) - 1| \rightarrow 0$ as $y \rightarrow \infty$.

(C1) states that each conditional law is heavy-tailed with index $\gamma(u)$, as in (1); (C2) strengthens the slow variation of L to uniform convergence toward a bounded positive limit, the form needed to control mixtures of the conditional laws.

The active set A collects the coordinates on which γ genuinely depends: $j \in A$ if and only if there exist u, v differing only in coordinate j with $\gamma(u) \neq \gamma(v)$. Iterating over the complementary coordinates shows that $\gamma(u) = g(u_A)$ for some function g , and that A is the smallest coordinate set with this property. Membership in A concerns the limiting exponent only, so a coordinate may shift the body of the conditional distribution, or its second-order behaviour, without being active.

Conditions (C1)–(C2) parallel conditions (C.1)–(C.2) of Gardes and Podgorny [9], except that we do not assume the law of $\gamma(U)$ to be absolutely continuous; the projection argument below handles atoms, which arise as soon as one coordinate is inactive.

Fix a trimming parameter $0 < \varepsilon < 1/2$ and write $I_\varepsilon = [\varepsilon, 1 - \varepsilon]$. For each coordinate j , fix a pointwise conditional distribution $K_j(u, dv)$ of U_{-j} given $U_j = u$, and define the conditional law of Y given $U_j = u$ to be compatible with it through the disintegration identity

$$\mathbb{P}(Y > y \mid U_j = u) = \int \mathbb{P}\{Y > y \mid U = (u, v)\} K_j(u, dv);$$

without such pointwise versions the statements below hold for almost every u . Define

$$\xi_j(u) = \text{ess sup}_{v \sim K_j(u, \cdot)} \gamma(u, v),$$

the essential supremum of $v \mapsto \gamma(u, v)$ under $K_j(u, \cdot)$; Proposition 2.1 shows that the conditional law of Y given $U_j = u$ is regularly varying with index $-1/\xi_j(u)$, so $\xi_j(u)$ is its tail index. One further condition is used only where stated:

(S) *Full fibre support.* For every j and $u \in I_\varepsilon$, the support of $K_j(u, \cdot)$ is all of $[0, 1]^{p-1}$. A continuous joint density, positive on the interior cube, is sufficient.

2.2 Projection turns the surface into an envelope

Conditioning on $U_j = u$ mixes the conditional laws of Y over the fibre $\{u\} \times [0, 1]^{p-1}$. Heuristically, the slowest-decaying component of a mixture of heavy tails dictates the tail of the mixture, so the projected index should equal the largest tail index available on the fibre, and a localized region of heavy tails should survive projection intact. The following proposition makes this precise.

For a sub- σ -field $\mathcal{G}$, the conditional essential supremum $\text{ess sup}\{\gamma(U) \mid \mathcal{G}\}$ is the smallest $\mathcal{G}$ -measurable random variable dominating $\gamma(U)$ almost surely.

Proposition 2.1 (Projection). *Assume (C1)–(C2) and let $\mathcal{G} \subseteq \sigma(U)$. Then $\mathbb{P}(Y > y \mid \mathcal{G})$ is almost surely regularly varying with index $-1/\xi$, where $\xi = \text{ess sup}\{\gamma(U) \mid \mathcal{G}\}$. In particular $\xi_j(u)$ is the conditional essential supremum of $\gamma(U)$ given $U_j = u$. Under (S),*

$$\xi_j(u) = \max_{v \in [0, 1]^{p-1}} \gamma(u, v), \quad u \in I_\varepsilon. \quad (3)$$

The essential-supremum identity requires neither full support nor absolute continuity; (S) enters only to replace the essential supremum by the maximum over the whole fibre. If dependence removes part of a fibre, the maximum runs over the conditional support actually available at u , and the population target changes accordingly.

2.3 Score, gap, and detectability

Write $\gamma^* = \max_u \gamma(u)$ for the global maximum and

$$\Psi_j = \frac{1}{1 - 2\varepsilon} \int_{I_\varepsilon} \xi_j(u) du, \quad \Delta_j = \gamma^* - \Psi_j, \quad \Delta_{\min} = \min_{j \in A} \Delta_j. \quad (4)$$

The screen ranks coordinates by increasing Ψ_j and keeps the d smallest. Small scores indicate activity: all inactive coordinates share the flat baseline γ^* , while fixing a detectable active coordinate lowers the attainable envelope. The algorithm uses only the ordering of the Ψ_j and never estimates γ^* .

Let $M = \arg \max_u \gamma(u)$, a nonempty compact set, and let $\pi_j(M)$ be its projection onto coordinate j .




Figure 1: Projected envelopes. Left: a monotone active coordinate depresses the envelope below the inactive baseline γ^* . Right: in the diagonal example every fibre meets the argmax set, so the envelope of an active coordinate stays flat.

Proposition 2.2 (Detectability). *Assume (C1)–(C2) and (S).*

1. If $j \notin A$, then $\xi_j \equiv \gamma^*$ on I_ε and $\Delta_j = 0$.
2. For every j and $u \in I_\varepsilon$: $\xi_j(u) = \gamma^*$ if and only if $u \in \pi_j(M)$.
3. $\Delta_j > 0$ if and only if $I_\varepsilon \setminus \pi_j(M)$ has positive Lebesgue measure.

If the restriction of γ to the active coordinates has a finite set of maximizers, every active coordinate is detectable.

Claim 3 separates two failure modes that finite-sample experiments tend to blur. When $\pi_j(M)$ covers I_ε , no sample size makes coordinate j identifiable from this score; the diagonal example $\gamma(u_1, u_2) = \gamma_0 - (u_1 - u_2)^2$, with $M = \{(t, t) : 0 \leq t \leq 1\}$, is of this kind, and both of its active coordinates are invisible. When $\pi_j(M)$ misses a set of positive measure, detection is a question of estimation accuracy, quantified by Section 3. Figure 1 contrasts the two situations.

Proposition 2.3 (Quantitative margin). *Assume (C1)–(C2), (S), and that for some $u_A^* \in [0, 1]^A$, $c > 0$, $\nu > 0$,*

$$\gamma(u) \leq \gamma^* - c \|u_A - u_A^*\|^\nu \quad \text{for all } u.$$

Then, for $j \in A$,

$$\gamma^* - \xi_j(u) \geq c |u - u_j^*|^\nu, \quad \Delta_j \geq c \mathbb{E}(|V - u_j^*|^\nu), \quad V \sim \text{Unif}(I_\varepsilon).$$

For $\nu = 2$ the right-hand side is at least $c\{(1 - 2\varepsilon)^2/12 + (1/2 - u_j^)^2\}$.*

Two consequences of (3) help with examples. If $\gamma(u) = G(\sum_{\ell \in A} g_\ell(u_\ell))$ with G strictly increasing, then $\xi_j(u) = G(g_j(u) + \sum_{\ell \in A, \ell \neq j} \max g_\ell)$, and $\Delta_j > 0$ whenever g_j lies below its maximum on a subset of I_ε of positive measure. If $\partial_j \gamma \geq c_j > 0$ throughout the cube, arbitrary interactions are allowed and $\Delta_j \geq c_j/2$.

For the benchmark $\gamma(u) = \frac{1}{2} \exp(-u_1 - u_2 - u_3 - u_4)$ used in Section 4, $\xi_j(u) = e^{-u}/2$ for $j \leq 4$ and $\xi_j \equiv 1/2$ otherwise, so $\Delta_j \approx 0.186$ at $\varepsilon = 0.05$. The gap is comfortable, yet estimating $\xi_j(u)$ near its maximum requires sample points whose remaining three active coordinates are simultaneously small; the simulations keep population detectability and finite-sample accessibility clearly separated.

3 Estimation and screening theory

3.1 Empirical-rank local Hill scores

The estimator follows the envelope construction coordinate by coordinate. Let R_{ij} be the rank of X_{ij} among $X_{1j}, \dots, X_{nj}$ and $\widehat{U}_{ij} = R_{ij}/(n+1)$. For $u \in I_\varepsilon$, form the rank window

$$\mathcal{W}_j(u) = \{i : |\widehat{U}_{ij} - u| \leq h\}, \quad M_j(u) = |\mathcal{W}_j(u)|, \quad k_j(u) = \lfloor \alpha M_j(u) \rfloor,$$

and, writing $Y_{j,u,(1)} \leq \dots \leq Y_{j,u,(M_j(u))}$ for the local order statistics, the local Hill estimate

$$\widehat{\xi}_j(u) = \frac{1}{k_j(u)} \sum_{r=1}^{k_j(u)} \log \frac{Y_{j,u,(M_j(u)-r+1)}}{Y_{j,u,(M_j(u)-k_j(u))}}, \quad (5)$$

defined when $\alpha M_j(u) > 1$. This is the local-fraction Hill statistic of Gardes and Podgorny [9, Definition 5] evaluated on one-coordinate rank windows. The score averages the profile over the deterministic grid $G = \{r/(n+1) : r = 1, \dots, n\} \cap I_\varepsilon$,

$$\widehat{\Psi}_j = \frac{1}{|G_j^{\text{val}}|} \sum_{u \in G_j^{\text{val}}} \widehat{\xi}_j(u), \quad G_j^{\text{val}} = \{u \in G : \alpha M_j(u) > 1\}, \quad (6)$$

with $\widehat{\Psi}_j = +\infty$ when no grid point is usable, so that underpopulation can never favour a coordinate. Because the window counts $M_j(u)$ depend on the ranks only through the grid position, G_j^{val} is in fact the same set for every j ; under the conditions of Theorem 3.1 every grid point is usable and $G_j^{\text{val}} = G$. The screened set $\widehat{A}_d$ collects the d smallest scores, ties broken by coordinate index.

The two localization parameters play different roles: h fixes how many observations resemble the target rank position, roughly $2nh$ of them, and α fixes how many of their largest responses enter the Hill statistic, roughly $2n\alpha h$. Rank windows make these counts deterministic up to boundary effects, which is what permits exact finite-sample union bounds over a diverging number of coordinates.

3.2 Assumptions

Ranks displace the unobserved transforms U_{ij} , windows displace the conditioning point, and the Hill statistic sees only the upper α -fraction of a projected distribution. The assumptions bound these three effects uniformly over the coordinates. Throughout this section every active coordinate is assumed detectable, $\Delta_{\min} > 0$ for every n , so that the conditions below are well posed. Let $Q_j(\cdot, u)$ denote the generalized upper conditional quantile of Y given $U_j = u$, $Q_j(s, u) = F_{Y|j}^{-1}(1 - s^{-1} | u)$ with $F^{-1}(t) = \inf\{y : F(y) \geq t\}$, and let $\tau_n = (pn)^{-2}$.

(E1) *Quantile representation and regularity of the projected index.* For every $j \leq p$ and every $u \in [0, 1]$ the conditional distribution function $F_{Y|j}(\cdot | u)$ is continuous, so that $V_{ij} = 1 - F_{Y|j}(Y_i | U_{ij})$ is conditionally standard uniform and $Y_i = Q_j(1/V_{ij}, U_{ij})$ almost surely. Moreover the projected indices are equicontinuous on I_ε , uniformly over coordinates and over n ,

$$\omega_\xi(r) = \max_{j \leq p} \sup_{u, v \in I_\varepsilon, |u-v| \leq r} |\xi_j(u) - \xi_j(v)| \longrightarrow 0 \quad (r \rightarrow 0),$$

and satisfy the grid-scale regularity condition $\omega_\xi(1/n) = o(\Delta_{\min})$.

(E2) *Uniform tail regularity.* Writing $Q_j(s, u) = s^{\xi_j(u)} \ell_j(s, u)$ with $\ell_j(\cdot, u)$ slowly varying,

$$\max_{j \leq p} \sup_{u \in I_\varepsilon} \sup_{s \geq (3\alpha)^{-1}, t > 1} \frac{|\log\{\ell_j(ts, u)/\ell_j(s, u)\}|}{\log t} = o(\Delta_{\min}).$$

(E3) *Uniform local quantile regularity.* There exists a constant $C_0 > 1$ such that

$$\max_{j \leq p} \sup_{u \in I_\varepsilon} \sup_{v \in [0, 1], |u-v| \leq C_0 h} \sup_{w \in [\tau_n, 1-\tau_n]} \left| \log \frac{Q_j(w^{-1}, v)}{Q_j(w^{-1}, u)} \right| = o(\Delta_{\min}).$$

The theorems use only this one C_0 ; under (7) any $C_0 > 1$ suffices.

(E1) asks for continuity only; strict monotonicity of the conditional distribution function is not needed, because flat pieces are visited with probability zero. (E2) is the analogue of condition (H.4) in Gardes and Podgorny [9], normalized by $\log t$ and required only beyond the working tail level $(3\alpha)^{-1}$; it accommodates slowly varying factors such as $\ell(s) \asymp (\log s)^{-2}$, for which the left side is of order $1/\log(1/\alpha) \asymp 1/\log n$ when $\alpha = n^{-a}$, so gaps $\Delta_{\min} \gg 1/\log n$ are covered and smaller gaps are not. (E3) adapts their condition (H.5): the truncation $\tau_n = (pn)^{-2}$ makes the probability that any of the pn conditional PIT variables leaves $[\tau_n, 1 - \tau_n]$ vanish, which is the high-dimensional replacement for their fixed-dimension truncation. Their density conditions (H.1)–(H.2) have no counterpart here: empirical-rank windows have deterministic counts, so no lower bound on a covariate density is required.

These conditions are stated on the projected quantiles, not derived from (C1)–(C2): projection can generate genuinely new slow variation. If $W \sim U(0, 1)$ and $\mathbb{P}(Y > y \mid W) = y^{-1/(1+W)}$ exactly, then $\mathbb{P}(Y > y) \sim 4y^{-1/2}/\log y$, so even exact Pareto conditionals produce a logarithmic factor after projection, and the left side of (E2) is then of exact order $1/\log(1/\alpha)$. Exhibiting primitive model classes whose coordinatewise projections satisfy (E1)–(E3) with useful rates is left open, together with the derivation of the conditions themselves from a fully primitive model.

3.3 Main results

Theorem 3.1 (Score concentration). *Assume (C1)–(C2), (S), (E1)–(E3), $\alpha \leq 1/3$, $h \leq 1 - \varepsilon$, and*

$$\log(pn) = o(nh^2), \quad \log(pn) = o(n\alpha h). \quad (7)$$

Then there are deterministic sequences $a_n = O(\sqrt{\log(pn)/(n\alpha h)})$ and $b_n = o(\Delta_{\min})$ with

$$\mathbb{P}\left(\max_{j \leq p} |\hat{\Psi}_j - \Psi_j| \leq a_n + b_n\right) \rightarrow 1;$$

in shorthand, $\max_{j \leq p} |\hat{\Psi}_j - \Psi_j| = O_{\mathbb{P}}(\sqrt{\log(pn)/(n\alpha h)}) + o(\Delta_{\min})$.

The stochastic term is the exponential-spacings fluctuation of $O(n)$ local Hill statistics per coordinate, each built from at least $c n \alpha h$ spacings, controlled simultaneously over p coordinates; $\sqrt{\log(pn)/(n\alpha h)}$ is the resulting effective-tail-sample-size rate. The $o(\Delta_{\min})$ term collects the tail bias (E2), the spatial bias (E3) evaluated at $C_0 h$ — the first condition in (7) makes the rank displacement smaller than $(C_0 - 1)h$, so windows never leave the range covered by (E3) — and the quadrature error $\omega_\xi(1/n) + O(1/n)$ from (E1).

Theorem 3.2 (Sure screening). *Assume the conditions of Theorem 3.1 (including $\Delta_{\min} > 0$, i.e. every active coordinate detectable in the sense of Proposition 2.2) and*

$$\log(pn) = o(n\alpha h \Delta_{\min}^2). \quad (8)$$

Then, for every $d \geq s$,

$$\mathbb{P}(A \subseteq \hat{A}_d) \rightarrow 1,$$

and with probability tending to one every active score is smaller than every inactive score.

Corollary 3.3 (Exact recovery). *Under the conditions of Theorem 3.2, if $d = s$ then $\mathbb{P}(\hat{A}_s = A) \rightarrow 1$.*

The proof passes through a uniform bound on the whole profile $\max_j \sup_{u \in I_\varepsilon} |\hat{\xi}_j(u) - \xi_j(u)|$, stated with explicit finite-sample constants as Theorem A.12 in the appendix; Theorems 3.1 and 3.2 are its asymptotic readings. The argument has four ingredients. A single Dvoretzky–Kiefer–Wolfowitz event per coordinate controls all rank displacements at once, contributing the $\log(pn) = o(nh^2)$ requirement. Rank windows have deterministic sizes, at least $\lfloor \{h + \min(h, \varepsilon)\}(n+1) \rfloor - 1$ observations each, and, as u moves, pass through at most $4n+1$ membership states per coordinate. Conditionally on a covariate column, the probability-integral transforms of the responses are i.i.d. uniform, and the Rényi representation turns each local Hill statistic into a mean of i.i.d. exponentials plus a slowly varying remainder bounded through (E2). A union bound over the $p(4n+1)$ states then gives the $\log(pn)$ factor; no independence across overlapping windows or dependent coordinates is used. Finally, score separation is deterministic: when every score error is below $\Delta_{\min}/2$, every active score lies below $\gamma^* - \Delta_{\min}/2$ and every inactive score above it. Gardes and Podgornyy [9, Theorem 3] establishes the corresponding uniform consistency in fixed dimension over a compact class of projection matrices; the present setting replaces their covering-number argument and density assumptions by the rank construction above, and their result does not imply the growing- p bound needed here.

Condition (8) quantifies the cost of dimension: screening p coordinates costs $\log p$ against the effective local extreme count $n\alpha h$, deflated by the squared gap. With $\alpha = n^{-a}$ and $h = n^{-b}/2$ and $b \leq a$, dimensions up to $\log p = o(n^{1-a-b} \Delta_{\min}^2)$ are admissible, a polynomial range of p for fixed gaps; for $b > a$ the rank-displacement condition $\log(pn) = o(nh^2) = o(n^{1-2b})$ binds first. These ranges treat (E3) as given; a spatial modulus of order $h \log(1/\tau_n)$, the natural benchmark, adds the constraint $\log(pn) = o(n^b)$.

4 Simulation study

Each coordinate is scored in $O(n \log n)$ time: one sort by empirical rank, then a sliding window over the rank grid whose local Hill statistics are maintained by a Fenwick tree indexed by global response ranks. Covariates are generated and processed in blocks, so the total cost is $O(pn \log n)$ time and $O(n)$ working memory beyond the scores. At $(n, p) = (2000, 1000)$ one complete screen takes under a second on a laptop core. Scores are sorted increasingly, ties broken by coordinate index; a coordinate whose grid is entirely underpopulated receives score $+\infty$.

4.1 Models

All four models share the covariate design and the tail-index active set $A_\gamma = \{1, 2, 3, 4\}$. Latent Gaussian AR(1) covariates $Z_{ij} = \rho Z_{i,j-1} + \sqrt{1 - \rho^2} \epsilon_{ij}$ are transformed to $U_{ij} = \Phi(Z_{ij})$; the tail-index functions are evaluated at the population transforms U , while the estimator sees only Z and

computes its own empirical ranks. Responses are generated by quantile inversion,

$$Y = V^{-\gamma(U)} \ell(1/V, \cdot), \quad V \sim \text{Unif}(0, 1),$$

the mechanism used by Gardes and Podgorny [9]. The tail-index functions are

$$\begin{aligned} \gamma_1(u) &= 0.5 \exp(-\sum_{j=1}^4 u_j), \\ \gamma_2(u) &= 0.5 \exp(-0.5 \sum_{j=1}^4 u_j), \\ \gamma_3(u) &= 0.55 \exp(-0.80 \sum_{j=1}^4 u_j - 0.20 u_1 u_2 - 0.15 u_3 u_4). \end{aligned}$$

Models M1, M3 and M4 carry the slowly varying factor of Gardes and Podgorny [9, Section 4.3],

$$\ell^{(1)}(s, u) = [1 + \exp\{v(u) - s\}]^{-1}, \quad v(u) = 2.5(u_5 + u_6 + u_7 + u_8) \in (0, 10),$$

matching the range of their $B_1^\top x$; since $\ell^{(1)} \rightarrow 1$ as $s \rightarrow \infty$, its effect vanishes in the tail.

Model M2 instead carries a scale factor that persists at every finite quantile. A single latent standard normal factor F , independent of the active coordinates and not itself observed, enters the response through

$$\log \ell^{(2)}(s, F) = -\frac{1}{2s} + \kappa_c \left\{ \Phi(F) - \frac{1}{2} \right\}, \quad \kappa_c = 0.4772,$$

so that the scale part of $\ell^{(2)}$ is strictly positive, bounded within $[e^{-\kappa_c/2}, e^{\kappa_c/2}]$ — compatible with the uniform constants in (C2) — and has $\text{Var}[\kappa_c \{\Phi(F) - \frac{1}{2}\}] = \kappa_c^2/12$ exactly. The analyst observes $s = 20$ proxies of F , occupying the coordinates $A_{\text{scale}} = \{5, \dots, 24\}$,

$$Z_{ij} = \lambda F_i + \sqrt{1 - \lambda^2} \epsilon_{ij}, \quad j \in A_{\text{scale}}, \quad \lambda = 0.7,$$

each marginally standard normal like every other covariate and correlated with the nuisance at level λ (pairwise intra-cluster correlation $\lambda^2 = 0.49$). The four models are

$$\text{M1} = (\gamma_1, \ell^{(1)}), \quad \text{M2} = (\gamma_1, \ell^{(2)}), \quad \text{M3} = (\gamma_2, \ell^{(1)}), \quad \text{M4} = (\gamma_3, \ell^{(1)}).$$

Each model isolates one difficulty. M1 is the monotone reference. M2 keeps the tail index of M1 and adds the persistent scale cluster: its tail index depends on A_γ only, while every finite conditional quantile also depends on the twenty scale coordinates, which are individually attractive to any screen built on finite quantiles yet carry no information about the limiting exponent. M3 halves the exponent range of the signal, weakening the trimmed population gaps from 0.186 (M1, M2) to 0.107. M4 combines a strong monotone signal (trimmed gap 0.173) with interactions among the active coordinates, so it probes departures from additivity rather than weakness of signal. All four models are implemented in **generate4.R**, with $(\kappa_c, s, \lambda) = (0.4772, 20, 0.7)$ fixed throughout.

None of the four models satisfies the projected-quantile conditions (E1)–(E3) with gaps of the order required at $n = 2000$: their tail-index surfaces are maximized at a corner of the active cube with no atom in the fibre-maximizer law, which places the projected quantiles in the logarithmic slow-variation regime discussed after (E3) in Section 3, where the bias modulus at $n = 2000$ is of order one, larger than every population gap. The simulations therefore probe the method beyond the regime its finite-sample theory certifies. Ranking nevertheless succeeds because much of the local Hill bias is shared across coordinates — every inactive coordinate estimates the same projected mixture — and the screen uses only score differences; the experiments measure this common-mode behaviour, which the theory does not exploit.

The common Monte Carlo design is

$$n = 2000, \quad p \in \{500, 1000, 2000\}, \quad \rho = 0.25, \quad d = 20, \quad \varepsilon = 0.05,$$

with 1,000 replications per cell and per model in the method comparison (binomial standard errors at most 0.016) and 40 replications per cell in the tuning study (standard errors at most 0.079); Sure- d denotes $\mathbb{P}(R_{\max} \leq d)$ with $R_{\max} = \max_{j \in A_\gamma} \text{rank}(j)$, and all methods within a replication are evaluated on the same dataset.

4.2 Tuning and the aggregated screen

The two smoothing parameters are parameterized as

$$\alpha = n^{-a}, \quad h = n^{-b}/2,$$

following the convention of Gardes and Podgorny [9]. We evaluate Sure-20 for each of the four models over the crossed grid $a \in \{.25, .30, .35, .40, .45, .50\}$, $b \in \{0, .05, .10, .15, .20, .30, .40\}$ at $n = 2000$, $p = 1000$, $\rho = 0.25$; Figure 2 shows the four heatmaps.

Three features stand out. The row $a = 0.25$ collapses for the three $\ell^{(1)}$ models (Sure-20 at most .095, .050 and .200 for M1, M3 and M4 over $b \leq .20$) but not for M2 (up to .475): at $\alpha = n^{-0.25} \approx 0.15$ the local Hill statistics reach into quantile regions where $\ell^{(1)}$, whose argument $v(u) - s$ is dominated by the U_5 – U_8 combination at moderate s , deforms the projected distributions, while the bounded scale factor of M2 produces no such deformation. Performance degrades toward large a and large b together, where the local extreme count collapses, and the degenerate bandwidth $b = 0$ is never competitive. In between, the row $a = 0.30$ is best for M1, M3 and M4 simultaneously, and

$$(a^*, b^*) = (0.30, 0.15)$$

attains the highest four-model mean Sure-20 on the grid (0.796, against 0.785 for (0.30, 0.10)), with per-model values .965, .710, .585 and .925. We fix (a^*, b^*) for every subsequent experiment, in simulation and in the application; at $n = 2000$ this gives $\alpha \approx 0.10$, $h \approx 0.16$, and about $2n\alpha h \approx 64$ effective local extremes per interior window. No model is given its own tuning.

The aggregated screen. A single pair (a, b) makes the final ranking depend entirely on one tuning. To stabilize the ranking against small variations of the tuning, we also evaluate an aggregated version of the screen, built on the 3×3 block of settings

$$\mathcal{N}_9 = \{0.30, 0.35, 0.40\} \times \{0.10, 0.15, 0.20\},$$

the neighbourhood of (a^*, b^*) within the well-behaved region of the grid (the row $a = 0.25$, which collapses for three of the four models, is excluded). For each setting $(a, b) \in \mathcal{N}_9$ and each coordinate j , compute the score $\widehat{\Psi}_j(a, b)$ and its rank $r_j(a, b)$ among the p coordinates, ranks increasing with the score; aggregate by

$$r_j^{\text{agg}} = \min_{(a, b) \in \mathcal{N}_9} r_j(a, b),$$

and rank the coordinates by increasing r_j^{agg} , ties broken by coordinate index. A coordinate is thus retained early if at least one setting of the block ranks it early. We make no optimality claim for this construction; it is the procedure used in the experiments below and in the application, and its effect is measured against the single-setting screen in Section 4.3. Its cost is nine score passes instead of one, leaving the $O(pn \log n)$ complexity unchanged up to the constant.

The two versions are reported side by side throughout: *Tail-index SIS, selected* (a^*, b^*) uses the central pair alone; *Tail-index SIS, aggregated* uses r_j^{agg} .




Figure 2: Sure-20 over the (a, b) tuning grid for the four models at $n = 2000$, $p = 1000$, $\rho = 0.25$ (40 replications per cell; $\alpha = n^{-a}$, $h = n^{-b}/2$; standard errors at most 0.079). The dashed rectangle marks the 3×3 aggregation block $\mathcal{N}_9$ of Section 4.2.

4.3 Comparison with competing screens

Six rules are compared on the same 1,000 simulated datasets per model and per dimension $p \in \{500, 1000, 2000\}$, at $n = 2000$, $\rho = 0.25$: the two versions of the tail-index screen; the conditional-Pickands screen of Yoshida and Umezu [19], implemented from their equations (2)–(4) with the Epanechnikov kernel and the tuning of their numerical work transplanted proportionally ($h = 1$, $k = \lfloor 0.072n \rfloor$), on 25 equally spaced evaluation points in $[.02, .98]$; and the quantile-adaptive SIS of He et al. [11] at the levels $\tau \in \{.90, .95, .99\}$, using cubic B-spline marginal quantile regression with three degrees of freedom, identical at every level. The competitors rank large utilities first; ours ranks small scores first. Tables 1–3 report, for each model and method, Sure-4 (exact recovery), Sure-20, and the mean and median of the worst active rank.

Sensitivity to the dimension. Read across the three tables, the dimension degrades every method, but at very different rates, and the mean–median contrast identifies the mechanism. For the single-setting tail screen on M1, $\mathbb{E}(R_{\max})$ grows from 5.7 to 12.9 as p goes from 500 to 2000 while $\text{Med}(R_{\max})$ stays at 4: the typical replication is unaffected by the extra competitors, and the degradation is carried by a minority of replications in which an active coordinate falls into the null

Table 1: Method comparison at $n = 2000$, $p = 500$, $\rho = 0.25$, on the same 1,000 replications per model. Sure- d is the probability that all four tail-index-active variables rank among the d smallest scores ($d = 4$ is exact recovery); $\mathbb{E}(R_{\max})$ and $\text{Med}(R_{\max})$ are the mean and median of the worst active rank $R_{\max} = \max_{j \in A} \text{rank}(j)$. Standard errors of the probabilities are at most 0.016.

Model	Method	Sure-4	Sure-20	$\mathbb{E}(R_{\max})$	$\text{Med}(R_{\max})$
M1	Tail-index SIS, selected (a^*, b^*)	0.849	0.979	5.7	4
	Tail-index SIS, aggregated	0.922	0.990	4.6	4
	Yoshida–Umezū	0.202	0.467	97.4	29
	Quantile SIS ($\tau = .90$)	0.299	0.894	11.7	6
	Quantile SIS ($\tau = .95$)	0.891	0.998	4.2	4
	Quantile SIS ($\tau = .99$)	0.125	0.852	13.0	9
M2	Tail-index SIS, selected (a^*, b^*)	0.331	0.757	26.9	6
	Tail-index SIS, aggregated	0.549	0.921	8.8	4
	Yoshida–Umezū	0.008	0.111	128.3	102
	Quantile SIS ($\tau = .90$)	0.000	0.039	23.8	24
	Quantile SIS ($\tau = .95$)	0.060	0.631	16.2	17
	Quantile SIS ($\tau = .99$)	0.041	0.688	20.5	14
M3	Tail-index SIS, selected (a^*, b^*)	0.256	0.655	38.2	9
	Tail-index SIS, aggregated	0.245	0.813	19.5	7
	Yoshida–Umezū	0.028	0.178	153.9	124
	Quantile SIS ($\tau = .90$)	0.099	0.597	33.8	14
	Quantile SIS ($\tau = .95$)	0.431	0.930	8.2	5
	Quantile SIS ($\tau = .99$)	0.005	0.268	58.5	37
M4	Tail-index SIS, selected (a^*, b^*)	0.793	0.967	7.1	4
	Tail-index SIS, aggregated	0.847	0.990	4.9	4
	Yoshida–Umezū	0.211	0.576	67.9	12
	Quantile SIS ($\tau = .90$)	0.397	0.932	9.3	5
	Quantile SIS ($\tau = .95$)	0.867	0.997	4.3	4
	Quantile SIS ($\tau = .99$)	0.065	0.758	17.6	12

range, where its rank scales with p . Aggregation compresses exactly that tail ($\mathbb{E}(R_{\max})$ $4.6 \rightarrow 6.1$ on M1, $8.8 \rightarrow 19.5$ against $26.9 \rightarrow 98.8$ on M2), and the quantile screen at $\tau = .95$ is the least dimension-sensitive on the $\ell^{(1)}$ models ($4.2 \rightarrow 4.8$ on M1). M3 is the most dimension-sensitive model for every method (Sure-20 of the aggregated screen $.813 \rightarrow .580$, of quantile $.95$ $.930 \rightarrow .782$): a weak signal loses the most from additional competition. Yoshida–Umezū degrades from a low base on all four models.

Sensitivity to the quantile level. Within every table, the quantile screen is non-monotone in τ , and in opposite directions on either side of $\tau = .95$. At $\tau = .90$ it fails on M2 in the most instructive way: its top four contain on average 3.7 scale variables, and all twenty scale coordinates fill its top 24, so exact recovery is essentially zero — correct behaviour for its own target, since those variables genuinely move the $.90$ quantile, and fatal for tail-index recovery. Raising τ to $.99$ almost eliminates the scale contamination (0.18–0.41 scale variables in the top four) because the quantile target approaches the tail index; the variance cost, however, is severe on every model

Table 2: Method comparison at $n = 2000$, $p = 1000$, $\rho = 0.25$, on the same 1,000 replications per model. Sure- d is the probability that all four tail-index-active variables rank among the d smallest scores ($d = 4$ is exact recovery); $\mathbb{E}(R_{\max})$ and $\text{Med}(R_{\max})$ are the mean and median of the worst active rank $R_{\max} = \max_{j \in A} \text{rank}(j)$. Standard errors of the probabilities are at most 0.016.

Model	Method	Sure-4	Sure-20	$\mathbb{E}(R_{\max})$	$\text{Med}(R_{\max})$
M1	Tail-index SIS, selected (a^*, b^*)	0.809	0.954	8.3	4
	Tail-index SIS, aggregated	0.882	0.985	4.9	4
	Yoshida–Umezu	0.179	0.398	188.1	58
	Quantile SIS ($\tau = .90$)	0.219	0.804	18.6	7
	Quantile SIS ($\tau = .95$)	0.830	0.997	4.6	4
	Quantile SIS ($\tau = .99$)	0.068	0.674	22.2	15
M2	Tail-index SIS, selected (a^*, b^*)	0.290	0.683	54.4	8
	Tail-index SIS, aggregated	0.466	0.866	15.8	5
	Yoshida–Umezu	0.003	0.066	258.4	202
	Quantile SIS ($\tau = .90$)	0.000	0.032	25.2	24
	Quantile SIS ($\tau = .95$)	0.054	0.593	17.0	18
	Quantile SIS ($\tau = .99$)	0.026	0.463	34.1	22
M3	Tail-index SIS, selected (a^*, b^*)	0.192	0.525	81.0	17
	Tail-index SIS, aggregated	0.183	0.695	38.7	9
	Yoshida–Umezu	0.021	0.138	320.1	256
	Quantile SIS ($\tau = .90$)	0.059	0.456	68.7	24
	Quantile SIS ($\tau = .95$)	0.303	0.873	14.8	6
	Quantile SIS ($\tau = .99$)	0.004	0.103	120.5	74
M4	Tail-index SIS, selected (a^*, b^*)	0.747	0.939	10.5	4
	Tail-index SIS, aggregated	0.766	0.980	6.4	4
	Yoshida–Umezu	0.202	0.507	123.9	20
	Quantile SIS ($\tau = .90$)	0.306	0.885	15.7	6
	Quantile SIS ($\tau = .95$)	0.799	0.994	4.7	4
	Quantile SIS ($\tau = .99$)	0.030	0.533	29.8	19

(Sure-4 at most .125 across models at $p = 500$ and lower at $p = 2000$). The level $\tau = .95$ is the best compromise on M1, M3 and M4, and on none of the three levels does the quantile screen approach the tail screen on M2 (Sure-4 at most .073 against .431 at $p = 2000$; median worst rank 17–18 against 4–5, the signature of one to two scale variables permanently occupying its leading positions). The direction of the τ -sensitivity does not depend on p ; its amplitude grows with p on the weak-signal M3. Yoshida–Umezu, which estimates the index through Pickands-type quantile differences at its own fixed levels, is dominated on all models and dimensions here, with 6.4–9.7 of the twenty M2 scale coordinates in its top 24.

Effect of the aggregation. The aggregated screen improves on the selected pair almost uniformly, and most where failures are worst. On M2 it lifts exact recovery from .331/.290/.263 to .549/.466/.431 across the three dimensions and divides $\mathbb{E}(R_{\max})$ by three to five; on M1 and M4 it adds two to seven points of Sure-4 and compresses the rank tail (M4 at $p = 2000$: $\mathbb{E}(R_{\max})$ 21.6 $\rightarrow$ 8.0); Sure-20 improves in all twelve model–dimension cells. The one systematic exception is exact recovery on

Table 3: Method comparison at $n = 2000$, $p = 2000$, $\rho = 0.25$, on the same 1,000 replications per model. Sure- d is the probability that all four tail-index-active variables rank among the d smallest scores ($d = 4$ is exact recovery); $\mathbb{E}(R_{\max})$ and $\text{Med}(R_{\max})$ are the mean and median of the worst active rank $R_{\max} = \max_{j \in A} \text{rank}(j)$. Standard errors of the probabilities are at most 0.016.

Model	Method	Sure-4	Sure-20	$\mathbb{E}(R_{\max})$	$\text{Med}(R_{\max})$
M1	Tail-index SIS, selected (a^*, b^*)	0.751	0.941	12.9	4
	Tail-index SIS, aggregated	0.851	0.983	6.1	4
	Yoshida–Umezū	0.125	0.305	432.2	168
	Quantile SIS ($\tau = .90$)	0.161	0.665	36.6	11
	Quantile SIS ($\tau = .95$)	0.758	0.992	4.8	4
	Quantile SIS ($\tau = .99$)	0.025	0.406	40.2	25
M2	Tail-index SIS, selected (a^*, b^*)	0.263	0.639	98.8	9
	Tail-index SIS, aggregated	0.431	0.832	19.5	5
	Yoshida–Umezū	0.009	0.059	480.7	368
	Quantile SIS ($\tau = .90$)	0.000	0.031	24.6	24
	Quantile SIS ($\tau = .95$)	0.073	0.604	17.1	18
	Quantile SIS ($\tau = .99$)	0.009	0.257	59.7	39
M3	Tail-index SIS, selected (a^*, b^*)	0.149	0.463	144.8	27
	Tail-index SIS, aggregated	0.113	0.580	76.0	14
	Yoshida–Umezū	0.015	0.084	640.2	545
	Quantile SIS ($\tau = .90$)	0.039	0.328	122.2	42
	Quantile SIS ($\tau = .95$)	0.234	0.782	24.7	7
	Quantile SIS ($\tau = .99$)	0.000	0.022	245.7	154
M4	Tail-index SIS, selected (a^*, b^*)	0.663	0.898	21.6	4
	Tail-index SIS, aggregated	0.728	0.965	8.0	4
	Yoshida–Umezū	0.151	0.430	282.3	44
	Quantile SIS ($\tau = .90$)	0.236	0.767	29.8	7
	Quantile SIS ($\tau = .95$)	0.694	0.986	5.4	4
	Quantile SIS ($\tau = .99$)	0.019	0.295	58.9	36

the weak-signal M3, where aggregation trades Sure-4 away ($.256 \rightarrow .245$ at $p = 500$, $.149 \rightarrow .113$ at $p = 2000$) while still improving Sure-20 ($.463 \rightarrow .580$ at $p = 2000$) and $\mathbb{E}(R_{\max})$ ($144.8 \rightarrow 76.0$): the minimum promotes any coordinate that some setting ranks early, which occasionally lets a lucky null into the top four when the active signal is weak, yet reliably rescues active coordinates from single-setting failures further down the list. Aggregation is thus a stabilizer of inclusion rather than of exact selection, which matches the intended first-stage use of a screen.

Overall, the quantile screen at a moderate level remains the strongest detector when high quantiles and the limiting exponent share their covariates and the signal is weak (M3); the aggregated tail screen is competitive or better on M1 and M4 and is the only method that recovers the tail-index-active set reliably in the presence of the M2 scale cluster, which distracts every fixed-quantile method at every level. For a question about the limiting exponent itself, the two rankings should be read side by side, as in Section 5: variables high on one list and low on the other are precisely where the two targets part ways.

5 Application: violent crime across U.S. communities

The analysis contains $n = 1,993$ communities and $p = 99$ candidate predictors. Under the tuning selected in Section 4.2, $\alpha = n^{-0.3} \approx 0.102$ and $h = n^{-0.15}/2 \approx 0.160$, so the sample contains $n\alpha \approx 204$ upper-tail observations in total and about $2n\alpha h \approx 65$ within a typical interior rank window. The dimension is therefore comparable to the number of observations available for local tail estimation. Unrestricted or nonparametric joint tail modelling with 99 covariates is poorly supported by roughly 200 upper-tail observations, and by fewer still within any local covariate neighbourhood; a marginal screen can rank the 99 candidates from exactly this information.

5.1 Data and protocol

The Communities and Crime data set [15], distributed by the UCI Machine Learning Repository in its unnormalized version, links socio-economic census attributes, law-enforcement survey attributes and FBI crime counts for 2,215 U.S. communities. The response is the violent-crime rate per 100,000 inhabitants. We keep the communities with an observed positive response ($n = 1,993$). Preprocessing of the 124 documented predictive attributes is response-blind: we retain numeric predictors with fewer than 5% missing values, which removes the 22 law-enforcement variables reported by only 343 communities, and with at least 100 distinct observed values, which removes 3 low-cardinality variables, keeping the application close to the continuous-marginal framework of the theory. This leaves $p = 99$ continuous predictors — demographic composition, income and poverty, education, employment, family structure, immigration, language, housing, and population density. The at most one missing value per retained predictor is median-imputed, and every method below works from the same imputed matrix.

The response is strongly right-skewed: median 374, 99th percentile 2,943, maximum 4,877. The left panel of Figure 3 shows the Hill curve of the response over $k \in [20, 600]$ upper order statistics; it is positive and moves slowly, between 0.283 and 0.374 for $k \in [100, 300]$, with value 0.328 at the working level $k = n\alpha = 204$. The middle panel shows the Pareto QQ plot of the 600 largest responses, approximately linear over the working range with least-squares slope 0.27. The gap between the QQ slope and the Hill value at the same level is mild evidence that regular variation is only approximate over the working range. The response exhibits a sufficiently heavy and stable upper tail to make a tail-index analysis plausible; neither diagnostic establishes regular variation beyond doubt, and no single threshold carries a structural interpretation.

The screening protocol is the one fixed in Section 4.2, applied unchanged: the aggregated tail-index screen computes the score at each of the nine settings $\mathcal{N}_9 = \{0.30, 0.35, 0.40\} \times \{0.10, 0.15, 0.20\}$, converts each pass into ranks, and orders the predictors by the minimum rank r_j^{agg} across the nine settings; the selected-pair screen at $(a^*, b^*) = (0.30, 0.15)$ is reported alongside. The two competing screens of Section 4.3 are run on the same imputed matrix: Yoshida–Umezū at its transplanted tuning and the quantile screen at $\tau \in \{.90, .95, .99\}$.

5.2 Results

Poverty and family structure dominate the aggregated ranking, and they dominate every competing ranking as well. The poverty rate (`pctPoverty`) is first, with per-setting ranks between 1 and 6 across the nine settings; the two-parent household variables `pct2Par` and `pctKids2Par` are third and fourth (ranks 1–5), and both are also leading variables for the quantile screen at every level (`pctKids2Par` is first at all three τ) and for Yoshida–Umezū (ranks 3–7). The related `pctKids4w2Par` and `pct12to17w2Par` complete the consensus block at aggregated ranks 7–8. For these variables the body and the tail of the crime distribution point at the same covariates, and every method agrees.




Figure 3: Communities and Crime application ($n = 1,993$, $p = 99$, $\alpha = n^{-0.3}$, $h = n^{-0.15}/2$). Left: Hill estimates for the violent-crime rate; the dashed line marks the working level $k = n\alpha = 204$. Middle: Pareto QQ plot of the 600 largest responses; the dashed line is fitted over the working range $i \leq 204$. Right: for the ten leading predictors of the aggregated screen, the nine per-setting ranks (open circles) and the aggregated rank (filled circles); the dotted line is the default retained size $d = 20$.

The informative disagreements are the tail-specific candidates. The share of workers using public transportation (`pctUsePubTrans`, aggregated rank 5) and the farm-employment rate (`pctWfarm`, rank 6) rank far lower under every fixed-quantile method: ranks 49–73 for the quantile screen across its three levels and 57–60 for Yoshida–Umezu. Their low tail-index scores are consistent with marginal variation in upper-tail heaviness across their values, of a kind that fixed-quantile screens are not designed to detect; Section 4.3 produces the same dissociation under controlled conditions, where the target is known. The two variables differ in stability across the aggregation block: `pctUsePubTrans` ranks within the top 15 at every one of the nine settings, while `pctWfarm` moves between 2 and 47 (right panel of Figure 3). The never-married male rate (`pctMaleNevMar`, aggregated rank 2, per-setting ranks 1–9, quantile ranks 28–35) sits between the consensus block and the tail-specific pair. We report all three as candidate tail-specific predictors for a second-stage analysis.

The minimum aggregation also exhibits its known failure mode, and the per-setting ranks make it visible: `pctLargHous` (aggregated rank 9) and `pctHousWOplumb` (rank 10) owe their positions to a single favourable setting each (per-setting ranks 6–74 and 6–55; selected-pair ranks 46 and 36). With 95 competitors, a coordinate promoted by one setting out of nine warrants no interpretation, and we draw none. The consensus block and the tail-specific candidates, by contrast, hold their positions across the whole block.

We do not interpret any ranking causally: the screen identifies covariates along which the conditional tail index varies marginally, and confounding among the 99 predictors is certain.

6 Discussion

The screen is a first stage. Its intended pipeline reduces p coordinates to a manageable d , after which a joint method — the central tail-index subspace of Gardes and Podgorny [9], a structured

tail regression [17, 18], or multivariate local estimation — can resolve interactions and correlated neighbours among the retained variables. The two stages solve different problems: rapid elimination of evidently irrelevant coordinates scales to $p \gg n$, while accurate joint tail inference does not.

Three limitations are intrinsic. First, activity without marginal detectability: when every fibre of an active coordinate meets the global argmax set, the score carries no information about it, at any sample size. Proposition 2.2 characterizes this failure exactly, and it is a property of the target, so enlarging d or n cannot repair it; iterative or grouped extensions would require new identification arguments. Second, mixture bias: when the projected slowly varying factor decays logarithmically, condition (E2) restricts the detectable gaps to $\Delta_{\min} \gg 1/\log n$, and the continuous-mixture example of Section 3 shows the restriction is intrinsic rather than an artifact of the proofs. Third, rare favourable configurations: even with a comfortable population gap, the envelope near its maximum is estimable only from observations whose remaining active coordinates are simultaneously near their maximizing values, and the mass of such configurations can decay exponentially with the number of active coordinates. The weak-signal model M3 in Section 4 shows the resulting cost.

The main open theoretical question is the gap between the primitive model (C1)–(C2) and the estimation conditions (E1)–(E3). Projection creates slow variation that the primitive model does not control, and we do not know useful primitive conditions implying uniform triangular-array moduli for all coordinatewise projections. The score functional also invites variants: a minimum or a low quantile of the profile would emphasize depressions confined to short intervals, at the price of new uniformity arguments for empirical profile extrema.

In practice, tuning matters more than the theory prescribes. The tail fraction trades Hill bias against variance; the bandwidth additionally determines how often near-maximizing configurations of the other active coordinates enter a window. We recommend the workflow of Sections 4 and 5: run the screen on the block of nine neighbouring tunings, rank by the aggregated minimum rank, read the per-setting ranks as the stability diagnostic — a variable promoted by a single setting warrants no interpretation, one holding its rank across the block does — retain a generous d , and reserve interpretation for variables that survive. Dependence deserves the same caution: it can help, by concentrating active configurations, or hurt, by promoting inactive neighbours, and the direction is design-specific.

Projection turns a tail-index surface into an upper envelope. Variation of that envelope is a scalable marginal signal for tail-index activity, and its absence, characterized by the argmax projections, tells the analyst exactly what a marginal screen can and cannot see.

A Proofs

Throughout the appendix the dimension is written $p = p_n$, and the auxiliary moduli suppressed in the main text reappear:

$$b_n(a) = \max_{j \leq p} \sup_{u \in I_\varepsilon} \sup_{s \geq a^{-1}, t > 1} \frac{|\log\{\ell_j(ts, u)/\ell_j(s, u)\}|}{\log t}, \quad \omega_n(r, \tau) = \max_{j \leq p} \sup_{u \in I_\varepsilon} \sup_{\substack{v \in [0, 1] \\ |v - u| \leq r}} \sup_{w \in [\tau, 1 - \tau]} \left| \log \frac{Q_j(w^{-1}, v)}{Q_j(w^{-1}, u)} \right|,$$

so that (E2) reads $b_n(3\alpha) = o(\Delta_{\min})$ and (E3) reads $\omega_n(C_0 h, \tau_n) = o(\Delta_{\min})$ with $\tau_n = (pn)^{-2}$; note that $\omega_n(\cdot, \tau)$ is nondecreasing in its first argument, so (E3) bounds $\omega_n(r, \tau_n)$ for every $r \leq C_0 h$. Put $\delta_n = (n + 1)^{-1}$,

$$m_n^- = \lfloor \{h + \min(h, \varepsilon)\}(n + 1) \rfloor - 1, \quad K_n = \lfloor \alpha m_n^- \rfloor, \quad S_n = 4n + 1.$$

Once $nh \geq 4$ and $\alpha m_n^- \geq 2$, one has $m_n^- \geq nh/2$ and $K_n \geq \alpha m_n^-/2 \geq nah/4$; the rate conditions (7) force both requirements for n large, which is the only regime where the conversion $K_n \asymp nah$ is

used. All conditional distributions are the pointwise compatible versions of Section 2, and constants are uniform over the triangular array. The appendix is organized as follows: preliminary lemmas (A.1); projection (A.2); population results (A.3); the finite-sample uniform profile bound (A.4); quadrature, score concentration and screening, including the proofs of Theorems 3.1 and 3.2 (A.5).

A.1 Preliminary lemmas

The first lemma transfers pointwise perturbations to order statistics, as in Gardes and Podgorny [9, Appendix, Lemma 3].

Lemma A.1 (Sorting is sup-norm nonexpansive). *Let $a_1, \dots, a_m$ and $b_1, \dots, b_m$ satisfy $a_i \leq b_i$ for all i . Then $a_{(i)} \leq b_{(i)}$ for all i . Consequently $|a_i - b_i| \leq \epsilon$ for all i implies $|a_{(i)} - b_{(i)}| \leq \epsilon$ for all i .*

Proof. Fix i . At least $m - i + 1$ indices r satisfy $a_r \geq a_{(i)}$, and for each such r , $b_r \geq a_r \geq a_{(i)}$; hence $b_{(i)} \geq a_{(i)}$. For the second claim apply the first to $(a_i - \epsilon, b_i)$ and $(b_i - \epsilon, a_i)$. $\square$

Lemma A.2 (Chernoff bound for exponential means). *Let L_k be the mean of k i.i.d. standard exponentials. For $0 < t \leq 1$, $\mathbb{P}(|L_k - 1| > t) \leq 2e^{-kt^2/4}$.*

Proof. Markov's inequality applied to $e^{\lambda k L_k}$, optimized at $\lambda = t/(1+t)$, gives $\mathbb{P}(L_k > 1+t) \leq e^{-k\{t - \log(1+t)\}}$. The function $\varphi(t) = t - \log(1+t) - t^2/4$ has $\varphi(0) = 0$ and

$$\varphi'(t) = 1 - \frac{1}{1+t} - \frac{t}{2} = \frac{t(1-t)}{2(1+t)} \geq 0 \quad \text{on } [0, 1],$$

so $t - \log(1+t) \geq t^2/4$ on $(0, 1]$. For the lower tail, $\mathbb{P}(L_k < 1-t) \leq e^{-k\{-t - \log(1-t)\}}$ and $-t - \log(1-t) \geq t^2/2 \geq t^2/4$. $\square$

Lemma A.3 (Threshold order statistic of uniforms). *Let $V_{(1)} < \dots < V_{(m)}$ be the order statistics of m i.i.d. standard uniforms, $0 < \alpha \leq 1/3$ and $k = \lfloor \alpha m \rfloor \geq 1$. Then $\mathbb{P}\{V_{(k+1)} > 3\alpha\} \leq e^{-(k+1)/4}$.*

Proof. If $3\alpha \geq 1$ the probability is zero. Otherwise $\{V_{(k+1)} > 3\alpha\} = \{N \leq k\}$ with $N \sim \text{Bin}(m, 3\alpha)$ and mean $\mu = 3\alpha m \geq 3k$. The multiplicative Chernoff bound $\mathbb{P}\{N \leq (1-\delta)\mu\} \leq e^{-\delta^2\mu/2}$ [3, Section 2.2] with $\delta = 2/3$ gives $\mathbb{P}(N \leq \mu/3) \leq e^{-2\mu/9} \leq e^{-2k/3} \leq e^{-(k+1)/4}$, the last step because $8k \geq 3(k+1)$ for $k \geq 1$. $\square$

A.2 Projection

Let $\mathcal{G} \subseteq \sigma(U)$, $T = \gamma(U)$, and $\xi = \text{ess sup}(T \mid \mathcal{G})$, the smallest $\mathcal{G}$ -measurable random variable dominating T almost surely.

Lemma A.4 (Uniform remainder replacement). *Define $M(y) = \mathbb{E}\{c(U)y^{-1/T} \mid \mathcal{G}\}$. Under (C1)–(C2), almost surely,*

$$\left| \frac{\mathbb{P}(Y > y \mid \mathcal{G})}{M(y)} - 1 \right| \leq \rho(y) := \sup_u \left| \frac{L(y, u)}{c(u)} - 1 \right| \rightarrow 0. \quad (\text{A.1})$$

Proof. By the tower property, $\mathbb{P}(Y > y \mid \mathcal{G}) = \mathbb{E}\{y^{-1/T} L(y, U) \mid \mathcal{G}\}$. Writing $L(y, U) = c(U)\{1 + r(y, U)\}$ with $|r(y, U)| \leq \rho(y)$,

$$|\mathbb{P}(Y > y \mid \mathcal{G}) - M(y)| = |\mathbb{E}\{c(U)y^{-1/T} r(y, U) \mid \mathcal{G}\}| \leq \rho(y)M(y),$$

and $\rho(y) \rightarrow 0$ is (C2). $\square$

The next lemma is the point at which the argument departs from the proof of Proposition 1 in Gardes and Podgorny [9]: their condition (C.1) assumes the law of $\gamma(X)$ absolutely continuous, which identifies $\{\gamma > t\}$ with $\{\gamma \geq t\}$ in their computation. The tilting argument below needs no such assumption and allows purely atomic conditional laws of T — the relevant case for inactive coordinates and constant-index null models.

Lemma A.5 (Tilted-measure concentration). *For $y > 1$ define the random probability measure*

$$K_y^*(A) = \frac{\mathbb{E}\{c(U)y^{-1/T}\mathbf{1}_{\{T \in A\}} \mid \mathcal{G}\}}{M(y)}, \quad A \in \mathcal{B}([\underline{\gamma}, \Gamma]).$$

On a common probability-one event, $K_y^ \Rightarrow \delta_\xi$ weakly as $y \rightarrow \infty$.*

Proof. Since $K_y^*\{T > \xi + \delta\} = 0$ for every $\delta > 0$ by definition of ξ , it suffices to prove, for fixed $\delta \in (0, \underline{\gamma})$,

$$K_y^*\{T \leq \xi - \delta\} \longrightarrow 0 \quad \text{almost surely.} \quad (\text{A.2})$$

Split by the $\mathcal{G}$ -measurable event $G_\delta = \{\xi \geq \underline{\gamma} + \delta\}$. On G_δ^c , $\xi - \delta < \underline{\gamma} \leq T$, so $\{T \leq \xi - \delta\}$ is empty and (A.2) is trivial. On G_δ , set $r = \xi - \delta$, $s = \xi - \delta/2$ and $\pi = \mathbb{P}(\overline{T} > s \mid \mathcal{G})$. Then $\pi > 0$ almost surely on G_δ : otherwise $\xi' = \xi - (\delta/2)\mathbf{1}_{G_\delta \cap \{\pi=0\}}$ would be a $\mathcal{G}$ -measurable almost-sure upper bound for T , strictly smaller than ξ with positive probability, contradicting minimality of the conditional essential supremum. Bounding numerator and denominator separately, using only $c_- \leq c \leq c_+$ and monotonicity of $g \mapsto y^{-1/g}$,

$$K_y^*\{T \leq r\} \leq \frac{c_+ y^{-1/r}}{c_- \pi y^{-1/s}} = \frac{c_+}{c_- \pi} y^{-(1/r - 1/s)} \longrightarrow 0.$$

A rational sequence $\delta \downarrow 0$ and a countable intersection give a common probability-one event. $\square$

Proof of Proposition 2.1. For fixed $t > 0$,

$$\frac{M(ty)}{M(y)} = \int_{[\underline{\gamma}, \Gamma]} t^{-1/g} K_y^*(dg) \longrightarrow t^{-1/\xi}$$

almost surely, by Lemma A.5 and boundedness and continuity of $g \mapsto t^{-1/g}$ on $[\underline{\gamma}, \Gamma]$. Lemma A.4 transfers the limit to $\mathbb{P}(Y > ty \mid \mathcal{G})/\mathbb{P}(Y > y \mid \mathcal{G})$. Convergence simultaneously for all $t > 0$ follows from rational t , monotonicity of the ratio in t , and continuity of $t \mapsto t^{-1/\xi}$.

For the fibre identity (3), fix $u \in I_\varepsilon$ and apply the above on the probability space $([0, 1]^{p-1}, K_j(u, \cdot))$ with the trivial σ -field; $\xi_j(u)$ is then the essential supremum of $v \mapsto \gamma(u, v)$ under $K_j(u, \cdot)$. Let $m = \max_v \gamma(u, v)$, attained by compactness and continuity. The essential supremum is at most m . Conversely, for $\eta > 0$ the set $\{v : \gamma(u, v) > m - \eta\}$ is open and nonempty, so has positive $K_j(u, \cdot)$ -probability under (S); the essential supremum is at least $m - \eta$ for every $\eta > 0$. $\square$

A.3 Population results

Lemma A.6 (Continuity of the envelope). *Under (C1), each $\xi_j(u) = \max_v \gamma(u, v)$ is continuous on $[0, 1]$, with modulus bounded by the modulus of continuity of γ .*

Proof. γ is uniformly continuous on the compact cube; if $|u - u'| \leq \delta$ then $|\gamma(u, v) - \gamma(u', v)| \leq \varpi(\delta)$ for every v , and taking maxima preserves the bound. $\square$

Proof of Proposition 2.2. Claim 1. For $j \notin A$, $\gamma(u, v) = g\{(u, v)_A\}$ does not involve u , and as v ranges over the fibre the active subvector ranges over all of $[0, 1]^s$; by (3), $\xi_j(u) = \max_w g(w) = \gamma^*$ for every u .

Claim 2. By (3), $\xi_j(u) = \gamma^*$ exactly when the fibre through u intersects M , i.e. when $u \in \pi_j(M)$.

Claim 3. $u \mapsto \gamma^* - \xi_j(u)$ is continuous (Lemma A.6), nonnegative, and by Claim 2 vanishes on I_ε exactly on $I_\varepsilon \cap \pi_j(M)$. A continuous nonnegative function has positive integral over I_ε if and only if its positivity set has positive Lebesgue measure.

Final statement. Write M^A for the maximizer set of g in $[0, 1]^s$; then $M = \{u : u_A \in M^A\}$ and, for $j \in A$, $\pi_j(M)$ is the projection of M^A , a finite set when M^A is finite; its intersection with I_ε is Lebesgue-null and Claim 3 applies. $\square$

Proof of Proposition 2.3. Every x on the fibre $x_j = u$ has $\|x_A - u_A^*\| \geq |u - u_j^*|$, so every point of the fibre satisfies $\gamma(x) \leq \gamma^* - c|u - u_j^*|^\nu$; taking the fibre maximum gives $\xi_j(u) \leq \gamma^* - c|u - u_j^*|^\nu$. Averaging over $u \in I_\varepsilon$ gives the score bound. For $\nu = 2$, $\mathbb{E}(V - u_j^*)^2 = \text{Var}(V) + (1/2 - u_j^*)^2 = (1 - 2\varepsilon)^2/12 + (1/2 - u_j^*)^2$. $\square$

A.4 Uniform profile bound

Inside proofs the empirical-rank quantities carry a superscript R when the distinction from frozen-location quantities matters. The scheme parallels the proof of Theorem 3 in Gardes and Podgorny [9] — oscillation control, exponential-spacings representation, slowly varying remainder — with the covering net over their compact matrix class replaced by exact union bounds over p coordinates and S_n window states, and their density conditions (H.1)–(H.2) replaced by the deterministic rank geometry.

Lemma A.7 (Rank displacement). *Let $\widehat{U}_{ij} = R_{ij}/(n+1)$ and*

$$D_n(\eta) = \left\{ \max_{j \leq p} \max_{i \leq n} |U_{ij} - \widehat{U}_{ij}| \leq \eta + \delta_n \right\}.$$

Then $\mathbb{P}\{D_n(\eta)^c\} \leq 2pe^{-2n\eta^2}$.

Proof. Fix j and let $F_{n,j}$ be the empirical distribution function of the U_{ij} , $i \leq n$. By the Dvoretzky–Kiefer–Wolfowitz inequality with Massart’s constant [14], $\sup_t |F_{n,j}(t) - t| \leq \eta$ outside probability $2e^{-2n\eta^2}$. On that event, continuity gives $F_{n,j}(U_{(r)j}) = r/n$, hence $|U_{(r)j} - r/n| \leq \eta$ and

$$|U_{(r)j} - r/(n+1)| \leq \eta + \frac{r}{n(n+1)} \leq \eta + \delta_n$$

for every rank r simultaneously. A union bound over j finishes; no union over ranks is needed. $\square$

Lemma A.8 (Deterministic window counts). *For every j , every realization and every $u \in I_\varepsilon$, $M_j(u) \geq m_n^-$ and $k_j(u) \geq K_n$; moreover $M_j(u)$ is a nonrandom function of u .*

Proof. The ranks form a permutation of $1, \dots, n$, so $M_j(u) = \#\{r \in \{1, \dots, n\} : |r/(n+1) - u| \leq h\}$, a deterministic function of u : the number of integers in $[x(u), y(u)]$ with $x(u) = \max\{(u-h)(n+1), 1\}$ and $y(u) = \min\{(u+h)(n+1), n\}$. Fix $u \in I_\varepsilon$ and write $m^- = m_n^- = \lfloor \{h + \min(h, \varepsilon)\}(n+1) \rfloor - 1$; note $h + \min(h, \varepsilon) \leq \min(2h, h + \varepsilon)$. Four cases.

If neither endpoint is clipped, the interval has length $2h(n+1)$ and contains at least $\lfloor 2h(n+1) \rfloor - 1 \geq m^-$ integers.

If only the left endpoint is clipped, then $x(u) = 1$ and the count is $\lfloor y(u) \rfloor$. Either $y(u) = (u+h)(n+1) \geq (\varepsilon+h)(n+1)$, so the count is at least $\lfloor (h+\varepsilon)(n+1) \rfloor \geq m^- + 1$; or $y(u) = n$ and the count is $n \geq \lfloor (h+\varepsilon)(n+1) \rfloor - 1 \geq m^-$, using $h \leq 1 - \varepsilon$, i.e. $(h+\varepsilon)(n+1) \leq n+1$.

If only the right endpoint is clipped, then $y(u) = n$ and the count is $n - \lceil x(u) \rceil + 1 \geq n - x(u) \geq n - (1 - \varepsilon - h)(n+1) = (h+\varepsilon)(n+1) - 1$, hence at least $\lceil (h+\varepsilon)(n+1) \rceil - 1 \geq m^-$.

If both endpoints are clipped, the count is n , which is at least m^- as in the second case. In every case $M_j(u) \geq m^-$, and $k_j(u) = \lfloor \alpha M_j(u) \rfloor \geq \lfloor \alpha m_n^- \rfloor = K_n$. $\square$

Lemma A.9 (Finite-state structure). *For fixed j and fixed ranks, $u \mapsto \mathcal{W}_j(u)$ takes at most $S_n = 4n + 1$ distinct values on $[0, 1]$.*

Proof. The membership indicator of observation i changes only at $\hat{U}_{ij} \pm h$: at most $2n$ breakpoints, hence at most $2n + 1$ constancy intervals plus the $2n$ breakpoint states. $\square$

Lemma A.10 (Conditional PIT and Rényi representation). *Fix j and condition on the column $\mathbf{U}_j = (U_{1j}, \dots, U_{nj})$. Under (E1):*

1. $V_{ij} = 1 - F_{Y|j}(Y_i | U_{ij})$, $i \leq n$, are i.i.d. standard uniform, and $Y_i = Q_j(1/V_{ij}, U_{ij})$ almost surely;
2. for any index set $\mathcal{W}$ of size m , deterministic given $\mathbf{U}_j$, with $V_{(1)} < \dots < V_{(m)}$ the ordered PIT values of its members and $k = \lfloor \alpha m \rfloor$,

$$L_k = \frac{1}{k} \sum_{r=1}^k \log \frac{V_{(k+1)}}{V_{(r)}}$$

is distributed as the mean of k i.i.d. standard exponentials, independent of $V_{(k+1)}$.

Proof. (i) Given $\mathbf{U}_j$, the pairs (U_{ij}, Y_i) are independent in i and Y_i has continuous conditional distribution function $F_{Y|j}(\cdot | U_{ij})$, so V_{ij} is standard uniform. For the representation, $Q_j(1/V_{ij}, U_{ij}) = F_{Y|j}^{-1}(F_{Y|j}(Y_i | U_{ij}) | U_{ij}) \leq Y_i$ always, with strict inequality only when Y_i lies in the interior of an interval on which $F_{Y|j}(\cdot | U_{ij})$ is constant; each such interval has conditional probability zero and there are countably many, so the representation holds almost surely. Strict monotonicity is not needed. Conditioning on the whole column, rather than on window counts as in Gardes and Podgorny [9, Appendix, Lemma 4], makes the window sets deterministic (Lemma A.8); only coordinate j is conditioned on, and no independence across coordinates is used.

(ii) With $W_i = -\log V_i$ standard exponentials and order statistics $W_{(1)} < \dots < W_{(m)}$, $W_{(m-r+1)} = -\log V_{(r)}$, Abel summation gives

$$kL_k = \sum_{r=1}^k \{W_{(m-r+1)} - W_{(m-k)}\} = \sum_{i=1}^k i \{W_{(m-i+1)} - W_{(m-i)}\},$$

and by the Rényi representation of exponential order statistics [16] the normalized top spacings $i\{W_{(m-i+1)} - W_{(m-i)}\}$, $i \leq k$, are i.i.d. standard exponential and independent of $W_{(m-k)}$, hence of $V_{(k+1)}$. $\square$

Lemma A.11 (Frozen-location Hill expansion). *Fix j , condition on $\mathbf{U}_j$, and fix a window state $\mathcal{W}$ of size m realized at $u \in I_\varepsilon$, with $k = \lfloor \alpha m \rfloor$ and Hill statistic H . On the event $D_n(\eta)$ intersected with $\{V_{ij} \in [\tau, 1 - \tau] \ \forall i\}$ and $\{V_{(k+1)} \leq 3\alpha\}$,*

$$|H - \xi_j(u)| \leq 2\omega_n(h + \eta + \delta_n, \tau) + \xi_j(u)|L_k - 1| + b_n(3\alpha)L_k. \quad (\text{A.3})$$

Proof. Spatial freezing. Each $i \in \mathcal{W}$ has $|\widehat{U}_{ij} - u| \leq h$, hence $|U_{ij} - u| \leq r_n := h + \eta + \delta_n$ on $D_n(\eta)$. With $V_{ij} \in [\tau, 1 - \tau]$, the definition of ω_n gives $|\log Q_j(1/V_{ij}, U_{ij}) - \log Q_j(1/V_{ij}, u)| \leq \omega_n(r_n, \tau)$ for every member. Let $\check{H}$ be the Hill statistic of the frozen responses $\check{Y}_i = Q_j(1/V_{ij}, u)$. By Lemma A.1, ordered log-responses differ by at most $\omega_n(r_n, \tau)$, and the Hill statistic is an average of differences of two of them, so $|H - \check{H}| \leq 2\omega_n(r_n, \tau)$.

Exact decomposition. Since $s \mapsto Q_j(s, u)$ is nondecreasing, $\check{Y}_{(m-r+1)} = Q_j(1/V_{(r)}, u)$, and $\log Q_j(s, u) = \xi_j(u) \log s + \log \ell_j(s, u)$ yields

$$\check{H} = \xi_j(u) L_k + R, \quad R = \frac{1}{k} \sum_{r=1}^k \log \frac{\ell_j(1/V_{(r)}, u)}{\ell_j(1/V_{(k+1)}, u)}.$$

Remainder. On $\{V_{(k+1)} \leq 3\alpha\}$, each ratio is $\ell_j(ts, u)/\ell_j(s, u)$ with $s = 1/V_{(k+1)} \geq (3\alpha)^{-1}$ and $t = V_{(k+1)}/V_{(r)} > 1$, so $|\log(\cdot)| \leq b_n(3\alpha) \log(V_{(k+1)}/V_{(r)})$; averaging gives $|R| \leq b_n(3\alpha) L_k$. $\square$

Theorem A.12 (Finite-sample uniform profile bound). *Assume (C1)–(C2), (S), the structural part of (E1) (continuity of every $F_{Y|j}(\cdot | u)$), hence the quantile representation), that $b_n(3\alpha)$ and $\omega_n(\cdot, \cdot)$ are finite, $0 < h \leq 1 - \varepsilon$, $0 < \alpha \leq 1/3$, and $K_n \geq 2$. For every $\eta > 0$, $0 < t \leq 1$, $0 < \tau < 1/2$,*

$$\begin{aligned} \mathbb{P} \left\{ \max_{j \leq p} \sup_{u \in I_\varepsilon} |\widehat{\xi}_j(u) - \xi_j(u)| > 2\omega_n(h + \eta + \delta_n, \tau) + \Gamma t + (1 + t)b_n(3\alpha) \right\} \\ \leq \underbrace{2pe^{-2n\eta^2}}_{\text{ranks}} + \underbrace{2pn\tau}_{\text{PIT truncation}} + \underbrace{2pS_n e^{-K_n t^2/4}}_{\text{spacings}} + \underbrace{pS_n e^{-(K_n+1)/4}}_{\text{threshold}}. \end{aligned} \quad (\text{A.4})$$

Proof. Call a window state *admissible* for coordinate j if it equals $\mathcal{W}_j(u)$ for some $u \in I_\varepsilon$; by Lemmas A.8 and A.9 there are at most S_n admissible states per coordinate and each has $k \geq K_n \geq 2$, so L_k and $V_{(k+1)}$ are well defined on every one of them. (States realized only outside I_ε may have smaller windows; they are simply not needed.) Define the exceptional events $T_1 = D_n(\eta)^c$; $T_2 = \{\exists i, j : V_{ij} \notin [\tau, 1 - \tau]\}$; $T_3 = \{\exists j, \text{ admissible state } \mathcal{W} : |L_k - 1| > t\}$; $T_4 = \{\exists j, \text{ admissible state } \mathcal{W} : V_{(k+1)} > 3\alpha\}$.

Outside $T_1 \cup T_2 \cup T_3 \cup T_4$, fix j and $u \in I_\varepsilon$: the realized window is an admissible state, and Lemma A.11 with $\xi_j \leq \Gamma$, $|L_k - 1| \leq t$, $L_k \leq 1 + t$ gives the displayed deviation bound; the modulus $\omega_n(h + \eta + \delta_n, \tau)$ covers every selected observation because its definition allows the displaced point $v = U_{ij}$ to range over all of $[0, 1]$.

Probabilities: $\mathbb{P}(T_1) \leq 2pe^{-2n\eta^2}$ is Lemma A.7; $\mathbb{P}(T_2) \leq 2pn\tau$ since each V_{ij} is uniform. For T_3, T_4 , condition on $\mathbf{U}_j$; the admissible states are then deterministic with $k \geq K_n$, Lemma A.10 applies to each, and Lemmas A.2 and A.3 give conditional bounds $2e^{-kt^2/4} \leq 2e^{-K_n t^2/4}$ and $e^{-(k+1)/4} \leq e^{-(K_n+1)/4}$, both decreasing in k . A union bound over the at most pS_n admissible states and integration over $\mathbf{U}_j$ complete the proof; overlapping states and dependent coordinates cost nothing beyond the union bound. $\square$

A.5 Scores, concentration, and screening

Recall the score grid $G = \{r\delta_n : r = 1, \dots, n\} \cap I_\varepsilon$, of size $L_n = |G|$, and that $K_n \geq 2$ makes every grid point usable, so $\widehat{\Psi}_j = L_n^{-1} \sum_{u \in G} \widehat{\xi}_j(u)$.

Lemma A.13 (Quadrature). *If $L_n \geq 1$, then for every j ,*

$$\left| L_n^{-1} \sum_{u \in G} \xi_j(u) - \Psi_j \right| \leq q_n := \omega_\xi(\delta_n) + \frac{4\Gamma\delta_n}{1 - 2\varepsilon}. \quad (\text{A.5})$$

Proof. Write the grid points $u_1 < \dots < u_{L_n}$, consecutive multiples of δ_n . If $L_n = 1$ then $|I_\varepsilon| < 2\delta_n$ and the left side is at most $\Gamma < 4\Gamma\delta_n/|I_\varepsilon|$. Assume $L_n \geq 2$ and partition I_ε exactly: $C_1 = [\varepsilon, u_1 + \delta_n/2]$, $C_{L_n} = [u_{L_n} - \delta_n/2, 1 - \varepsilon]$, and $C_r = [u_r - \delta_n/2, u_r + \delta_n/2]$ otherwise. Interior cells have length δ_n , boundary cells length in $[\delta_n/2, 3\delta_n/2]$, and every point of C_r is within δ_n of u_r . Then

$$\left| \frac{1}{L_n} \sum_r \xi_j(u_r) - \frac{1}{|I_\varepsilon|} \int_{I_\varepsilon} \xi_j \right| \leq \frac{1}{|I_\varepsilon|} \sum_r \int_{C_r} |\xi_j - \xi_j(u_r)| + \Gamma \sum_r \left| \frac{\lambda(C_r)}{|I_\varepsilon|} - \frac{1}{L_n} \right|.$$

The first term is at most $\omega_\xi(\delta_n)$. For the second, both weight vectors sum to one; $|I_\varepsilon| = \sum_r \lambda(C_r) \in [(L_n - 1)\delta_n, (L_n + 1)\delta_n]$ gives $||I_\varepsilon|/L_n - \delta_n| \leq \delta_n/L_n$, so interior cells contribute at most δ_n in total and the two boundary cells at most $\delta_n/2 + \delta_n/L_n \leq \delta_n$ each; the sum is at most $3\delta_n$, and the second term is at most $3\Gamma\delta_n/|I_\varepsilon| \leq 4\Gamma\delta_n/|I_\varepsilon|$. $\square$

Theorem A.14 (Finite-sample score bound). *Under the conditions of Theorem A.12 (with in addition $\omega_\xi(\delta_n) < \infty$) and $L_n \geq 1$, for every $\eta > 0$, $0 < t \leq 1$, $0 < \tau < 1/2$,*

$$\mathbb{P} \left\{ \max_{j \leq p} |\hat{\Psi}_j - \Psi_j| > B_n(\eta, t, \tau) + q_n \right\} \leq 2pe^{-2n\eta^2} + 2pn\tau + 2pS_ne^{-K_nt^2/4} + pS_ne^{-(K_n+1)/4}, \quad (\text{A.6})$$

where $B_n(\eta, t, \tau) = 2\omega_n(h + \eta + \delta_n, \tau) + \Gamma t + (1 + t)b_n(3\alpha)$. If moreover $B_n(\eta, t, \tau) + q_n < \Delta_{\min}/2$, the same right-hand side bounds the probability that some active score fails to precede every inactive score; on the complementary event, $A \subseteq \hat{A}_d$ for every $d \geq s$, and $\hat{A}_s = A$.

Proof. Pathwise,

$$|\hat{\Psi}_j - \Psi_j| \leq \sup_{u \in I_\varepsilon} |\hat{\xi}_j(u) - \xi_j(u)| + \left| L_n^{-1} \sum_{u \in G} \xi_j(u) - \Psi_j \right|,$$

and Lemma A.13 bounds the second term by q_n on every sample path; reusing the same observations for estimation and averaging therefore costs no extra probability. Theorem A.12 gives (A.6). On the event that every score error is below $\Delta_{\min}/2$: inactive coordinates have $\Psi_j = \gamma^*$ (Proposition 2.2), hence $\hat{\Psi}_j > \gamma^* - \Delta_{\min}/2$; active coordinates have $\Psi_j \leq \gamma^* - \Delta_{\min}$, hence $\hat{\Psi}_j < \gamma^* - \Delta_{\min}/2$. Every active score precedes every inactive score, and the screening and recovery statements follow deterministically. $\square$

Proof of Theorem 3.1. Choose $\eta_n = \sqrt{\log(pn)/n}$, $\tau = \tau_n = (pn)^{-2}$, and $t_n = \sqrt{(4/K_n) \log\{pS_n(pn)\}}$. The four probability terms in (A.6) are then each $o(1)$: the first is $2pe^{-2\log(pn)} = 2p(pn)^{-2} \rightarrow 0$; the second is $2pn(pn)^{-2} \rightarrow 0$; the third is $2pS_n\{pS_n(pn)\}^{-1} = 2(pn)^{-1} \rightarrow 0$; the fourth is $pS_ne^{-(K_n+1)/4} \rightarrow 0$ because $\log(pS_n) \leq C \log(pn) = o(K_n)$ by the second condition in (7) and $K_n \geq c\alpha nh$. The same condition makes $t_n \leq 1$ for large n , as required by Lemma A.2, and $\Gamma t_n = O(\sqrt{\log(pn)/(n\alpha h)})$.

It remains to check that the bias part of $B_n(\eta_n, t_n, \tau_n) + q_n$ is $o(\Delta_{\min})$. First, $\log(pn) = o(nh^2)$ gives $\eta_n + \delta_n = o(h)$, so $h + \eta_n + \delta_n \leq C_0 h$ eventually and $2\omega_n(h + \eta_n + \delta_n, \tau_n) \leq 2\omega_n(C_0 h, \tau_n) = o(\Delta_{\min})$ by (E3). Second, $(1 + t_n)b_n(3\alpha) \leq 2b_n(3\alpha) = o(\Delta_{\min})$ by (E2). Third, $q_n = \omega_\xi(\delta_n) + O(1/n)$: the first part is $o(\Delta_{\min})$ by (E1), and $1/n \leq \sqrt{\log(pn)/(n\alpha h)}$ since $\alpha h \leq 1 \leq n \log(pn)$, so the second part is absorbed into the stochastic rate. Collecting terms, $\max_j |\hat{\Psi}_j - \Psi_j| \leq O(\sqrt{\log(pn)/(n\alpha h)}) + o(\Delta_{\min})$ outside a vanishing probability. $\square$

Proof of Theorem 3.2 and Corollary 3.3. Under (8), $\sqrt{\log(pn)/(n\alpha h)} = o(\Delta_{\min})$, so the choices in the previous proof give $B_n(\eta_n, t_n, \tau_n) + q_n < \Delta_{\min}/2$ for n large, and the second part of Theorem A.14 applies: the failure probability of active–inactive separation is bounded by the four vanishing terms. Sure screening for every $d \geq s$ and exact recovery at $d = s$ follow. $\square$

References

- [1] Anass Aghbalou, François Portier, Anne Sabourin, and Chen Zhou. Tail inverse regression: Dimension reduction for prediction of extremes. *Bernoulli*, 30(1):503–533, 2024. doi: 10.3150/23-BEJ1606.
- [2] Julyan Arbel, Stéphane Girard, and Hadrien Lorenzo. Shrinkage for extreme partial least-squares, 2024. arXiv:2403.09503v2.
- [3] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, Oxford, 2013. doi: 10.1093/acprof:oso/9780199535255.001.0001.
- [4] Meryem Bousebata, Geoffroy Enjolras, and Stéphane Girard. Extreme partial least-squares. *Journal of Multivariate Analysis*, 194:105101, 2023. doi: 10.1016/j.jmva.2022.105101.
- [5] Abdelaati Daouia, Laurent Gardes, and Stéphane Girard. On kernel smoothing for extremal quantile regression. *Bernoulli*, 19(5B):2557–2589, 2013. doi: 10.3150/12-BEJ466.
- [6] Laurens de Haan and Ana Ferreira. *Extreme Value Theory: An Introduction*. Springer, New York, 2006. doi: 10.1007/0-387-34471-3.
- [7] Jianqing Fan and Jinchi Lv. Sure independence screening for ultrahigh dimensional feature space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(5): 849–911, 2008. doi: 10.1111/j.1467-9868.2008.00674.x.
- [8] Laurent Gardes. Tail dimension reduction for extreme quantile estimation. *Extremes*, 21(1): 57–95, 2018. doi: 10.1007/s10687-017-0300-x.
- [9] Laurent Gardes and Alexandre Podgorny. Dimension reduction for the estimation of the conditional tail index. *Scandinavian Journal of Statistics*, 52(3):1444–1476, 2025. doi: 10.1111/sjos.12792.
- [10] Laurent Gardes and Gilles Stupfler. Estimation of the conditional tail index using a smoothed local Hill estimator. *Extremes*, 17(1):45–75, 2014. doi: 10.1007/s10687-013-0174-5.
- [11] Xuming He, Lan Wang, and Hyokyoung Grace Hong. Quantile-adaptive model-free variable screening for high-dimensional heterogeneous data. *The Annals of Statistics*, 41(1):342–369, 2013. doi: 10.1214/13-AOS1087.
- [12] Bruce M. Hill. A simple general approach to inference about the tail of a distribution. *The Annals of Statistics*, 3(5):1163–1174, 1975. doi: 10.1214/aos/1176343247.
- [13] Runze Li, Wei Zhong, and Liping Zhu. Feature screening via distance correlation learning. *Journal of the American Statistical Association*, 107(499):1129–1139, 2012. doi: 10.1080/01621459.2012.695654.
- [14] Pascal Massart. The tight constant in the Dvoretzky–Kiefer–Wolfowitz inequality. *The Annals of Probability*, 18(3):1269–1283, 1990. doi: 10.1214/aop/1176990746.
- [15] Michael Redmond and Alok Baveja. A data-driven software tool for enabling cooperative information sharing among police departments. *European Journal of Operational Research*, 141(3):660–678, 2002. doi: 10.1016/S0377-2217(01)00264-8.

- [16] Alfréd Rényi. On the theory of order statistics. *Acta Mathematica Academiae Scientiarum Hungaricae*, 4:191–231, 1953. doi: 10.1007/BF02127580.
- [17] Yuya Sasaki, Jing Tao, and Yulong Wang. High-dimensional tail index regression, 2026. arXiv:2403.01318v3.
- [18] Hansheng Wang and Chih-Ling Tsai. Tail index regression. *Journal of the American Statistical Association*, 104(487):1233–1240, 2009. doi: 10.1198/jasa.2009.tm08458.
- [19] Takuma Yoshida and Yuta Umezu. Sure independence screening for covariate-dependent extreme value index estimation, 2026. arXiv:2410.15705v3.