



# **Mapping complex traits in populations: Genome-wide association studies (GWAS)**



# QTL mapping in a cross/ pedigree

- Localize gene(s) affecting phenotypic difference between original strains/ people
  - Let's say you map a gene causing diabetes in one family to 18p22 (strong effect next to a marker)
  - You map in ***another*** family having diabetes, and you see no effect of 18p22 region
    - What happened?



# Another approach: mapping in populations by association studies (e.g., GWAS)

- We discussed this last week for simple traits
  - If a marker is very close to the disease-causing gene, individuals having one allele will be more likely to have the disease than individuals having the other allele
- Same principle true even if many genes involved




# Test many, many, many markers

- Some have an allele associated with disease
  - AA: prob disease: 10%
  - aa: prob disease: 1%
  - Chi-square or other statistic says unlikely difference by chance
- Many have no association
  - BB: prob disease: 1.5%
  - bb: prob disease: 1.6%

Visualize with  
something like  
LOD plot

# Test many, many, many markers


- Some have an allele associated with disease
  - AA: prob disease: 10%
  - aa: prob disease: 1%
  - Chi-square or other statistic says unlikely difference by chance
- Many have no association
  - BB: prob disease: 1.5%
  - bb: prob disease: 1.6%



From Feb 2012 *Nature Genetics*  
Azoospermia in Chinese men

# Often find not one but many possible factors

- From Feb 2012 issue of *Nature Genetics*:
  - A genome-wide association study in Han Chinese identifies multiple susceptibility loci for IgA nephropathy (8p23, 17p13, 22q12).



# Problem: multiple comparisons

- From lab, chi square test gave probability of getting that association “by chance”
  - If you roll two dice and get snake-eyes twice in a row, are they loaded?
    - ~0.1% chance that would happen, but not ZERO



# Problem: multiple comparisons

- From lab, chi square test gave probability of getting that association “by chance”
  - If you roll two dice and get snake-eyes twice in a row, are they loaded?
    - ~0.1% chance that would happen, but not ZERO
  - What if rolled dice 1,000,000 times- what is chance you’d get snake-eyes twice in a row SOMETIME in there?





# Problem: multiple comparisons

- From lab, chi square test gave probability of getting that association “by chance”
  - If you roll two dice and get snake-eyes twice in a row, are they loaded?
    - ~0.1% chance that would happen, but not ZERO
  - What if rolled dice 1,000,000 times- what is chance you’d get snake-eyes twice in a row SOMETIME in there? **>99.99%!**



# Problem: multiple comparisons

- Same problem when test 1,000,000 SNPs for association with disease of interest
  - Very likely to get an association that looks strong by chance but no real biology
- As a result, instead of using 5% probability as a cut-off, often use  $\sim 0.00001\%$  probability




# Problem: multiple comparisons

- Same problem when test 1,000,000 SNPs for association with disease of interest
  - Very likely to get an association that looks strong by chance but no real biology
- As a result, instead of using 5% probability as a cut-off, often use  $\sim 0.00001\%$  probability
  - BUT this requires a VERY strong association...



# Pedigree/cross vs. Population

- Pedigree/ cross analyses find genetic factors causing differences *in that family*.
  - Typically, high power for small number of genes
- Population association studies try to find genetic factors causing variation *across the population*.
  - Typically, lower power but can find more genes and localizes more precisely



# Why???

- In cross/ pedigree, limited to mapping genes with effects in that cross, BUT the alleles you're mapping are abundant
  - In simple cross, initially 25% or higher


$Zz \times zz$   
↓

- In population, you can theoretically map alleles at any gene causing effect, BUT you may have *different* very rare variants causing the same effect/disease

aazz aazz aazz aazz **A**aaz  
aa**Z**z aazz aazz aazz aazz

# Genome-wide association studies in populations work for mapping “common disease variants”

- ... but are most disease variants common?
- ... how often do *different rare* mutations cause the same disease?



## **Also, can see differences among ethnic groups for associations**

- Sometimes see a marker allele associated with disease in one ethnic group but not another



# Type II Diabetes

- Four SNPs identified associated with Type II diabetes in East Asian subjects
- One of these four SNPs (on chr 11) had no detectable effect in studies of European populations





# Image Credits, Unit 7-3

- Microarray, © 2006 Schutz, CC by-SA 3.0, en.wikipedia.org.
- Afghan child receiving vaccination, © 2010 UNICEF Sverige, CC by 2.0, [www.fotopedia.com](http://www.fotopedia.com).
- LOD scatterplot, © 2012 Hu Z. et al and Nature Genetics, all rights reserved, <http://www.nature.com/ng/journal/v44/n2/full/ng.1040.html>, Nature Genetics 44, 183–186 (2012)
- Nephropathy kidney cells, © 2008 Karamadoukis et al, CC by 2.0, [www.jmedicalcasereports.com](http://www.jmedicalcasereports.com)
- Stick figures and DNA, © vishnukumar, all rights reserved, [www.photoxpress.com](http://www.photoxpress.com)
- Multicolor handprints, © Willee Cole, all rights reserved, [www.photoxpress.com](http://www.photoxpress.com)