



NATURAL SELECTION: Types acting on traits

Duke
UNIVERSITY




Natural Selection:

Quantitative Traits vs. Single Locus

- Natural selection can also be studied in the context of phenotypes, as in our discussions of heritability
 - Don't necessarily need to know underlying genes to infer type of selection operating




Types of selection, inferred from phenotypes



Directional selection (phenotypes)


- Individuals at *one* end of distribution favored
 - e.g., “big” or “small individuals
- Causes change in *mean* of population over time



Directional selection (phenotypes)

- Individuals at *one* end of distribution favored
 - e.g., “big” or “small individuals
- Causes change in *mean* of population over time

Pink salmon weight, in Canada
1951-1974




Directional selection (phenotypes)

Pink salmon weight, in Canada
1951-1974


Stabilizing selection (phenotypes)

- Individuals in *middle* of distribution favored
- No change in mean, but loss of extremes




Stabilizing selection (phenotypes)

- Individuals in *middle* of distribution favored
- No change in mean, but loss of extremes



Bumpus 1898 study of 136 house sparrows after Rhode Island storm
64 died, 72 survived



Mean of survivors = Mean of dead
All >28 or <23 died


Stabilizing selection (phenotypes)

- Individuals in *middle* of distribution favored
 - No change in mean, but loss of extremes
- Other example:
HUMAN BIRTHWEIGHT




Disruptive selection (phenotypes)

- Individuals at *both ends* of distribution favored
- No change in mean, but loss of intermediate phenotypes




Disruptive selection (phenotypes)

- Individuals at *both ends* of distribution favored
- No change in mean, but loss of intermediate phenotypes




Beak size in female African finch:



Specializes on seeds of two sedge species

Types of selection, inferred from phenotypes





Final thoughts – is selection “good” or “bad” for population?



- Natural selection preferentially reduces/ eliminates “bad” genotypes
- “Average” fitness of all individuals *remaining in population after selection* goes up!
 - Since “bad alleles” are removed, simple directional selection gives long-term improvement to population



Directional selection improves average population fitness

- **Fisher's fundamental theorem of natural selection**
 - The rate of increase in fitness is equals the genetic variance in fitness.



Image Credits, Unit 12-3

- Selection types, © 2007 Pearson Education Inc., all rights reserved, Figure 9.5, Evolutionary Analysis 4/e by Freeman & Herron
- Graph of salmon weight selection, © Wiley & Sons, all rights reserved,
- Salmon fishing, © Newsweek, all rights reserved
- Baby being weighed, © 2005 Drpoulette, CC by 2.0, en.wikipedia.org
- Human birthweight selection, © Wiley & Sons, all rights reserved
- Female African finch beak sizes, © 1993 T. B. Smith, Disruptive selection and the genetic basis of bill size polymorphism in the African finch, *Pyrenestes*. *Nature* 363: 618–620
- Photo by Bletchley, © 2008, CC by-SA 3.0, en.wikipedia.org