



ORDER / DISORDER


Temperature-path fingerprints



ANGULAR ATLAS / Glycerol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Ethylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / DMSO-inspired


10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Propylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum



ANGULAR ATLAS / Formamide-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Benzyl alcohol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




Glycerol-inspired / Partial-charge Coulomb + LJ
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Ethylene glycol-inspired / Partial-charge Coulomb + LJ
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




DMSO-inspired / Partial-charge Coulomb + LJ
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Propylene glycol-inspired / Partial-charge Coulomb + LJ
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Formamide-inspired / Partial-charge Coulomb + LJ
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Benzyl alcohol-inspired / Partial-charge Coulomb + LJ
PCHIP in T, log(P) · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




ORDER / DISORDER


Temperature-path fingerprints



ANGULAR ATLAS / Glycerol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Ethylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / DMSO-inspired


10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Propylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum



ANGULAR ATLAS / Formamide-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Benzyl alcohol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




Glycerol-inspired / Lennard-Jones only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Ethylene glycol-inspired / Lennard-Jones only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




DMSO-inspired / Lennard-Jones only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Propylene glycol-inspired / Lennard-Jones only
PCHIP in T , $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Formamide-inspired / Lennard-Jones only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Benzyl alcohol-inspired / Lennard-Jones only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




ORDER / DISORDER


Temperature-path fingerprints



ANGULAR ATLAS / Glycerol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum



ANGULAR ATLAS / Ethylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / DMSO-inspired


10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Propylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum



ANGULAR ATLAS / Formamide-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Benzyl alcohol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




Glycerol-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Ethylene glycol-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




DMSO-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Propylene glycol-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Formamide-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Benzyl alcohol-inspired / Partial-charge Coulomb + WCA
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




ORDER / DISORDER


Temperature-path fingerprints



ANGULAR ATLAS / Glycerol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Ethylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / DMSO-inspired


10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Propylene glycol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum



ANGULAR ATLAS / Formamide-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




ANGULAR ATLAS / Benzyl alcohol-inspired 10 kPa · pooled maps and instantaneous multipole spectrum




Glycerol-inspired / WCA repulsion only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Ethylene glycol-inspired / WCA repulsion only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




DMSO-inspired / WCA repulsion only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Propylene glycol-inspired / WCA repulsion only
PCHIP in T , $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order




Formamide-inspired / WCA repulsion only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order




FCC reference order



Benzyl alcohol-inspired / WCA repulsion only
PCHIP in T, $\log(P)$ · gold = simulated samples · NOT a phase boundary

O_6 multipolar order



FCC reference order

