



**UNIVERSIDAD
DE GRANADA**

Facultad de Ciencias

Physics Degree

Complex Systems Project

Quantum Chaos and the Statistics of Energy Levels

**María de los Ángeles Lara Consuegra
Pablo Orellana Chornyak**

05/2026

Abstract

In this report, we study how classical chaos shows up in quantum mechanics by looking at the statistical properties of energy spectra in 2D billiards. To do this, we solved the time-independent Schrödinger equation using a finite difference approach and diagonalized the resulting sparse matrices. We analyzed two specific shapes with completely different classical dynamics: a rectangular billiard (which is an integrable system) and a quarter-stadium billiard (which is a chaotic system).

Índice

1	Introduction	3
2	Methods	3
2.1	Discretisation of the system: Mascara Method	4
2.2	Finite Difference Method	4
2.3	Eigenvalues distribution	6
3	Results	6
3.1	Statistical analysis of spacing distribution	6
3.2	High-energy states visualization	8
3.2.1	Ergodicity in the quantum system	9
4	Conclusions	12
	Referencias	12

1 Introduction

The study of complex systems allows to explore the transition between the classical order and the statistical manifestations in the quantum regime. As in classical mechanics chaos is defined as the exponential sensitivity to initial conditions, the linearity in the Schrödinger equation does not allow for a direct translation of this definition to quantum mechanics. However, the phenomena known as 'Quantum Chaos' is manifested throughout the statistical properties of the energy spectrum of the system.

The core of this project is the Bohigas-Giannoni-Schmit (BGS) conjecture, which states that the energy levels of a quantum system whose classical counterpart is chaotic will exhibit the same statistical properties as the eigenvalues of a random matrix ensemble.

To validate this hypothesis, we'll investigate the distribution of spacing between adjacent energy levels in two systems: one integrable and one chaotic.

The first one is the two-dimensional **rectangular billiard**, which is an **integrable system**. In this system, the random matrix theory (RMT) predicts that the distribution of level spacings follows a Poisson distribution, which is given by:

$$P(s) = e^{-s} \quad (1.1)$$

The second system is the **stadium billiard**, which is a **chaotic system**. In this case, RMT predicts that the distribution of level spacings follows the Wigner-Dyson distribution, which is given by:

$$P(s) = \frac{\pi}{2} s e^{-\frac{\pi}{4}s^2} \quad (1.2)$$

where s is the normalized spacing between adjacent energy levels in both cases.

To integrate the time independent Schrödinger equation for these systems, we will use the **Finite Difference Method** (FDM): this method allows us to discretize the spatial domain and solve for the energy eigenvalues and eigenfunctions of the system.

2 Methods

To be able to solve the Schrödinger Equation $H\psi = E\psi$, we'll first need to discretize the space and then solve the resulting eigenvalue problem. We'll need to impose the Dirichlet conditions to the domain, which means that the wavefunction ψ must vanish at the boundaries of the billiard.

$$\psi|_{\partial\Omega} = 0 \quad (2.1)$$

All the methods used in this project are implemented in Python, and the code is available in the GitHub repository [1]. To explain the followed procedure, we'll first introduce the Mascara method, which is used to discretize the system, and then the Finite Difference Method, which is used to solve the resulting eigenvalue problem.

2.1 Discretisation of the system: Mascara Method

To easily create different geometries, we decided to implement a finite difference method that takes different masks and computes the corresponding eigenvalues. The mask represents a grid containing the geometry of the problem. For each pixel, we determine whether it belongs to the geometry by assigning it a true or false value.

To carry out this project, we created three different masks: a rectangular mask, a stadium-shaped mask, and a quarter-stadium mask. The following figure shows these masks:




Figure 1: Different mask geometries used in this experiment. From left to right: a rectangular mask, a stadium-shaped mask, and a quarter-stadium mask.

The project aimed to investigate the stadium geometry, which produces chaotic behavior in the eigenvalue spectrum. However, we encountered a problem: the stadium has two symmetries, one with respect to the x -axis and another with respect to the y -axis. In Chapter 1.4 of the thesis by *Wisniacki, Diego A.*, it is explained that the eigenvalues are divided into four different sets because of these symmetries:

- even | even
- even | odd
- odd | even
- odd | odd

This produces four different Wigner–Dyson distributions, which, when combined, result in a Poisson distribution. By removing the symmetries and considering only a quarter of the stadium, the chaotic behavior becomes visible. [2]

2.2 Finite Difference Method

Once we have a grid of the system, where $x_i = i\Delta x$ and $y_j = j\Delta y$, being $i = 1, 2, \dots, n_x$ and $j = 1, 2, \dots, n_y$. As we know, the Hamiltonian for a two-dimensional system is given by:

$$H = -\frac{\hbar^2}{2m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + V(x, y) \quad (2.2)$$

Since in our systems we have a free particle, the potential $V(x, y)$ is zero everywhere inside the billiard and infinite outside. With this consideration, the Schrödinger equation can be rewritten as the Helmholtz equation:

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + k^2 \right) \psi(x, y) = 0 \quad (2.3)$$

where $k^2 = E$ (using units such that $\hbar^2/2m = 1$).

Now, we can discretize the second derivatives using finite differences.[3] If we consider one of the dimensions, for example the x -direction, we can approximate the wave function in the nearest points with a Taylor expansion:

$$\psi(x_i \pm \Delta x) = \psi(x_i) \pm \Delta x \psi'(x_i) + \frac{\Delta x^2}{2} \psi''(x_i) \pm O(\Delta x^3) \quad (2.4)$$

Reordenating terms, we can get an expression for the second derivative:

$$\psi''(x_i) = \frac{\psi(x_i + \Delta x) - 2\psi(x_i) + \psi(x_i - \Delta x)}{\Delta x^2} + O(\Delta x^2) \approx \frac{\psi_{i+1} - 2\psi_i + \psi_{i-1}}{\Delta x^2} \quad (2.5)$$

Taking this in consideration, and applying the same procedure for the y -direction, we'll apply the Helmholtz equation in its discretized form 2.6 to the point (x_i, y_j) of the grid.

$$\left(\frac{\psi_{i+1,j} - 2\psi_{i,j} + \psi_{i-1,j}}{\Delta x^2} + \frac{\psi_{i,j+1} - 2\psi_{i,j} + \psi_{i,j-1}}{\Delta y^2} + k^2 \psi_{i,j} \right) = 0 \quad (2.6)$$

Is more efficient to write the resulting equation in a matrix form, where the matrix H is given by:

$$H' = \begin{pmatrix} D & \frac{1}{\Delta x^2} & 0 & \dots & \frac{1}{\Delta y^2} & 0 & \dots \\ \frac{1}{\Delta x^2} & D & \frac{1}{\Delta x^2} & 0 & \dots & \frac{1}{\Delta y^2} & \dots \\ 0 & \frac{1}{\Delta x^2} & D & \frac{1}{\Delta x^2} & 0 & \dots & \dots \\ \vdots & 0 & \ddots & \ddots & \ddots & 0 & \vdots \\ \frac{1}{\Delta y^2} & \vdots & 0 & \frac{1}{\Delta x^2} & D & \frac{1}{\Delta x^2} & 0 \\ 0 & \frac{1}{\Delta y^2} & \vdots & \dots & \frac{1}{\Delta x^2} & D & \frac{1}{\Delta x^2} \\ \vdots & \dots & \dots & \dots & 0 & \frac{1}{\Delta x^2} & D \end{pmatrix} \quad (2.7)$$

This matrix is called the finite difference matrix (FDM) and it is a sparse matrix, since most of its elements are zero. It has a total of $n_x \times n_y$ elements. The diagonal elements are given by $D = k^2 - \left(\frac{2}{\Delta x^2} + \frac{2}{\Delta y^2} \right)$. The second diagonal elements are going by $\frac{1}{\Delta x^2}$ and it corresponds to the coupling between the nearest points in the x -direction. The elements that are n_x positions away from the diagonal are given by $\frac{1}{\Delta y^2}$ and it corresponds to the coupling between the nearest points in the y -direction. Since in our systems we have a free particle, the potential $V(x, y)$ is zero everywhere inside the billiard and infinite outside.

2.3 Eigenvalues distribution

Then we calculate the eigenvalues and the eigenvectors of the sparse matrix (2.7), with the help of the library `scipy`. Now we calculate the difference between consecutive eigenvalues, which is called the spacing distribution (and normalized), and we plot an histogram with those values. The number of **eigenvalues** calculated will be, for every case, of $n = 2000$. With this sample, we'll use **35 bins**, with each bin containing 57 values. To provide a statistical point of view, we'll also plot error bars that represents a trust interval of 95 % ($\pm 2\sigma$), where $\sigma = \sqrt{N}$, being N the number of values in a bin.

This histogram will give us information about the chaoticity of the system, since, as we know, the spacing distribution for a chaotic system follows a Wigner-Dyson distribution, while for an integrable system it follows a Poisson distribution.

3 Results

3.1 Statistical analysis of spacing distribution

First, we calculate the first 10000 eigenvalues of each system by diagonalizing the sparse matrix that we obtained applying the FDM 2.7. Then, with those arrays, we compute a new one that has the spacing distribution between consecutive eigenvalues, this is $s = E_{i+1} - E_i$. Since the Poisson and Wigner distributions are defined in the normalized space, we must normalize then the spacing by doing $s \rightarrow s / \langle s \rangle$.

Finally, we plot an histogram of the normalized spacing between adjacent eigenvalues, as it follows:




Figure 2: Blue bars represent the histogram spacings, with associated errorbars that follows a counting Poisson error in a 95 % ($\pm 2\sigma$) confidence interval. The red line is the expected Poisson distribution 1.1. The considered system is the rectangle mascara.

As expected, we can clearly see in Figure 2 that the spacing distribution for the integrable rectangle follows a Poisson distribution. This graphic shows how this




Figure 3: Blue bars represent the histogram spacings, with associated errorbars that follows a counting Poisson error in a 95 % ($\pm 2\sigma$) confidence interval. The red line is the distribution that fits the better the tendency of the histogram. The considered system is the complete billiard.




Figure 4: Blue bars represent the histogram spacings, with associated errorbars that follows a counting Poisson error in a 95 % ($\pm 2\sigma$) confidence interval. The green line is the expected WD distribution 1.2 for a non-integrable system. The considered system is a quarter of the billiard.

systems, even if it is integrable, that their eigenvalues are not correlated, then the probability of finding two eigenvalues close to each other is high ($P(s) \rightarrow 1$ when $s \rightarrow 0$).

Also, in the case of the non-integrable billiard (the complete stadium showed in Figure 3), we can see that it also follows a near Poisson distribution, due to the symmetries of the system. This symmetries make the system to have four pairs of solutions, then, the degenerate levels will increase the probability of finding two eigenvalues close to each other, even if the system is non-integrable.

On the other hand, if we eliminate the x – axis and y – axis symmetries, having just a quarter of the stadium as we see in [Figure 4](#), it is noticeable that the spacing distribution for the non-integrable billiard follows a Wigner-Dyson distribution. In this case, the opposite happens: the system may seem chaotic, but its eigenvalues are correlated, in a way that makes them repel each other or, in other words, the probability of finding two near-degenerate eigenvalues is very low ($P(s) \rightarrow 0$ when $s \rightarrow 0$). [\[4\]](#)

3.2 High-energy states visualization

To determine the chaocity of the system, we must look at high-energy states, and compare them to the non-chaotic system. This is due to the fact that, in the low-energy states, the eigenvectors may seem like those of a string, therefore, they don't contain the necessary information to see if the system is chaotic or not. We then plot the probability density of some of the high-energy states for both systems.

The three graphics of [Figure 5](#) show the probability density for three of the high-energy states of the integrable rectangle. We see that the patterns are very regular, and they look like the fundamental modes of a string, with nodes and antinodes. We can determine then that this system is not chaotic, as expected, since there are periodic patterns in the probability density.


In contrast, the same eigenvalues are shown for the non-integrable quarter billiard in [Figure 6](#). In this system, the probability density is not random or uniformly distributed, but it shows some patterns that are not regular, and they don't look like the fundamental modes of a string.

This is a striking result: in a classically chaotic system, one would expect an erratic way of filling the space, without any specific path or visual pattern. Yet, as seen in [Figure 6](#), certain high-energy eigenstates show a clear concentration of probability density along specific curves in this quarter of the billiard.


If we reverse the same reasoning that we stated to divide the complete billiard in just a quarter to eliminate the symmetries, now we can take advantage of this symmetries to visualize better this trajectory along the billiard, since it will be a closed trajectory and, therefore, it's directly comparable to a classical orbit.

Now, in [Figure 7](#) we can see for those cases that this closed trajectory resemble of a classical one. This is what is known as '**Quantum Scars**', and they correspond to the unstable periodic orbits of the classical system. The origin of quantum scars lies in wave interference: the quantum wavefunction can constructively interfere along the path of a periodic orbit, even if that orbit is unstable (which is the case, because in this system, another initial condition will make that the final result is totally different).


It's important to note that we'd only uploaded the files that we'd see representative. Any other state can be checked in the provided code in the github. [\[1\]](#)



(a) Probability density for the eigenvalue 1000.



(b) Probability density for the eigenvalue 1500.




(c) Probability density for the eigenvalue 1800.


Figure 5: Visualization of various of the high-energy states for the **integrable rectangle**. In the left, it shows the heatmap of the probability density for the eigenvalues 1000, 1500, and 1800; in the right, the same is shown, but in 3D, having axis x and y , and the third dimension represents the probability density.

3.2.1 Ergodicity in the quantum system


A system is said to be ergodic when, after evolving for a sufficiently long time, a single trajectory visits every region of the accesible phase space. According to Shnirelman's theorem (also known as the quantum ergodicity theorem), this classical ergodicity is reflected in the corresponding quantum system: for almost all high-energy eigenstates, the probability density becomes equidistributed over the spatial domain in the semiclassical limit.



(a) Probability density for the eigenvalue 1000.



(b) Probability density for the eigenvalue 1500.




(c) Probability density for the eigenvalue 1800.


Figura 6: Visualization of various of the high-energy states for the **non-integrable quarter billiard**. In the left, it shows the heatmap of the probability density for the eigenvalues 1000, 1500, and 1800; in the right, the same is shown, but in 3D, having axis x and y , and the third dimension represents the probability density.

This theorem is showed in our figures: the rectangular system shows regions with zero probability of the particle to be in that position, as we can see at any example of the [Figure 5](#). Meanwhile, the quantum system has a non zero probability in every point of the space.


If we imagine the system as a ball hitting the edges of the surface and bouncing during a long time, the ball will never cross certain areas in the case of the rectangle, but, in the billiard, eventually it will have crossed the entire surface at least one time.



(a) Probability density for the eigenvalue 1200.



(b) Probability density for the eigenvalue 1400.



(c) Probability density for the eigenvalue 1800.

Figure 7: Visualization of various of the high-energy states for the **non-integrable complex billiard**. In the left, it shows the heatmap of the probability density for the eigenvalues 1200, 1400, and 1800 of the billiard. In the right, the same is shown, but in 3D, having axis x and y , and the third dimension represents the probability density.

At last, it's interesting to discern the quantum scars from the ergodicity theorem. Although Shnirelman's theorem predicts that the probability density of high-energy eigenstates in a chaotic system should eventually become equidistributed, the presence of quantum scars introduce a nuance to this ergodicity.

This phenomenon is explained with the Gutzwiller's Trace Formula, which relates the quantum density of states to a summation over all classical periodic orbits: eigenvalues emerge as singularities where wave phases interfere, as we said, constructively along these paths. Consequently, while the system tends toward ergodicity in the semiclassical limit, these scars represent a visual suppression of classical chaos. [5]

4 Conclusions

The finite difference method combined with sparse matrices to perform the diagonalization results in a powerful technique for solving the 2D Schrödinger equation in arbitrary geometries. Using this approach, we obtain the eigenvalue spacings and observe signatures of chaotic behavior.

We show that the level spacing distribution in the rectangular billiard is Poissonian, which can be interpreted as the eigenvalues being uncorrelated. As a consequence, there is a high probability of finding two eigenvalues very close to each other or even degenerate. On the other hand, the spacing distribution in the stadium billiard follows Wigner-Dyson statistics, which indicates that the eigenvalues are correlated. Therefore, the probability of finding two very close or degenerate eigenvalues is strongly suppressed.

It is important to note that the complete stadium billiard possesses spatial symmetries (even and odd parity), causing the spectrum to become a superposition of independent subsystems. By considering the quarter-stadium geometry, these symmetry sectors are separated, allowing the pure Wigner-Dyson statistics to emerge without contamination from states of different parity. This isolation of the spatial symmetries is essential for obtaining undistorted spectral statistics that properly reflect the chaotic nature of the system.

In summary, this work successfully demonstrates how the complex statistical properties predicted by Random Matrix Theory naturally emerge when numerically solving a simple physical system under specific geometric boundary conditions.

Referencias

- [1] María de los Ángeles Lara and Pablo Orellana. PCS-Project. <https://github.com/l3l3ss/PCS-Project->, 2026. Accessed: 2026-05-01.
- [2] Diego Ariel Wisniacki. *Caos cuántico en sistemas dependientes de un parámetro*. PhD thesis, Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales, 1999.
- [3] Antonio Carrillo Ledesma, Karla Ivonne González Rosas, and Omar Mendoza Bernal. Introducción al método de diferencias finitas y su implementación computacional. *Bernal, Facultad de Ciencias, UNAM*, 2015.
- [4] Hiroki Nakayama. Random Matrix Theory, Quantum Chaos, and Eigenstate Thermalization, July 2020. Submitted as coursework for PH470, Stanford University, Spring 2020. <http://large.stanford.edu/courses/2020/ph470/nakayama/docs/nakayama-ph470-stanford-spring-2020.pdf>.
- [5] Wikipedia contributors. Quantum scar. https://en.wikipedia.org/wiki/Quantum_scar, 2026. Wikipedia, The Free Encyclopedia. Accessed: 2026-06-01.