

AAAFM UCLA
American Association for Advances in
Functional Materials
Materials For Life
2021
18-20 August

Ultrasound as Functional Influence Tool on FeB Pair Association in Silicon Solar Cells

Oleg Olikh¹, Vitaliy Kostylyov²,
Victor Vlasiuk², Roman Korkishko²



**¹Taras Shevchenko National University
of Kyiv, Kyiv, Ukraine**



**²V.Lashkaryov Institute of
Semiconductor Physic, Kyiv, Ukraine**



НАН УКРАЇНИ

ІФН


ISP

NAS UKRAINE



Main goal:

to investigate experimentally the FeB pair association in silicon solar cells under ultrasound loading conditions






НАН УКРАЇНИ



IФН

ISP

NAS UKRAINE



Samples:



- 1 - front metal electrode (Al);
- 2 - Si_3N_4 (40 nm);
- 3 - SiO_2 (30 nm);
- 4 - induced n++ - layer;
- 5 - diffusion n+ - layer (0.7 μm);
- 6 - base region (p-Si:B, 10 $\Omega\cdot\text{cm}$, 380 μm);
- 7 - diffusion p+ -layer (0.6 μm);
- 8 - rear metallization (Al)




НАН УКРАЇНИ


IФН

ISP

NAS UKRAINE




Experimental setup:





Ultrasound loading (USL)



- 1 – Cu foil
- 2 – piezoelectric transducer
- 3 – contacts for IV measuring
- 4 – contacts for US excitation

1) longitudinal wave (thickness vibrations)

$$f_{\text{US}} = 2.4; 4.1; 5.4; 9.0; 14; 18; 31 \text{ MHz}$$

2) transverse wave (radial vibrations)

$$f_{\text{US}} = 0.3 \text{ MHz}$$

ultrasound intensity $W_{\text{US}} \leq 0.6 \text{ W/cm}^2$



UCLA 2021 817. Ultrasound as Functional Influence Tool on FeB Pair Association in Silicon Solar Cells

$$I_{SC} = \frac{W_{ph}(1-R)q\beta\lambda}{hc} \frac{\alpha\sqrt{D\tau}}{1+\alpha\sqrt{D\tau}}$$

$$\tau^{-1} = \tau_{rad}^{-1} + \tau_{Aug}^{-1} + (\tau_{SRH}^{Fe_i})^{-1} + (\tau_{SRH}^{FeB})^{-1} + \tau_{other}^{-1}$$

$$N_{Fe}(t) = (N_{Fe,0} - N_{Fe,eq}) \exp\left(-\frac{t}{\tau_{ass}}\right) + N_{Fe,eq}$$

$$\tau_{SRH}^{Fe_i,FeB} = \frac{\tau_{p0}(n_0 + n_1) + \tau_{n0}(N_A + p_1)}{N_A + n_0}$$


$$N_{Fe,eq} = \frac{N_{Fe,0}}{\left[1 + N_A 10^{-23} \exp\left(\frac{0.582}{kT}\right)\right] \left[1 + \exp\left(\frac{E_F - 0.394}{kT}\right)\right]}$$

$$\tau_{n0,p0} = (N_{trap} \sigma_{n,p} v_{th,n,p})^{-1}$$

calculated values

experiment parameters

extracted parameters



points – experiment,
lines - fitting



НАН УКРАЇНИ


ІФН

ISP

NAS UKRAINE




Results





Results





НАН УКРАЇНИ




IФН


ISP

NAS UKRAINE




Results





Results





НАН УКРАЇНИ


ІФН


ISP

NAS UKRAINE



Results






Discussion




$$\tau_{ass} = \frac{\varepsilon \varepsilon_0 k_B T}{q^2 D_{Fe} N_A} = \frac{\varepsilon \varepsilon_0 k_B T}{q^2 D_{0,Fe} N_A} \exp\left(\frac{E_{m,Fe_i}}{k_B T}\right)$$

iron atom migration energy



$$\Delta E_{US} \leq 10 \text{ meV}$$





CONCLUSION

- ✓ The effect of the acoustically induced acceleration of FeB pairing in silicon solar cell has revealed
- ✓ The efficiency of ultrasound influence decreases with the acoustic waves frequency rising and the temperature drop and does not depend on iron concentration. The effect saturation with ultrasound intensity increase is observed
- ✓ The analysis has shown that these effects are caused by the acoustically induced decrease in the migration energy of iron atom




НАН УКРАЇНИ

ІФН

ISP

NAS UKRAINE



olegolikh@knu.ua

**THANK YOU
FOR YOUR
ATTENTION**

