

Injection dependence of spontaneous radiative recombination in crystalline silicon: Experimental verification and theoretical analysis

P. P. Altermatt^{a)}

Department of Solar Energy Research, Institute of Solid-State Physics, University of Hannover, Appelstrasse 2, 30167 Hannover, Germany

F. Geelhaar^{b)}

Integrated Systems Laboratory, Swiss Federal Institute of Technology, Gloriastrasse 35, 8092 Zurich, Switzerland

T. Trupke, X. Dai, and A. Neisser^{c)}

Centre of Excellence for Advanced Silicon Photovoltaics and Photonics, University of New South Wales, Sydney, New South Wales 2052, Australia

E. Daub

Siltronic AG, Johannes-Hess-Strasse 24, 84489 Burghausen, Germany

(Received 17 February 2006; accepted 12 May 2006; published online 26 June 2006)

The radiative recombination coefficient B in crystalline bulk silicon is enhanced by the Coulomb attraction between electrons and holes. This effect is weakened at high carrier densities due to screening. We measure the resulting dependence of B on the free-carrier density (i) by reinterpreting published data and (ii) with photoluminescence and photovoltaic measurements. We calculate the Coulomb enhancement by determining the electron-hole pair correlation function at zero interparticle distance, assuming a Debye interaction potential. Both bound and scattering state contributions are fully taken into account. Due to screening, B decreases with increasing free-carrier density. © 2006 American Institute of Physics. [DOI: 10.1063/1.2218041]

The quantification of the radiative recombination rate in crystalline silicon is important to the present development of efficient light-emitting devices based on this material. Such devices have reached external quantum efficiencies of near 1% for electroluminescence¹ and over 10% for photoluminescence (PL).² This is high considering that silicon has an indirect band gap and also influences the performance of solar cells that operate under concentrated sunlight. The rate R of spontaneous radiative recombination is proportional to the density of free electrons n and free holes p . Considering only excess carriers, there holds

$$R = B(np - n_{i,\text{eff}}^2), \quad (1)$$

where B is the radiative recombination coefficient and $n_{i,\text{eff}}$ is the effective intrinsic carrier density. It has been known since 1974 that B is enhanced by the Coulomb attraction between electrons and holes,³ which leads to an increased hole density in the vicinity of an electron, and vice versa. This enhancement depends on the amount of screening among the free carriers: if there is a sufficient density of free electrons between the mentioned electron and its surrounding holes, their Coulomb attraction is screened. Screening depends on both temperature³ and carrier injection level.⁴ The temperature dependence of B has been thoroughly examined,^{3,5-7} while no quantification of the injection dependence is currently available, although it is the dominant effect in light-emitting devices operated at constant ambient temperature.

In this letter, we present both measurements and theoretical calculations of the injection dependence.

B can be determined experimentally using two different approaches. First, by measuring R and the density of excess carriers;³ second, by measuring the absorption coefficient for band-to-band transitions α_{bb} and by applying the van Roosbroeck theory⁸ or the generalized Planck equation,^{7,9}

$$B = \frac{1}{n_{i,\text{eff}}^2 \pi^2 \hbar^3 c_0^2} \int_0^\infty N^2 E_{\text{ph}}^2 \alpha_{\text{bb}} e^{-E_{\text{ph}}/kT} dE_{\text{ph}}. \quad (2)$$

This formula involves the index of refraction N , the speed of light in vacuum c_0 , and the photon energy E_{ph} .

We scale B to B/B_{low} , where B_{low} is obtained at very low carrier densities. We do this because, in contrast to measurements of B , B/B_{low} does not require absolute measurements of the emitted photon flux j_γ ; neither are light-trapping and reabsorption processes nor the exact value of $n_{i,\text{eff}}$ involved. Furthermore, calculations relative to B_{low} do not depend on small variations in the effective masses or in the static dielectric constant, so that our values for B/B_{low} will be unaffected by possible future refinements of silicon parameters.

In the symmetrical plasma ($n=p$), we determine the injection dependence of B with the first experimental approach, using the well-established data of Schlangerotto *et al.*³ While Eq. (1) predicts that j_γ is proportional to np , the measurements depicted in Fig. 1 of Ref. 3 deviate from this simple behavior, which we plot in our Fig. 1 in terms of B/B_{low} (symbols). The data scatter considerably above $B/B_{\text{low}}=0.8$, partly because our digital scan of the plot in Ref. 3 has a limited precision, and partly due to the restricted accuracy of the original experiments. At lower values of B/B_{low} , the data show that B declines with increasing injection density in a temperature-dependent manner.

^{a)}Electronic mail: altermatt@solar.uni-hannover.de

^{b)}Present address: Advanced Micro Devices, Inc., One AMD Place, M/S 143, Sunnyvale, CA 94088.

^{c)}Present address: Hahn-Meitner Institute, Glienicke Strasse 100, 14109 Berlin, Germany.




FIG. 1. The scaled radiative recombination coefficient B/B_{low} for a symmetrical plasma ($n=p$) at various temperatures. B_{low} is obtained at very low carrier densities. Symbols: measurements, as extracted from Ref. 3; lines: our calculations.

In the asymmetrical case ($n \neq p$), we determine B with the second experimental approach, where α_{bb} is required for various phosphorus or boron dopant densities N_{dop} . At high N_{dop} , spectroscopic transmission measurements do not render α_{bb} , but rather the sum of α_{bb} and the coefficient for free-carrier absorption α_{fc} ; the latter may completely dominate at the photon energies E_{ph} of interest. To circumvent this obstacle, we obtain α_{bb} from PL spectra in the same way as in Ref. 10, i.e., from relative measurements of the photon flux per energy interval and by applying⁷

$$\alpha_{\text{bb}} \propto \frac{dj_{\gamma}}{dE_{\text{ph}}} E_{\text{ph}}^{-2} e^{E_{\text{ph}}/kT}. \quad (3)$$

As it is difficult to measure dj_{γ} in absolute units, we scale dj_{γ} such that the resulting α_{bb} matches well-established data from spectroscopic transmission measurements¹¹ in a suitable photon energy range. In order to calculate B from these absorption data, we apply Eq. (2) involving $n_{i,\text{eff}}$ and N . We obtain $n_{i,\text{eff}}$ via

$$n_{i,\text{eff}} = n_i e^{\Delta E_g/(2kT)}, \quad (4)$$

where the intrinsic carrier density n_i is taken from Ref. 12 and the band gap shrinkage ΔE_g from Ref. 13. Finally, we use N given in Ref. 14 for lowly doped silicon because the changes in N due to doping are negligibly small.¹⁵ The resulting B and B/B_{low} values are depicted in Fig. 2; they decline with increasing hole or electron density.

To complement these data, we fabricated solar cells with wafers of various boron dopant densities,¹⁶ and we determined α_{bb} with spectral response measurements as in Refs. 17 and 18. As the cells are only 250 μm thin, free-carrier absorption attenuates the incoming light only at very high N_{dop} , where we correct the data with ray-tracing techniques (the optical properties of the cells are known very




FIG. 2. The radiative recombination coefficient B for the asymmetrical plasma ($n \neq p$) at 300 K, scaled to B/B_{low} . Symbols: our measurements obtained using photovoltaic (PV) and photoluminescence (PL) methods; lines: our calculations.

precisely). Because individual devices of various dopant densities are used, B/B_{low} cannot be determined directly. Instead, we calculate B from the absorption data in absolute terms via Eq. (2) and scale it to B/B_{low} , which is independent of the choice of n_i and N . Our results are plotted in Fig. 2 as circles; they coincide with the PL data shown as squares.

In the following, we outline our calculations of the injection dependence of B , shown as lines in Figs. 1 and 2. The radiative recombination is influenced by the Coulomb attraction between the electrons and holes, which increases the band-to-band recombination coefficient from its hypothetical free-carrier value B_0 to the empirical value $B = g_{eh} B_0$. The factor g_{eh} accounts for the enhanced probability of finding an electron and a hole in the immediate vicinity of each other and is given by the pair correlation function at zero interparticle distance. In contrast to Schlagenotto *et al.*,³ who described the electron-hole interaction in terms of the bare Coulomb potential, we incorporate screening effects by employing the Debye potential. This is given by $w_{eh}(r) = -e^2/(4\pi\epsilon_r\epsilon_0)\exp(-\kappa r)/r$, where ϵ_r and ϵ_0 are the relative and vacuum dielectric constants, respectively, r is the distance and κ is the inverse screening length. A Padé approximation¹⁹ has been used to obtain κ as a function of the carrier densities n and p . Denoting the radial s -wave functions for the relative motion of the two charge carriers by $R_{\nu 0}(r)$, there holds

$$g_{eh} = \frac{\Lambda_{eh}^3}{4\pi} \sum_{\nu} |R_{\nu 0}(0)|^2 \exp(-E_{\nu 0}/kT), \quad (5)$$

where $\Lambda_{eh}^2 = 2\pi\hbar^2/kTm_{eh}$ is the thermal wavelength squared and m_{eh} is the reduced electron-hole mass. The summation index ν comprises both discrete and continuous quantum numbers. The energy levels $E_{\nu 0}$ and wave functions $R_{\nu 0}(r)$ in Eq. (5) vary with κ , so that g_{eh} depends not only on the temperature T but also on the particle densities. This is a direct consequence of the screening effect. In particular, when Boltzmann statistics can be applied, one has $\kappa^2 = (e^2/kT\epsilon_r\epsilon_0)(n+p)$, and the enhancement factor becomes a function of $n+p$. The bound-state portion of g_{eh} has been computed from a variational solution of the radial

Schrödinger equation by means of the quadrature discretization method;^{20–22} the details of our procedure are given in Ref. 23. The scattering state portion of g_{eh} has been derived from the Noyes-Kowalski half-shell functions^{24,25} for the two-particle T matrix, as outlined in Refs. 26 and 27. For low carrier densities ($\kappa \rightarrow 0$), our numerical results correctly reproduce the analytical results for the bare Coulomb potential given in Ref. 28. For high carrier densities ($\kappa \rightarrow \infty$), g_{eh} decreases to unity because the screening is then so strong that the electron-hole system can be described by the Hartree-Fock approximation. From Eq. (5) and the Boltzmann expression for κ given above, it can be seen that the decrease of g_{eh} sets in at lower carrier densities when the temperature is reduced.

The theoretical curves describe the experiments well within the measurement's precision, as shown in Figs. 1 and 2. It has to be kept in mind that the experimental values in the asymmetrical plasma were obtained with either $n \gg p$ or $n \ll p$, while the calculations were performed for the symmetrical plasma ($n=p$). We can compare the two because at carrier densities below around $2 \times 10^{19} \text{ cm}^{-3}$, Boltzmann statistics are valid, and g_{eh} is a function of $n+p$.

In order to obtain the absolute values of B , B/B_{low} needs to be multiplied with the experimentally determined B_{low} values published in Ref. 7.

In conclusion, we experimentally confirmed that the radiative recombination coefficient B in crystalline bulk silicon depends on the injection density of free carriers, and we explained our measurements in terms of numerical results for the electron-hole pair correlation function using Debye screening. We showed that B/B_{low} (with B_{low} obtained at very low carrier densities) decreases with increasing carrier densities and that this decrease starts at lower carrier densities as the temperature is reduced.

Part of this work was conducted while one of the authors (P.P.A.) worked at the Australian National University, Canberra, Australia. The author is also grateful for discussions with Andres Cuevas and Daniel Macdonald from the ANU.

Another author (F.G.) would like to thank the Swiss National Science Foundation for financial support under the research Project No. 21-58811.99. Another author (T.T.) is grateful for financial support from the Alexander von Humboldt foundation.

- ¹M. A. Green, J. Zhao, A. Wang, P. J. Reece, and M. Gal, *Nature* (London) **412**, 805 (2001).
- ²T. Trupke, J. Zhao, A. Wang, R. Corkish, and M. A. Green, *Appl. Phys. Lett.* **82**, 2996 (2003).
- ³H. Schlagenotto, H. Maeder, and W. Gerlach, *Phys. Status Solidi A* **21**, 357 (1974).
- ⁴A. Hangleiter, thesis, University of Stuttgart, 1992.
- ⁵Y. P. Varshni, *Phys. Status Solidi* **19**, 459 (1967).
- ⁶W. Michaelis and M. H. Pilkuhn, *Phys. Status Solidi* **36**, 311 (1969).
- ⁷T. Trupke, M. A. Green, P. Würfel, P. P. Altermatt, A. Wang, J. Zhao, and R. Corkish, *J. Appl. Phys.* **94**, 4930 (2003).
- ⁸W. van Roosbroeck and W. Shockley, *Phys. Rev.* **94**, 1558 (1954).
- ⁹P. Würfel, *J. Phys. C* **15**, 3967 (1982).
- ¹⁰E. Daub and P. Würfel, *Phys. Rev. Lett.* **74**, 1020 (1995).
- ¹¹G. G. Macfarlane, T. P. McLean, J. E. Quarrington, and V. Roberts, *Phys. Rev.* **111**, 1245 (1958).
- ¹²P. P. Altermatt, A. Schenk, F. Geelhaar, and G. Heiser, *J. Appl. Phys.* **93**, 1598 (2002).
- ¹³A. Schenk, *J. Appl. Phys.* **84**, 3684 (1998).
- ¹⁴M. A. Green and M. J. Keevers, *Prog. Photovoltaics* **3**, 189 (1995).
- ¹⁵H. C. Huang, S. Yee, and M. Soma, *J. Appl. Phys.* **67**, 2033 (1990).
- ¹⁶X. Dai, Ph.D. thesis, University of New South Wales, 1994.
- ¹⁷M. J. Keevers and M. A. Green, *Appl. Phys. Lett.* **66**, 174 (1995).
- ¹⁸A. Neisser and M. A. Green, *Proceedings of the 2nd World Conference on Photovoltaic Solar Energy Conversion* (James&James, London, 1998), pp. 136–139.
- ¹⁹R. Zimmermann, *Many-Particle Theory of Highly Excited Semiconductors* (BSB Teubner, Leipzig, 1988).
- ²⁰F. Calogero, *Lett. Nuovo Cimento Soc. Ital. Fis.* **37**, 9 (1983).
- ²¹L. Durand, *Polynômes Orthogonaux et Applications* (Springer-Verlag, New York, 1985), Vol. 1171.
- ²²C. Mavriplis, *J. Chem. Phys.* **80**, 480 (1989).
- ²³F. Geelhaar, *Coulomb Correlation Effects in Silicon Devices*, Series in Microelectronics Vol. 147 (Hartung-Gorre, Konstanz, 2004).
- ²⁴H. P. Noyes, *Phys. Rev. Lett.* **15**, 538 (1965).
- ²⁵K. L. Kowalski, *Phys. Rev. Lett.* **15**, 798 (1965); **15**, 908 (1965).
- ²⁶G. D. Mahan, *Phys. Rev.* **153**, 882 (1966).
- ²⁷S. Schmitt-Rink, C. Ell, and H. Haug, *Phys. Rev. B* **33**, 1183 (1986).
- ²⁸B. Davies and R. G. Storer, *Phys. Rev.* **171**, 150 (1968).