



XXII International Seminar on Physics and Chemistry of Solids

Book of Abstracts



**June 17-19, 2020
Lviv, Ukraine**



Lviv City Council
Lviv Convention Bureau
Ivan Franko National University of Lviv
Jan Dlugosz University in Czestochowa

XXII International Seminar on Physics and Chemistry of Solids

Book of Abstracts

June 17-19, 2020
Lviv, Ukraine

SCIENTIFIC COMMITTEE

Zygmunt Bąk
Wojciech Ciesielski
Grygoriy Dmytriv
Jozef Drabowicz
Jacek Filipecki
Roman Gladyshevskii (chairman)
Janusz Kapuśniak (co-chairman)
Volodymyr Kapustianyk
Małgorzata Makowska-Janusik
Stepan Mudry
Vasyl Stadnyk
Volodymyr Tkachuk
Anatolii Voloshinovskii
Petro Yakibchuk

ORGANIZING COMMITTEE

Oleg Bovgyra
Yaroslav Chornodolskyy
Andryj Korolyshyn
Stepan Mudry (chairman)
Volodymyr Pavlyuk
Yuriy Plevachuk
Viktor Prysyazhnyuk
Ihor Shtablavyi
Vasyl Sklyarchuk
Anatolii Voloshinovskii (co-chairman)
Ivan Shcherba

SEMINAR SECRETARY

Ihor Shtablavyi
Faculty of Physics
Ivan Franko National University of Lviv
Kyryla and Mefodia Str., 8a, Lviv, 79005, Ukraine.
Phone: + 38-032-239-45-94
Phone: + 38-067-451-59-52
e-mail: ispcs2020@gmail.com

MODELING OF IDEALITY FACTOR VALUE IN SILICON SOLAR CELLS

Olikh O.Ya., Zavhorodnii O. V.

¹ Faculty of Physics, Taras Shevchenko National University of Kyiv, Kyiv 01601, Ukraine
olikh@univ.kiev.ua

Ideality factor (n) is one of a commonly used parameter of solar cell (SC). In the simplest case, $n = 2$ is used for Shockley-Read-Hall recombination current. But in fact, n depends on ambient conditions and recombination center parameters. The purpose of this work is to evident such dependencies for Si-SC.

The current-voltage characteristics of silicon n^+p-p^+ structure were calculated by using of one-dimensional code SCAPS in the temperature range 290-340 K. The both base depth d and acceptor (boron) concentration N_A were varied over 150-240 μm and $10^{15}\div 10^{17} \text{ cm}^{-3}$ respectively. The iron atoms are suggested to be present in p-layers with $N_{\text{Fe}} = 10^{10}\div 10^{13} \text{ cm}^{-3}$. The case of Fe_iB_s pairs as well as the case of interstitial Fe_i was under consideration. The ideality factor was determined according to the two-diode model. Some results are presented in Fig.1.




Figure 1: Dependencies of ideality factor in equilibrium state (n , upper surfaces) and alteration in ideality factor after Fe-B pair dissociation (δn , lower surfaces).
 $N_{\text{Fe}} = 10^{10} \text{ cm}^{-3}$, $d = 240 \mu\text{m}$ (a), $N_A = 10^{17} \text{ cm}^{-3}$, $T = 340 \text{ K}$ (b).

It was established that i) the base depth affects the ideality factor value in the case of minority carrier diffusion length $L_n \gg d$; ii) the influence of temperature and doping level deals with change in a recombination level population mainly; iii) the dependence of n on the iron concentration is a monotonic function.